

TOPOLOGICAL FLATNESS OF LOCAL MODELS FOR RAMIFIED UNITARY GROUPS. II. THE EVEN DIMENSIONAL CASE

BRIAN D. SMITHLING

ABSTRACT. Local models are schemes, defined in terms of linear-algebraic moduli problems, which give étale-local neighborhoods of integral models of certain p -adic PEL Shimura varieties defined by Rapoport and Zink. In the case of a unitary similitude group whose localization at $\mathbb{Q}_p$ is ramified, quasi-split GU_n , Pappas has observed that the original local models are typically not flat, and he and Rapoport have introduced new conditions to the original moduli problem which they conjecture to yield a flat scheme. In a previous paper we proved that their new local models are topologically flat when n is odd. In the present paper we prove topological flatness when n is even. Along the way, we characterize the μ -admissible set for certain cocharacters μ in types B and D , and we show that for these cocharacters admissibility can be characterized in a vertexwise way, confirming a conjecture of Pappas and Rapoport.

CONTENTS

1. Introduction	3
Acknowledgments	7
Notation	7
2. Unitary local models	8
2.1. Pairings	8
2.2. Standard lattices	9
2.3. Naïve local models	9
2.4. Wedge and spin conditions	10
2.5. Lattice chain automorphisms	12
3. Unitary similitude group	12
3.1. Unitary similitudes	12
3.2. Tori	12
3.3. Kottwitz homomorphism	13
3.4. Parahoric subgroups	13
3.5. Affine roots	13
4. Preliminary combinatorics	14
4.1. The group $\tilde{W}_n$	14
4.2. Faces of type (n, I)	14
4.3. The vector $\mu_i^{\mathbf{v}}$	15
4.4. Further lemmas	16
5. Iwahori-Weyl groups	17

2010 *Mathematics Subject Classification*. Primary 14G35; Secondary 05E15; 11G18; 17B22.

Key words and phrases. Local model; unitary group; Iwahori-Weyl group, admissible set.

5.1.	Iwahori-Weyl groups	17
5.2.	Bruhat order	18
5.3.	Admissible and permissible sets	18
5.4.	Case of GU_{2m}	19
5.5.	Faces of type I	20
5.6.	Kottwitz homomorphism on $\widetilde{W}_G$	21
5.7.	Relation to type B_m	21
5.8.	The $\{\mu_{r,s}\}$ -admissible set	23
6.	Affine flag variety	23
6.1.	Affine flag variety	23
6.2.	Lattice-theoretic description	24
6.3.	Schubert cells and varieties	25
6.4.	Embedding the geometric special fiber	26
7.	Schubert cells in the geometric special fiber	27
7.1.	The image of the geometric special fiber	27
7.2.	Naive permissibility	27
7.3.	Wedge-permissibility	28
7.4.	Spin-permissibility	29
7.5.	Topological flatness of M_I^{spin}	33
8.	Type D combinatorics	33
8.1.	The root datum $\mathcal{R}_{D_m}$	34
8.2.	μ -spin-permissibility	35
8.3.	Proper elements	36
8.4.	The integer c_k^w	36
8.5.	The vector ν_k^w	37
8.6.	μ -permissibility	39
8.7.	Intervals	42
8.8.	Reflections	43
8.9.	Reflections and the Bruhat order	44
8.10.	μ -admissibility implies μ -spin-permissibility	44
8.11.	Lemmas on reflections and μ -spin-permissibility	45
8.12.	μ -spin-permissibility implies μ -admissibility	48
8.13.	Parahoric variants and faces of type I	60
8.14.	μ -admissibility and μ -spin-permissibility in the parahoric case	62
8.15.	Vertexwise admissibility	67
8.16.	μ -permissibility in the parahoric case	70
9.	Type B combinatorics	70
9.1.	Steinberg fixed-point root data	71
9.2.	The root datum $\mathcal{R}_{B_m}$	73
9.3.	Relation to $\mathcal{R}_{D_{m+1}}$	75
9.4.	Parahoric variants	75
9.5.	μ -spin-permissibility	77
9.6.	μ -spin-permissibility and μ -admissibility	79
9.7.	Vertexwise admissibility	80
9.8.	μ -permissibility	81
	References	82

1. INTRODUCTION

One of the basic problems in the arithmetic theory of Shimura varieties is the construction of reasonable integral models over the ring of integers $\mathcal{O}$ of the reflex field, or over certain localizations of $\mathcal{O}$. One would like to construct models that are faithfully flat with mild singularities, and in cases in which the Shimura variety can be described as a moduli space of abelian varieties with additional structure, to describe the model in terms of an integral version of the moduli problem. A major intended application is the calculation of the Hasse-Weil zeta function of the Shimura variety.

Let $(\mathbf{G}, X)$ be a Shimura datum, and let $K \subset \mathbf{G}(\mathbb{A}_f)$ be a sufficiently small compact open subgroup of the form $K = K^p K_p$, where $K_p \subset \mathbf{G}(\mathbb{Q}_p)$, $K^p \subset \mathbf{G}(\mathbb{A}_f^p)$, and $\mathbb{A}_f^p$ denotes the ring of finite adeles over $\mathbb{Q}$ with trivial p -component. Let $\mathbf{E}$ denote the reflex field, let $\mathrm{Sh}_K(\mathbf{G}, X)$ denote the canonical model of the Shimura variety over $\mathbf{E}$, let v be a place of $\mathbf{E}$ over p , and let $\mathcal{O}_{\mathbf{E}_v}$ denote the ring of integers in the v -adic completion $\mathbf{E}_v$. When K_p is a *hyperspecial* subgroup of $\mathbf{G}(\mathbb{Q}_p)$, smooth $\mathcal{O}_{\mathbf{E}_v}$ -models of $\mathrm{Sh}_K(\mathbf{G}, X)$ were constructed in most cases of PEL type by Kottwitz [Ko1], and, more generally, in cases of abelian type by Kisin [Ki] and Vasiu (see [V] and the references therein).

Our concern in this paper is the more general setting that K_p is a *parahoric*, but not necessarily hyperspecial, subgroup. Then it is no longer reasonable to expect a smooth integral model. For PEL Shimura varieties, when K_p is a parahoric subgroup that can be described as the stabilizer of a lattice chain, Rapoport and Zink constructed natural integral models over $\mathcal{O}_{\mathbf{E}_v}$ in [RZ]. When the localization $\mathbf{G}_{\mathbb{Q}_p}$ is unramified with simple factors of types only A and C , Görtz [G1, G2] showed that Rapoport and Zink's models are flat with reduced special fiber, and that the irreducible components of the special fiber are normal with rational singularities. By contrast, when $\mathbf{G}_{\mathbb{Q}_p}$ is *ramified*, Pappas [P] observed that the Rapoport–Zink models may fail to be flat.

The key tool to investigate flatness of the Rapoport–Zink models, as well as other questions of a local nature, is the *naive local model*, also introduced by Rapoport and Zink in [RZ]. The naive local model gives an étale-local neighborhood of the Rapoport–Zink integral Shimura model, but it is defined in terms of a purely linear-algebraic moduli problem, whereas the Shimura model is defined in terms of abelian varieties. Therefore one expects the naive local model to be easier to study. Here “naive” signifies that the naive local model, like the integral Shimura model, may fail to be flat.

In light of Pappas's observation, it is necessary to correct the definition of the integral Shimura models — or, equivalently, of the naive local models — to obtain flat schemes over $\mathcal{O}_{\mathbf{E}_v}$. The most straightforward solution is to define the corrected models and local models to be the scheme-theoretic closure of the generic fiber in the original models and local models, respectively. Then the corrected models and local models are flat essentially by definition. Zhu's recent proof [Z] of the *coherence conjecture* of Pappas–Rapoport [PR3] has strong implications for the geometry of the corrected models and local models whenever $\mathbf{G}_{\mathbb{Q}_p}$ is tamely ramified; see [PR3, Th. 11.3] for the local models we shall consider in this paper. However, defined this way, the corrected models and local models carry the disadvantage of not admitting ready moduli-theoretic descriptions, since the process of scheme-theoretic closure can be difficult to interpret in a moduli-theoretic way. Thus it is

of interest to describe, when possible, the corrected models and local models via a refinement of the original moduli problem. This is the problem of concern in this paper and its prequel [Sm3], in the case of a unitary similitude group attached to an imaginary quadratic number field ramified at $p \neq 2$.

More precisely, let F/F_0 be a ramified quadratic extension of discretely valued, non-Archimedean fields with perfect residue field of characteristic not 2. Endow F^n , $n \geq 2$, with the F/F_0 -Hermitian form ϕ specified by the values $\phi(e_i, e_j) = \delta_{i, n+1-j}$ on the standard basis vectors $e_1, \dots, e_n$, and consider the reductive group $G := GU_n := GU(\phi)$ over F_0 . In this paper we shall consider exclusively the case that n is even; the case of odd n was treated in [Sm3]. Let $m := n/2$. In the even case, each conjugacy class of parahoric subgroups in $G(F_0)$ contains a unique parahoric of the form P_I° , where I is a nonempty subset of $\{0, 1, \dots, m\}$ with the property

$$(1.1) \quad m+1 \in I \implies m \in I;$$

see §3.4 for definitions, and [PR4, §1.2.3(b)] for more details. We remark that P_I° is the full stabilizer of a lattice chain only when $m \in I$, owing to the existence of elements with nontrivial Kottwitz invariant.

Identifying $G \otimes_{F_0} F \simeq GL_{n,F} \times \mathbb{G}_{m,F}$, let $\mu_{r,s}$ denote the cocharacter $(1^{(s)}, 0^{(r)}, 1)$ of $D \times \mathbb{G}_{m,F}$, where D denotes the standard maximal torus of diagonal matrices in $GL_{n,F}$, and r and s are nonnegative integers with $r+s=n$. Let $\{\mu_{r,s}\}$ denote the geometric conjugacy class of $\mu_{r,s}$. In the special case that $F_0 = \mathbb{Q}_p$ and (F^n, ϕ) is isomorphic to the $\mathbb{Q}_p$ -localization of a Hermitian space (K^n, ψ) with K an imaginary quadratic number field, the pair (r, s) denotes the signature of (K^n, ψ) , and $\{\mu_{r,s}\}$ is the conjugacy class of minuscule cocharacters obtained in the usual way from the Shimura datum attached to the associated unitary similitude group, as in [PR4, §1.1].

Let $E := F_0$ if $r = s$, and $E := F$ if $r \neq s$. Then E is the reflex field of the conjugacy class $\{\mu_{r,s}\}$. Attached to the triple $(G, \{\mu_{r,s}\}, P_I^\circ)$ is the naive local model M_I^{naive} [RZ, PR4], a projective $\mathcal{O}_E$ -scheme whose explicit definition we shall recall in §2.3. As observed by Pappas [P], M_I^{naive} is not flat over $\mathcal{O}_E$ in general. Let M_I^{loc} denote the honest local model, that is, the scheme-theoretic closure of the generic fiber of M_I^{naive} in M_I^{naive} . The papers [P, PR4] propose to describe M_I^{loc} by adding new conditions to the moduli problem defining M_I^{naive} : first Pappas adds the *wedge condition* to define a closed subscheme $M_I^\wedge \subset M_I^{\text{naive}}$, the *wedge local model*; and then Pappas and Rapoport add a further condition, the *spin condition*, to define a third closed subscheme $M_I^{\text{spin}} \subset M_I^\wedge$, the *spin local model* (see §2.4). The schemes

$$M_I^{\text{loc}} \subset M_I^{\text{spin}} \subset M_I^\wedge \subset M_I^{\text{naive}}$$

all have common generic fiber, and Pappas and Rapoport conjecture the following.

Conjecture 1.2 (Pappas–Rapoport [PR4, 7.3]). *M_I^{spin} coincides with the local model M_I^{loc} inside M_I^{naive} , or in other words, M_I^{spin} is flat over $\mathcal{O}_E$.*

Although the full conjecture remains open, the main result of this paper is the following version of it, for even n .

Theorem 1.3. *Suppose that n is even. Then the scheme M_I^{spin} is topologically flat over $\mathcal{O}_E$, or in other words, the underlying topological spaces of M_I^{spin} and M_I^{loc} coincide.*

See §7.5. Recall that a scheme over a regular, integral, 1-dimensional base scheme is *topologically flat* if its generic fiber is dense. We proved the version of Theorem 1.3 for odd n in [Sm3], where we also showed that the wedge local model $M_I^\wedge$ is topologically flat (though typically not flat, as its scheme structure typically differs from that of M_I^{spin}). For even n , we shall see later that $M_I^\wedge$ and M_I^{spin} typically do not even coincide topologically. For example, when $m \in I$, $M_I^\wedge$ and M_I^{spin} coincide topologically only for the trivial signatures $(r, s) = (n, 0)$ and $(r, s) = (0, n)$.

Following Görtz [G1] (see also [G2, G3, PR1, PR2, PR3, PR4, Sm1, Sm3]), the key technique in the proof of Theorem 1.3 is to embed the geometric special fiber of M_I^{naive} in an affine flag variety for GU_n , where it and the geometric special fibers of $M_I^\wedge$, M_I^{spin} , and M_I^{loc} decompose (as topological spaces) into unions of finitely many Schubert varieties; see §6.4. We show the following.

Theorem 1.4. *With respect to the embedding into the affine flag variety specified in §6.4, the Schubert varieties contained in the geometric special fiber of M_I^{spin} are indexed by the $\{\mu_{r,s}\}$ -admissible set.*

See §5.3 for the definition of the $\{\mu_{r,s}\}$ -admissible set, and §7.5 for the proof of the theorem. The $\{\mu_{r,s}\}$ -admissible set was defined by Kottwitz–Rapoport [KR] and Rapoport [R]. It is a subset of a certain double quotient of the *Iwahori–Weyl group* (see §5.1), which indexes the Schubert varieties in the relevant affine flag variety. Pappas and Rapoport show in [PR4, Prop. 3.1] that the $\{\mu_{r,s}\}$ -admissible Schubert varieties are contained in the geometric special fiber of M_I^{loc} , which together with Theorem 1.4 implies Theorem 1.3.

The analog of Theorem 1.4 for odd n is proved in [Sm3]. However, the proof for even n that we shall give here is considerably more laborious. For simplicity, let us explain this in the case that our parahoric subgroup is an Iwahori subgroup. Then the Schubert varieties in the affine flag variety are indexed by the Iwahori–Weyl group $\widetilde{W}_G$ itself. The inclusion relations between the Schubert varieties are given by the *Bruhat order* $\leq$ on $\widetilde{W}_G$, which is defined in a purely combinatorial way; see §5.2. Thus the problem of identifying which Schubert varieties are contained in the geometric special fiber of the spin local model amounts to the problem of identifying a certain finite subset of $\widetilde{W}_G$ closed under the Bruhat order.

Now let $\mathcal{A}$ be an apartment in the building for the adjoint group of G , and let $\mathcal{H}$ be the collection of affine root hyperplanes in $\mathcal{A}$. By [B, VI §2.5 Prop. 8], upon choosing a special vertex in $\mathcal{A}$ as origin, there exists a unique reduced root system Σ on $\mathcal{A}$ such that the hyperplanes $\mathcal{H}$ are precisely the affine root hyperplanes for Σ . The Bruhat order on $\widetilde{W}_G$ is governed by the affine Weyl group for Σ , and indeed the study of $(\widetilde{W}_G, \leq)$, at least for the purposes of this paper, reduces to that of an extended affine Weyl group for Σ , as is explained in [PRS, Rems. 4.4.7, 4.4.8]. For odd n , Σ is of type C , and Theorem 1.4 is proved in [Sm3] by exploiting a result of Haines–Ngô [HN] relating admissible sets for GSp_{2m} to *permissible* sets for GL_{2m} . See §5.3 for the definition of permissibility; the force of their result is that it is relatively easy to determine when a given element is permissible, at least as compared to when an element is admissible.

By contrast, for even $n = 2m$, Σ is of type B_m , and the results of Haines–Ngô do not apply. We are therefore forced to work through the combinatorics largely from scratch. To fix notation, let us make the harmless assumption that $s \leq r$, as we shall do in the body of the paper. We shall see in §§5.7–5.8 that $\widetilde{W}_G$, equipped with

its Bruhat order, identifies with the Iwahori-Weyl group $\widetilde{W}_{B_m}$ of split GO_{2m+1} , and that in this way the $\{\mu_{r,s}\}$ -admissible set identifies with the μ_B -admissible set in $\widetilde{W}_{B_m}$, where

$$\mu_B := (2^{(s)}, 1^{(2m+1-2s)}, 0^{(s)}).$$

In §§7.2–7.4 we shall obtain simple combinatorial conditions in $\widetilde{W}_G$ that describe the Schubert varieties contained in the geometric special fiber of the spin local model. To prove Theorem 1.4, we must then show that these conditions, translated to $\widetilde{W}_{B_m}$, characterize the μ_B -admissible set.

Rather than working directly with $\widetilde{W}_{B_m}$, we shall approach this problem by passing to the Iwahori-Weyl group $\widetilde{W}_{D_{m+1}}$ of split GO_{2m+2} . The root datum of GO_{2m+1} is a Steinberg fixed-point root datum (see §9.1) obtained from the root datum of GO_{2m+2} . Therefore one has a natural inclusion $\widetilde{W}_{B_m} \hookrightarrow \widetilde{W}_{D_{m+1}}$, and the Bruhat order on $\widetilde{W}_{D_{m+1}}$ restricts to the Bruhat order on $\widetilde{W}_{B_m}$. The cocharacter μ_B in $\widetilde{W}_{B_m}$ has image

$$\mu_D := (2^{(s)}, 1^{(2m+2-2s)}, 0^{(s)})$$

in $\widetilde{W}_{D_{m+1}}$. Motivated by the conditions characterizing the Schubert varieties that occur in the geometric special fiber of the spin local model, in (8.2.1) we introduce the notion of μ_D -spin-permissibility for elements in $\widetilde{W}_{D_{m+1}}$. The fundamental combinatorial result in this paper is the following.

Theorem 1.5. *An element $w \in \widetilde{W}_{D_{m+1}}$ is μ_D -admissible $\iff w$ is μ_D -spin-permissible.*

The proof of Theorem 1.5 will occupy almost all of §§8.3–8.12. We shall extend it to the general parahoric case in (8.14.5). With Theorem 1.5 (and its parahoric variants) in hand, the characterization of the μ_B -admissible set in $\widetilde{W}_{B_m}$ (and its parahoric variants) that we need for Theorem 1.4 follows easily, via the formalism of Steinberg fixed-point root data.

In the course of working out the combinatorics in §§8–9, we shall also show that for the cocharacters μ_B and μ_D , admissibility is equivalent to the property of *vertexwise admissibility*; see §8.15 and §9.7. This result for μ_B immediately implies the conjecture [PR4, Conj. 4.2] of Pappas and Rapoport. Vertexwise admissibility is defined in general in [PRS, §4.5]; particular instances of it appeared earlier (sometimes only implicitly) in papers of Pappas and Rapoport [PR1, PR2, PR4]. The definition is motivated by Pappas and Rapoport’s idea to describe the local model for general parahorics by first taking the flat closure in the maximal parahoric case, and then taking the intersection of the inverse images of these flat closures in the naive local model. See, for example, p. 512 and §4.2 in [PR4].

We shall also obtain very explicit descriptions of the μ_D -permissible set in $\widetilde{W}_{D_{m+1}}$ and its parahoric variants, and of the μ_B -permissible set in $\widetilde{W}_{B_m}$ and its parahoric variants; see (8.6.4), (8.16.1), and (9.8.1). It is a general result of Kottwitz–Rapoport [KR] that, for any cocharacter μ in a root datum, μ -admissibility implies μ -permissibility. Haines and Ngô [HN] subsequently showed that the converse can fail. More precisely, they showed that every irreducible root system of rank ≥ 4 and not of type A has coweights μ for which the μ -permissible set is strictly bigger than the μ -admissible set. For our particular cocharacters μ_B and μ_D as above, we shall show that if $m, s \geq 3$, then the μ_B - (resp. μ_D -)admissible and μ_B - (resp.

μ_D -)permissible sets in $\widetilde{W}_{B_m}$ (resp. $\widetilde{W}_{D_{m+1}}$) are not equal; see (8.6.5) and (9.8.3). In the case of $\widetilde{W}_{B_3}$, this gives an example of a cocharacter in a rank 3 root datum for which admissibility and permissibility do not coincide. In the case of $\widetilde{W}_{D_{m+1}}$, we find that admissibility and permissibility do not coincide for the cocharacter

$$(2^{(m)}, 1, 1, 0^{(m)}) = (1^{(m+1)}, 0^{(m+1)}) + (1^{(m)}, 0, 1, 0^{(m)}),$$

which is a sum of minuscule cocharacters for D_{m+1} contained in the same closed Weyl chamber. This gives a counterexample to Rapoport's conjecture [R, §3, p. 283] that admissibility and permissibility are equivalent for any cocharacter which is a sum of dominant minuscule cocharacters. See (8.6.5) for some further discussion.

Let us now outline the contents of the paper. In §2 we review the definitions of the various local models from [PR4]. In §3 we review some basic group-theoretic aspects of ramified, quasi-split GU_{2m} . In §4 we introduce some preliminary combinatorial considerations, essentially covering just enough so that we can translate the wedge and spin conditions into combinatorics later in §7. In §5 we review the general formalism of Iwahori-Weyl groups, and then specialize this to the case of our group GU_{2m} . In §6 we review the embedding of the geometric special fiber of the local model into an appropriate affine flag variety for GU_{2m} from [PR4]. In §7 we obtain combinatorial descriptions of the Schubert varieties contained in the geometric special fibers of M_I^{naive} , $M_I^\wedge$, and M_I^{spin} inside the affine flag variety, and we prove Theorems 1.3 and 1.4 modulo the later result (9.6.1), which characterizes the μ_B -admissible set in $\widetilde{W}_{B_m}$. Section 8 is the technical heart of the paper. In it we cover all of the type D combinatorics, most notably proving Theorem 1.5 and its parahoric generalization (8.14.5). We conclude the paper in §9 by working through the type B combinatorics, essentially all of which follows from §8 via the formalism of Steinberg fixed-point root data.

Acknowledgments. I thank Michael Rapoport for his encouragement to work on this problem and for introducing me to the subject. I also thank him, Thomas Haines, Robert Kottwitz, Stephen Kudla, and George Pappas for many helpful conversations.

Notation. Throughout the paper F/F_0 denotes a ramified quadratic extension of discretely valued, non-Archimedean fields with respective rings of integers $\mathcal{O}_F$ and $\mathcal{O}_{F_0}$ and respective uniformizers π and π_0 satisfying $\pi^2 = \pi_0$. We assume that the residue field is perfect of characteristic not 2, and we denote by k an algebraic closure of it. We also employ an auxiliary ramified quadratic extension K/K_0 of discretely valued, non-Archimedean Henselian fields with perfect residue field of characteristic not 2, with respective rings of integers $\mathcal{O}_K$ and $\mathcal{O}_{K_0}$, and with respective uniformizers u and t satisfying $u^2 = t$; eventually K and K_0 will be the fields of Laurent series $k((u))$ and $k((t))$, respectively. We put $\Gamma := \text{Gal}(K/K_0)$, and we write $x \mapsto \bar{x}$ for the action of the nontrivial element of Γ on K , so that $\bar{u} = -u$. Abusing notation, we continue to write $x \mapsto \bar{x}$ for the R -algebra automorphism of $K \otimes_{K_0} R$ induced by any base change $K_0 \rightarrow R$.

We fix once and for all an even integer $2m \geq 2$ and nonnegative integers r and s with $2m = r + s$.¹ It will be easy to see later on that the cocharacters $\mu_{r,s}$ and $\mu_{s,r}$

¹Strictly speaking, the paper of Pappas and Rapoport [PR4] restricts to unitary groups GU_n for $n \geq 3$, but all of our appeals to their paper will still go through for $n = 2m = 2$. On the other hand, local models for GU_2 are worked out explicitly in [PRS, Rem. 2.6.13].

defined above specify isomorphic local models, and that these cocharacters even have the same image in the Galois coinvariants $X_*(D \times \mathbb{G}_m)_\Gamma$. Hence for simplicity of notation we shall always assume $s \leq r$. Although almost everything we shall do will depend on m , r , and s , we shall usually not embed them into the notation. We continue to take $E := F_0$ if $r = s$ and $E := F$ if $r \neq s$.

We relate objects by writing $\simeq$ for isomorphic, $\cong$ for canonically isomorphic, and $=$ for equal.

For $x \in \mathbb{R}$, we write $\lfloor x \rfloor$ for the greatest integer $\leq x$ and $\lceil x \rceil$ for the least integer $\geq x$.

For any positive integer n , given a vector $v \in \mathbb{R}^n$, we write $v(j)$ for the j th entry of v , Σv for the sum of the entries of v , and v^* for the vector in $\mathbb{R}^n$ defined by $v^*(j) = v(n+1-j)$. Given another vector w , we write $v \geq w$ if $v(j) \geq w(j)$ for all j . We write $\mathbf{d}$ for the vector $(d, d, \dots, d)$, leaving it to context to make clear the number of entries. The expression $(d^{(i)}, e^{(j)}, \dots)$ denotes the vector with d repeated i times, followed by e repeated j times, and so on.

We write S_n for the symmetric group on $1, \dots, n$; S_n^* for the subgroup

$$S_n^* := \{ \sigma \in S_n \mid \sigma(n+1-j) = n+1-\sigma(j) \text{ for all } j \in \{1, \dots, n\} \};$$

and S_n° for the subgroup

$$S_n^\circ := \{ \sigma \in S_n^* \mid \sigma \text{ is even in } S_n \}.$$

Given facets $\mathbf{f}$ and $\mathbf{f}'$ in an apartment, we write $\mathbf{f} \preceq \mathbf{f}'$ if $\mathbf{f}$ is a subfacet of $\mathbf{f}'$, that is, if $\mathbf{f} \subset \overline{\mathbf{f}'}$. We say that a point v is a *vertex* of $\mathbf{f}$ if v is contained in a minimal subfacet of $\mathbf{f}$. In this paper, the apartments we shall consider will typically be attached to reductive groups which are not semisimple, and the minimal facets in such apartments are not themselves points.

2. UNITARY LOCAL MODELS

We begin by recalling the definition and some of the discussion of local models for even ramified unitary groups from [PR4]. Let $I \subset \{0, \dots, m\}$ be a nonempty subset satisfying property (1.1).

2.1. Pairings. Let $V := F^{2m}$. In this subsection we introduce some pairings on V and notation related to them.

Let $e_1, e_2, \dots, e_{2m}$ denote the standard ordered F -basis in V , and let

$$\phi: V \times V \longrightarrow F$$

be the F/F_0 -Hermitian form on V whose matrix with respect to the standard basis is

$$(2.1.1) \quad \begin{pmatrix} & & & 1 \\ & & \ddots & \\ & & & \\ 1 & & & \end{pmatrix}.$$

We attach to ϕ the respective alternating and symmetric F_0 -bilinear forms $V \times V \rightarrow F_0$

$$(2.1.2) \quad \langle x, y \rangle := \frac{1}{2} \operatorname{Tr}_{F/F_0}(\pi^{-1} \phi(x, y)) \quad \text{and} \quad (x, y) := \frac{1}{2} \operatorname{Tr}_{F/F_0}(\phi(x, y)).$$

For any $\mathcal{O}_F$ -lattice $\Lambda \subset V$, we denote by $\widehat{\Lambda}$ the ϕ -dual of Λ ,

$$(2.1.3) \quad \widehat{\Lambda} := \{ x \in V \mid \phi(\Lambda, x) \subset \mathcal{O}_F \}.$$

Then $\widehat{\Lambda}$ is also the $\langle \cdot, \cdot \rangle$ -dual of Λ ,

$$\widehat{\Lambda} = \{ x \in V \mid \langle \Lambda, x \rangle \subset \mathcal{O}_{F_0} \};$$

and $\widehat{\Lambda}$ is related to the $(\cdot, \cdot)$ -dual $\widehat{\Lambda}^s := \{ x \in V \mid (\Lambda, x) \subset \mathcal{O}_{F_0} \}$ by the formula $\widehat{\Lambda}^s = \pi^{-1}\widehat{\Lambda}$. Both $\widehat{\Lambda}$ and $\widehat{\Lambda}^s$ are $\mathcal{O}_F$ -lattices in V , and the forms $\langle \cdot, \cdot \rangle$ and $(\cdot, \cdot)$ induce perfect $\mathcal{O}_{F_0}$ -bilinear pairings

$$(2.1.4) \quad \Lambda \times \widehat{\Lambda} \xrightarrow{\langle \cdot, \cdot \rangle} \mathcal{O}_{F_0} \quad \text{and} \quad \Lambda \times \widehat{\Lambda}^s \xrightarrow{(\cdot, \cdot)} \mathcal{O}_{F_0}$$

for all Λ .

2.2. Standard lattices. For $i = 2mb + c$ with $0 \leq c < 2m$, we define the *standard $\mathcal{O}_F$ -lattice*

$$(2.2.1) \quad \Lambda_i := \sum_{j=1}^c \pi^{-b-1} \mathcal{O}_F e_j + \sum_{j=c+1}^{2m} \pi^{-b} \mathcal{O}_F e_j \subset V.$$

Then $\widehat{\Lambda}_i = \Lambda_{-i}$ for all i , and the Λ_i 's form a complete, periodic, self-dual lattice chain

$$\cdots \subset \Lambda_{-2} \subset \Lambda_{-1} \subset \Lambda_0 \subset \Lambda_1 \subset \Lambda_2 \subset \cdots,$$

which we call the *standard lattice chain*. More generally, for our given subset $I \subset \{0, \dots, m\}$, we denote by Λ_I the periodic, self-dual subchain of the standard chain consisting of all lattices of the form Λ_i for $i \in 2m\mathbb{Z} \pm I$. Thus $\Lambda_{\{0, \dots, m\}}$ denotes the standard chain itself.

The standard lattice chain admits the following obvious trivialization. Let $\beta_i: \mathcal{O}_F^{2m} \rightarrow \mathcal{O}_F^{2m}$ multiply the i th standard basis element ϵ_i by π and send all other standard basis elements to themselves. Then there is a unique isomorphism of chains of $\mathcal{O}_F$ -modules

$$(2.2.2) \quad \begin{array}{ccccccc} \cdots & \hookrightarrow & \Lambda_0 & \hookrightarrow & \Lambda_1 & \hookrightarrow & \cdots \hookrightarrow \Lambda_{2m} \hookrightarrow \cdots \\ & & \uparrow \sim & & \uparrow \sim & & \uparrow \sim \\ \cdots & \xrightarrow{\beta_{2m}} & \mathcal{O}_F^{2m} & \xrightarrow{\beta_1} & \mathcal{O}_F^{2m} & \xrightarrow{\beta_2} & \cdots \xrightarrow{\beta_{2m}} \mathcal{O}_F^{2m} \xrightarrow{\beta_1} \cdots \end{array}$$

such that the leftmost displayed vertical arrow identifies the ordered $\mathcal{O}_F$ -basis $\epsilon_1, \dots, \epsilon_{2m}$ of $\mathcal{O}_F^{2m}$ with the ordered basis $e_1, \dots, e_{2m}$ of Λ_0 . Restricting to subchains in the top and bottom rows in (2.2.2), we get an analogous trivialization of Λ_I for any I .

2.3. Naive local models. We now review the definition of the naive local models from [PR4, §1.5].

Recall our fixed partition $2m = r + s$ with $s \leq r$. The *naive local model* M_I^{naive} is the following contravariant functor on the category of $\mathcal{O}_E$ -algebras. Given an $\mathcal{O}_E$ -algebra R , an R -point in M_I^{naive} consists of, up to an obvious notion of isomorphism,

- a functor

$$\begin{aligned} \Lambda_I &\longrightarrow (\mathcal{O}_F \otimes_{\mathcal{O}_{F_0}} R\text{-modules}) \\ \Lambda_i &\longmapsto \mathcal{F}_i, \end{aligned}$$

where Λ_I is regarded as a category by taking the morphisms to be the inclusions of lattices in V ; together with

- an inclusion of $\mathcal{O}_F \otimes_{\mathcal{O}_{F_0}} R$ -modules $\mathcal{F}_i \hookrightarrow \Lambda_i \otimes_{\mathcal{O}_{F_0}} R$ for each $i \in 2m\mathbb{Z} \pm I$, functorial in Λ_i ;

satisfying the following conditions for all $i \in 2m\mathbb{Z} \pm I$.

- (LM1) Zariski-locally on $\text{Spec } R$, $\mathcal{F}_i$ embeds in $\Lambda_i \otimes_{\mathcal{O}_{F_0}} R$ as a direct R -module summand of rank $2m$.
- (LM2) The isomorphism $\Lambda_i \otimes_{\mathcal{O}_{F_0}} R \xrightarrow{\sim} \Lambda_{i-2m} \otimes_{\mathcal{O}_{F_0}} R$ obtained by tensoring $\Lambda_i \xrightarrow[\sim]{\pi} \pi\Lambda_i = \Lambda_{i-2m}$ identifies $\mathcal{F}_i$ with $\mathcal{F}_{i-2m}$.
- (LM3) The perfect R -bilinear pairing

$$(\Lambda_i \otimes_{\mathcal{O}_{F_0}} R) \times (\Lambda_{-i} \otimes_{\mathcal{O}_{F_0}} R) \xrightarrow{\langle \cdot, \cdot \rangle \otimes R} R$$

induced by (2.1.4) identifies $\mathcal{F}_i^\perp \subset \Lambda_{-i} \otimes_{\mathcal{O}_{F_0}} R$ with $\mathcal{F}_{-i}$.

- (LM4) The element $\pi \otimes 1 \in \mathcal{O}_F \otimes_{\mathcal{O}_{F_0}} R$ acts on $\mathcal{F}_i$ as an R -linear endomorphism with characteristic polynomial

$$\det(T \cdot \text{id} - \pi \otimes 1 \mid \mathcal{F}_i) = (T - \pi)^s (T + \pi)^r \in R[T].$$

When $E = F_0$, i.e. when $r = s = m$, the polynomial on the right-hand side of the display is to be interpreted as $(T^2 - \pi_0)^m$. The functor M_I^{naive} is plainly represented by a closed subscheme, which we again denote M_I^{naive} , of a finite product of Grassmannians over $\text{Spec } \mathcal{O}_E$. The F -generic fiber $M_I^{\text{naive}} \otimes_{\mathcal{O}_E} F$ can be identified with the Grassmannian of s -planes in a $2m$ -dimensional vector space; see [PR4, §1.5.3].

2.4. Wedge and spin conditions. In this subsection we review the wedge and spin conditions, which Pappas and Rapoport conjecture to define the honest local model inside M_I^{naive} .

The *wedge condition*, due to Pappas [P, §4], on an R -point $(\mathcal{F}_i)_i$ of M_I^{naive} is the condition that

- (LM5) if $r \neq s$, then for all $i \in 2m\mathbb{Z} \pm I$,

$$\bigwedge_R^{s+1} (\pi \otimes 1 + 1 \otimes \pi \mid \mathcal{F}_i) = 0 \quad \text{and} \quad \bigwedge_R^{r+1} (\pi \otimes 1 - 1 \otimes \pi \mid \mathcal{F}_i) = 0.$$

(There is no condition when $r = s$.)

We denote by $M_I^\wedge$ the subfunctor of M_I^{naive} of points that satisfy the wedge condition, and we call it the *wedge local model*. Plainly $M_I^\wedge$ is a closed subscheme of M_I^{naive} . As noted in [PR4, §1.5.6], the generic fibers of $M_I^\wedge$ and M_I^{naive} coincide.

To formulate the spin condition it is necessary to introduce some notation. For brevity we shall recall only the minimum that we need from [PR4, §7]; compare also [Sm1, §2.3; Sm3, §2.5]. Regarding V as a $4m$ -dimensional vector space over F_0 , consider the ordered F_0 -basis

$$-\pi^{-1}e_1, \dots, -\pi^{-1}e_m, e_{m+1}, \dots, e_{2m}, e_1, \dots, e_m, \pi e_{m+1}, \dots, \pi e_{2m},$$

which we denote by $f_1, \dots, f_{4m}$. Then $f_1, \dots, f_{4m}$ is *split* for the form $(\cdot, \cdot)$ from (2.1.2), that is, we have $(f_i, f_j) = \delta_{i, 4m+1-j}$ for all i and j .

Using the basis $f_1, \dots, f_{4m}$, we define an operator a on $\bigwedge_{F_0}^{2m} V$ as follows. For any subset $E \subset \{1, \dots, 4m\}$ of cardinality $2m$, let

$$(2.4.1) \quad f_E := f_{i_1} \wedge \dots \wedge f_{i_{2m}} \in \bigwedge_{F_0}^{2m} V,$$

where $E = \{i_1, \dots, i_{2m}\}$ with $i_1 < \dots < i_{2m}$. Given such E , we also let

$$E^\perp := (4m + 1 - E)^c = 4m + 1 - E^c,$$

where the set complements are taken in $\{1, \dots, 4m\}$. Then $E^\perp$ consists of the elements $j \in \{1, \dots, 4m\}$ such that $(f_i, f_j) = 0$ for all $i \in E$. We now define a by defining it on the basis elements f_E of $\bigwedge_{F_0}^{2m} V$ for varying E ,

$$a(f_E) := \text{sgn}(\sigma_E) f_{E^\perp},$$

where σ_E is the permutation on $\{1, \dots, 4m\}$ sending $\{1, \dots, 2m\}$ to the elements of E in increasing order, and sending $\{2m+1, \dots, 4m\}$ to the elements of $E^\perp$ in increasing order.

Remark 2.4.2. The definition of a is in entirely the same spirit as [Sm3, §2.5]. In analogy with [Sm3, Rem. 2.5.2], our a agrees only up to a sign of $(-1)^m$ with the analogous operators denoted $a_{f_1 \wedge \dots \wedge f_{4m}}$ in [PR4, disp. 7.6] and a in [Sm1, §2.3].

It follows easily from the definition of σ_E , or directly from [PR4, Prop. 7.1] and the preceding remark, that $\text{sgn}(\sigma_E) = \text{sgn}(\sigma_{E^\perp})$. Hence $a^2 = \text{id}_{\bigwedge_{F_0}^{2m} V}$. Hence $\bigwedge_{F_0}^{2m} V$ decomposes as

$$\bigwedge_{F_0}^{2m} V = \left(\bigwedge_{F_0}^{2m} V \right)_1 \oplus \left(\bigwedge_{F_0}^{2m} V \right)_{-1},$$

where

$$\left(\bigwedge_{F_0}^{2m} V \right)_{\pm 1} := \text{span}_{F_0} \{ f_E \pm \text{sgn}(\sigma_E) f_{E^\perp} \}_E$$

is the ± 1 -eigenspace for a ; here E ranges through the subsets of $\{1, \dots, 4m\}$ of cardinality $2m$.

Now consider the $\mathcal{O}_F$ -lattice $\Lambda_i \subset V$ for some i and regard it as an $\mathcal{O}_{F_0}$ -lattice. Then $\bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_i$ is naturally an $\mathcal{O}_{F_0}$ -lattice in $\bigwedge_{F_0}^{2m} V$. We set

$$\left(\bigwedge_{F_0}^{2m} \Lambda_i \right)_{\pm 1} := \left(\bigwedge_{F_0}^{2m} \Lambda_i \right) \cap \left(\bigwedge_{F_0}^{2m} V \right)_{\pm 1}.$$

To state the spin condition, recall our partition $2m = r + s$ and note that, given an R -point $(\mathcal{F}_i)_i$ of M_I^{naive} , the R -module $\bigwedge_R^{2m} \mathcal{F}_i$ is naturally contained in $\bigwedge_R^{2m} (\Lambda_i \otimes_{\mathcal{O}_{F_0}} R)$ for all i . The *spin condition*, due to Pappas and Rapoport [PR4, §7.2.1], is that

(LM6) for all $i \in 2m\mathbb{Z} \pm I$, $\bigwedge_R^{2m} \mathcal{F}_i$ is contained in

$$\text{im} \left[\left(\bigwedge_{F_0}^{2m} \Lambda_i \right)_{(-1)^s} \otimes_{\mathcal{O}_{F_0}} R \longrightarrow \bigwedge_R^{2m} (\Lambda_i \otimes_{\mathcal{O}_{F_0}} R) \right].$$

For fixed i , we say that $\mathcal{F}_i$ satisfies the spin condition if $\bigwedge_R^{2m} \mathcal{F}_i$ is contained in the displayed image. We denote by M_I^{spin} the subfunctor of $M_I^\wedge$ of points satisfying the spin condition, and we call it the *spin local model*. Plainly M_I^{spin} is a closed subscheme of $M_I^\wedge$. As noted in [PR4, §7.2.2], the generic fiber of M_I^{spin} agrees with the common generic fiber of M_I^{naive} and $M_I^\wedge$.

Remark 2.4.3. As we remarked in the odd case in [Sm3, Rem. 2.5.3], the statement of the spin condition in [PR4, §7.2.1] contains a sign error which traces to the sign discrepancy observed in (2.4.2). Indeed, in the even case the element $f_1 \wedge \dots \wedge f_{2m}$ lies in the $(-1)^m$ -eigenspace for the operator $a_{f_1 \wedge \dots \wedge f_{4m}}$ of [PR4]. Thus the argument of [PR4, §7.2.2] shows that the sign of $(-1)^s$ in the statement of the spin condition in [PR4] should be replaced with $(-1)^{s+m}$. For us, since $f_1 \wedge \dots \wedge f_{2m}$ lies in the $+1$ -eigenspace of our operator a , the same argument shows that we get a sign of $(-1)^s$.

2.5. Lattice chain automorphisms. Regarding Λ_I as a lattice chain over $\mathcal{O}_{F_0}$, the perfect pairings $\langle \cdot, \cdot \rangle$ of (2.1.4) give a *polarization* of Λ_I in the sense of [RZ, Def. 3.14] (with $B = F$, $b^* = \bar{b}$ in the notation of [RZ]). Let $\underline{\text{Aut}}(\Lambda_I)$ denote the $\mathcal{O}_{F_0}$ -group scheme of automorphisms of the lattice chain Λ_I that preserve the pairings $\langle \cdot, \cdot \rangle$ for variable $\Lambda_i \in \Lambda_I$ up to common unit scalar. Then $\underline{\text{Aut}}(\Lambda_I)$ is smooth and affine over $\mathcal{O}_{F_0}$; see [P, Th. 2.2], which in turn relies on [RZ, Th. 3.16]. The base change $\underline{\text{Aut}}(\Lambda_I)_{\mathcal{O}_E}$ acts naturally on M_I^{naive} , and it is easy to see that this action preserves the closed subschemes $M_I^\wedge$, M_I^{spin} , and M_I^{loc} . We shall return to this point in §6.4.

3. UNITARY SIMILITUDE GROUP

In this section we review a number of basic group-theoretic matters from [PR4]; these will become relevant when we begin to consider the affine flag variety. We switch to working with respect to the auxiliary field extension K/K_0 . We write $i^* := 2m + 1 - i$ for $i \in \{1, \dots, 2m\}$.

3.1. Unitary similitudes. Let h denote the Hermitian form on K^{2m} whose matrix with respect to the standard ordered basis is (2.1.1). We denote by

$$G := GU_{2m} := GU(h)$$

the algebraic group over K_0 of *unitary similitudes* of h : for any K_0 -algebra R , $G(R)$ is the group of elements $g \in GL_n(K \otimes_{K_0} R)$ satisfying $h_R(gx, gy) = c(g)h_R(x, y)$ for some $c(g) \in R^\times$ and all $x, y \in (K \otimes_{K_0} R)^n$, where h_R is the induced form on $(K \otimes_{K_0} R)^{2m}$. As the form h is nonzero, the scalar $c(g)$ is uniquely determined, and c defines an exact sequence of K_0 -groups

$$1 \longrightarrow U_{2m} \longrightarrow G \xrightarrow{c} \mathbb{G}_m \longrightarrow 1$$

with evident kernel $U_{2m} := U(h)$ the unitary group of h . The displayed sequence splits (noncanonically), so that the choice of a splitting presents G as a semidirect product $U_{2m} \rtimes \mathbb{G}_m$.

After base change to K , we get the standard identification

$$(3.1.1) \quad G_K \xrightarrow[\sim]{(\varphi, c)} (GL_{2m})_K \times (\mathbb{G}_m)_K,$$

where $\varphi: G_K \rightarrow (GL_{2m})_K$ is given on R -points by the map on matrix entries $K \otimes_{K_0} R \rightarrow R$ sending $x \otimes y \mapsto xy$.

3.2. Tori. We denote by S the standard diagonal maximal split torus in G : on R -points,

$$S(R) = \{ \text{diag}(a_1, \dots, a_{2m}) \in GL_n(R) \mid a_1 a_{2m} = \dots = a_m a_{m+1} \}.$$

The centralizer T of S is the standard maximal torus of all diagonal matrices in G ,

$$T(R) = \{ \text{diag}(a_1, \dots, a_{2m}) \in GL_n(K \otimes_{K_0} R) \mid a_1 \bar{a}_{2m} = \dots = a_m \bar{a}_{m+1} \}.$$

The isomorphism (3.1.1) identifies T_K with the split torus $D \times (\mathbb{G}_m)_K$, where D denotes the standard diagonal maximal torus in $(GL_n)_K$. The standard identification $X_*(D \times (\mathbb{G}_m)_K) \cong \mathbb{Z}^{2m} \times \mathbb{Z}$ then identifies the inclusion $X_*(S) \subset X_*(T)$ with

$$(3.2.1) \quad \{ (x_1, \dots, x_{2m}, y) \mid x_1 + x_{2m} = \dots = x_m + x_{m+1} = y \} \subset \mathbb{Z}^{2m} \times \mathbb{Z};$$

note that this is not the description of $X_*(S)$ given in [PR4, §2.4.1], which appears to contain an error.

We write $\mathcal{A}_G$ for the standard apartment $X_*(S) \otimes_{\mathbb{Z}} \mathbb{R}$, and we regard it as a subspace of $\mathbb{R}^{2m} \times \mathbb{R}$ via (3.2.1).

3.3. Kottwitz homomorphism. Let L be the completion of a maximal unramified extension of K_0 , let $\bar{L}$ be an algebraic closure of L , and let $\mathcal{J} := \text{Gal}(\bar{L}/L)$. Let $\mathcal{G}$ be a connected reductive group defined over L . In [Ko2, §7] Kottwitz defines a surjective functorial surjection

$$\kappa_{\mathcal{G}}: \mathcal{G}(L) \twoheadrightarrow \pi_1(\mathcal{G})_{\mathcal{J}},$$

where the target is the $\mathcal{J}$ -coinvariants of the fundamental group in the sense of Borovoi [Bo].

In the case of our group G , it is harmless to work over K_0 itself, and we recall from [PR4, §1.2.3(b)] that κ_G may be described as the surjective map

$$\kappa_G: G(K_0) \twoheadrightarrow \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})$$

given on the first factor by $g \mapsto \text{ord}_t c(g)$ and on the second factor by $g \mapsto \text{ord}_u x \bmod 2$; here $\det(g)c(g)^{-m} \in K^\times$ has norm 1 and, using Hilbert's Theorem 90, $x \in K^\times$ is any element satisfying $x\bar{x}^{-1} = \det(g)c(g)^{-m}$.

3.4. Parahoric subgroups. We next recall the description of the parahoric subgroups of $G(K_0)$ from [PR4]. In analogy with §2.2, for $i = 2mb + c$ with $0 \leq c < 2m$, we define the $\mathcal{O}_K$ -lattice

$$(3.4.1) \quad \lambda_i := \sum_{j=1}^c u^{-b-1} \mathcal{O}_K e_j + \sum_{j=c+1}^{2m} u^{-b} \mathcal{O}_K e_j \subset K^{2m},$$

where now $e_1, \dots, e_{2m}$ denotes the standard ordered basis in K^{2m} . For any nonempty subset $I \subset \{0, \dots, m\}$, we write λ_I for the chain consisting of all lattices λ_i for $i \in 2m\mathbb{Z} \pm I$, and we define

$$\begin{aligned} P_I &:= \{ g \in G(K_0) \mid g\lambda_i = \lambda_i \text{ for all } i \in 2m\mathbb{Z} \pm I \} \\ &= \{ g \in G(K_0) \mid g\lambda_i = \lambda_i \text{ for all } i \in I \}. \end{aligned}$$

In contrast with the odd case [PR4, §1.2.3(a)], P_I is not a parahoric subgroup in general since it may contain elements with nontrivial Kottwitz invariant. Let

$$P_I^\circ := P_I \cap \ker \kappa_G.$$

Proposition 3.4.2 (Pappas–Rapoport [PR4, §1.2.3(b)]). *P_I° is a parahoric subgroup of $G(K_0)$, and every parahoric subgroup of $G(K_0)$ is conjugate to P_I° for a unique nonempty $I \subset \{0, \dots, m\}$ satisfying (1.1). For such I , we have $P_I^\circ = P_I$ exactly when $m \in I$. The special maximal parahoric subgroups are exactly those conjugate to $P_{\{m\}}^\circ = P_{\{m\}}$.* $\square$

3.5. Affine roots. In terms of the identification (3.2.1), the *relative roots* of S in G consist of the maps $\{\pm\alpha_{i,j}\}_{i < j < i^*} \cup \{\pm\alpha_{i,i^*}\}_{i \leq m}$ on $X_*(S)$, where

$$\alpha_{i,j}: (x_1, \dots, x_{2m}, y) \mapsto x_i - x_j.$$

The *affine roots* are calculated in [PR4, §2.4.1] (modulo the description of $X_*(S)$ there), but in a way that implicitly chooses coordinates on $\mathcal{A}_G$ such that the origin is a special vertex. Later on we shall wish to identify the action of G on its building

with the action of G on lattices given by its standard representation, and to do so in the most straightforward way will require a different choice of coordinates: (3.4.2) says that the parahoric subgroup attached to λ_m , and not to λ_0 (which will correspond to the origin), is special.

We shall calculate the affine roots with regard to the following choices. Let $\nu: T(K_0) \rightarrow \mathcal{A}_G$ be the unique map satisfying

$$\chi(\nu(x)) = \text{ord}_t(\chi(x)) \quad \text{for all } x \in T(K_0) \text{ and } \chi \in X^*(T);$$

here ν is the opposite of the map defined in Tits's article [T, §1.2], but this will have no impact on the calculation of the affine roots. Let N be the normalizer of S in G . Then N is the scheme of monomial matrices in G , and we have an internal semidirect product $N(K_0) = T(K_0) \rtimes S_{2m}^*$, where we identify S_{2m}^* with the permutation matrices in $G(K_0)$. We let $N(K_0)$ act by affine transformations on $\mathcal{A}_G$ by letting $T(K_0)$ act by translations via ν , and by letting S_{2m}^* act via its natural linear action on $X_*(S)$. With respect to this choice of action, a calculation completely analogous to the one for quasi-split SU_n with n odd in [T, §1.15] shows that the affine roots for G consist of the functions

$$(3.5.1) \quad \pm \alpha_{i,j} + \frac{1}{2}\mathbb{Z} \quad \text{for } i < j < i^* \quad \text{and} \quad \pm \alpha_{i,i^*} + \frac{1}{2} + \mathbb{Z} \quad \text{for } i < m+1.$$

4. PRELIMINARY COMBINATORICS

In this section we introduce some preliminary combinatorial considerations, covering just enough to be able to translate the wedge and spin conditions into combinatorics later in §7. Let n be a positive integer, and for $j \in \{1, \dots, n\}$, let $j^* := n+1-j$.

4.1. The group $\widetilde{W}_n$. Let

$$(4.1.1) \quad X_{*n} := \{v \in \mathbb{Z}^n \mid v + v^* = \mathbf{d} \text{ for some } d \in \mathbb{Z}\}.$$

Then the group S_n^* acts on X_{*n} by permuting the factors of $\mathbb{Z}^n$, and we form the semidirect product

$$(4.1.2) \quad \widetilde{W}_n := X_{*n} \rtimes S_n^*.$$

Note that if n is odd, then the integer d appearing the definition of X_{*n} must be even, since $\lceil n/2 \rceil^* = \lceil n/2 \rceil$.

4.2. Faces of type (n, I) . Let I be a nonempty subset of $\{1, \dots, \lfloor n/2 \rfloor\}$.

Definition 4.2.1. Let $d \in \mathbb{Z}$. A d -face of type (n, I) is a family $\mathbf{v} = (v_i)_{i \in n\mathbb{Z} \pm I}$ of vectors $v_i \in \mathbb{Z}^n$ such that

- (F1) $v_{i+n} = v_i - \mathbf{1}$ for all $i \in n\mathbb{Z} \pm I$;
- (F2) $v_i \geq v_j$ for all $i, j \in n\mathbb{Z} \pm I$ with $i \leq j$;
- (F3) $\Sigma v_i - \Sigma v_j = j - i$ for all $i, j \in n\mathbb{Z} \pm I$; and
- (F4) $v_i + v_{-i}^* = \mathbf{d}$ for all $i \in n\mathbb{Z} \pm I$.

A family of vectors $\mathbf{v} = (v_i)_{i \in n\mathbb{Z} \pm I}$ is a *face of type (n, I)* if it is a d -face of type (n, I) for some d .

When the integer n is clear from context, we typically say just “face of type I .” Faces of type I were defined by Kottwitz–Rapoport in [KR, §§9–10], at least for even n , where they are called “ G -faces of type $n\mathbb{Z} \pm I$.” (Kottwitz–Rapoport use the term “face of type $n\mathbb{Z} \pm I$ ” for the notion (4.2.1) without condition (F4).)

As in [Sm1, Sm3], we have adopted a different convention from Kottwitz–Rapoport in formulating (4.2.1), corresponding to a different choice of base alcove, to better facilitate working with the affine flag variety later on.

Note that if n is odd, then it is an easy consequence of (F3) and (F4) that d -faces only occur for even d .

For $i = bn + c$ with $b \in \mathbb{Z}$ and $0 \leq c < n$, let

$$(4.2.2) \quad \omega_i := ((-1)^{(c)}, 0^{(n-c)}) - \mathbf{b}.$$

Then the family

$$(4.2.3) \quad \omega_I := (\omega_i)_{i \in n\mathbb{Z} \pm I}$$

is a 0-face of type I , which we call the *standard face of type I* .

The group $\widetilde{W}_n$ acts naturally on $\mathbb{Z}^n$ by affine transformations, and this induces an action of $\widetilde{W}_n$ on faces of type I ,

$$w \cdot (v_i)_{i \in n\mathbb{Z} \pm I} = (wv_i)_{i \in n\mathbb{Z} \pm I},$$

which one easily sees to be transitive. In the “Iwahori” case $I = \{0, \dots, \lfloor n/2 \rfloor\}$, $\widetilde{W}_n$ acts simply transitively. For nonempty $J \subset I$, there is a natural $\widetilde{W}_n$ -equivariant surjection

$$\begin{aligned} \{\text{faces of type } I\} &\twoheadrightarrow \{\text{faces of type } J\} \\ (v_i)_{i \in n\mathbb{Z} \pm I} &\longmapsto (v_i)_{i \in n\mathbb{Z} \pm J}. \end{aligned}$$

4.3. The vector $\mu_i^{\mathbf{v}}$. We continue with I a nonempty subset of $\{0, \dots, \lfloor n/2 \rfloor\}$. Given a face $\mathbf{v} = (v_i)_i$ of type I , let

$$(4.3.1) \quad \mu_i^{\mathbf{v}} := v_i - \omega_i \quad \text{for } i \in n\mathbb{Z} \pm I.$$

Then condition (F1) is equivalent to the periodicity relation

$$(4.3.2) \quad \mu_i^{\mathbf{v}} = \mu_{i+n}^{\mathbf{v}} \quad \text{for all } i,$$

and (F3) is equivalent to

$$(4.3.3) \quad \Sigma \mu_i^{\mathbf{v}} = \Sigma \mu_j^{\mathbf{v}} \quad \text{for all } i, j.$$

If $\mathbf{v}$ is a d -face, then condition (F4) is equivalent to

$$(4.3.4) \quad \mu_i^{\mathbf{v}} + (\mu_{-i}^{\mathbf{v}})^* = \mathbf{d} \quad \text{for all } i,$$

from which we also determine

$$(4.3.5) \quad \Sigma \mu_i^{\mathbf{v}} = nd/2.$$

We now prove a couple of lemmas. For $i \in I$, we define subsets of $\{1, \dots, n\}$

$$(4.3.6) \quad A_i := \{1, 2, \dots, i, i^*, i^* + 1, \dots, n\} \quad \text{and} \quad B_i := \{i + 1, i + 2, \dots, n - i\}.$$

For arbitrary $i \in \mathbb{Z}$, we define $A_i := A_{i'}$ and $B_i := B_{i'}$, where i' is the unique element in $\{0, \dots, \lfloor n/2 \rfloor\}$ that is congruent mod n to i or $-i$.

Lemma 4.3.7 (Basic inequalities). *Let $\mathbf{v}$ be a d -face of type I . Then for any $i \in I$, we have*

$$\begin{aligned} d &\leq \mu_i^{\mathbf{v}}(j) + \mu_i^{\mathbf{v}}(j^*) \leq d + 1, \\ d - 1 &\leq \mu_{-i}^{\mathbf{v}}(j) + \mu_{-i}^{\mathbf{v}}(j^*) \leq d \end{aligned} \quad \text{for } j \in A_i$$

and

$$\begin{aligned} d - 1 &\leq \mu_i^{\mathbf{v}}(j) + \mu_i^{\mathbf{v}}(j^*) \leq d, \\ d &\leq \mu_{-i}^{\mathbf{v}}(j) + \mu_{-i}^{\mathbf{v}}(j^*) \leq d + 1 \end{aligned} \quad \text{for } j \in B_i.$$

Proof. Since faces of type $\{0, \dots, \lfloor n/2 \rfloor\}$ surject onto faces of type I , it suffices to assume that $I = \{0, \dots, \lfloor n/2 \rfloor\}$. Modulo conventions related to the choice of base alcove, the inequalities in this case for $\mu_i^{\mathbf{v}}(j) + \mu_i^{\mathbf{v}}(j^*)$ are proved for n even and $d = 0$ in [Sm2, Lem. 4.4.1], and the argument there works just as well for arbitrary n and d . The inequalities for $\mu_{-i}^{\mathbf{v}}(j) + \mu_{-i}^{\mathbf{v}}(j^*)$ then follow from (4.3.4). $\square$

Of course, periodicity gives analogous inequalities for $\mu_i^{\mathbf{v}}(j) + \mu_i^{\mathbf{v}}(j^*)$ for any $i \in n\mathbb{Z} \pm I$.

Remark 4.3.8. Suppose that n is odd. Then d is even, and the basic inequalities immediately imply that $\mu_i^{\mathbf{v}}(\lfloor n/2 \rfloor) = d/2$ for all i .

Definition 4.3.9. Let $\mathbf{v}$ be a d -face of type I , and let $i \in n\mathbb{Z} \pm I$. We say that $\mu_i^{\mathbf{v}}$ is *self-dual* if $\mu_i^{\mathbf{v}} = \mu_{-i}^{\mathbf{v}}$, or in other words, if $\mu_i^{\mathbf{v}} + (\mu_i^{\mathbf{v}})^* = \mathbf{d}$.

Lemma 4.3.10. Let $\mathbf{v}$ be a d -face of type I . Let $i \in n\mathbb{Z} \pm I$, and suppose that the equality $\mu_i^{\mathbf{v}}(j) + \mu_i^{\mathbf{v}}(j^*) = d$ holds for all $j \in A_i$ or for all $j \in B_i$. Then $\mu_i^{\mathbf{v}}$ is self-dual.

Proof. By (4.3.5) we have $\Sigma \mu_i^{\mathbf{v}} = nd/2$, and the conclusion is then an easy consequence of the basic inequalities. $\square$

4.4. Further lemmas. We continue with $I \subset \{0, \dots, \lfloor n/2 \rfloor\}$ nonempty. The aim of this subsection is to prove (4.4.3) below, which we will have occasion to use at a number of points later on in our study of admissibility and spin-permissibility. For S a subset of $\{1, \dots, n\}$, we write $S^* := n + 1 - S$.

Lemma 4.4.1. Let $\mathbf{v}$ be a 2-face of type I , and let $i \in I$. Suppose that $\mathbf{0} \leq \mu_i^{\mathbf{v}} \leq \mathbf{2}$. Then

$$\#\{j \mid \mu_i^{\mathbf{v}}(j) = 2 \text{ and } \mu_i^{\mathbf{v}}(j^*) = 1\} = \#\{j \mid \mu_i^{\mathbf{v}}(j) = 0 \text{ and } \mu_i^{\mathbf{v}}(j^*) = 1\}.$$

Proof. Let

$$\begin{aligned} E &:= \{j \mid \mu_i^{\mathbf{v}}(j) = 2 \text{ and } \mu_i^{\mathbf{v}}(j^*) = 1\}, \\ F &:= \{j \mid \mu_i^{\mathbf{v}}(j) = 0 \text{ and } \mu_i^{\mathbf{v}}(j^*) = 1\}, \\ G &:= \{j \mid \mu_i^{\mathbf{v}}(j) = 2 \text{ and } \mu_i^{\mathbf{v}}(j^*) = 0\}, \text{ and} \\ H &:= \{j \mid \mu_i^{\mathbf{v}}(j) = 1 \text{ and } \mu_i^{\mathbf{v}}(j^*) = 1\}, \end{aligned} \tag{4.4.2}$$

and let e, f, g , and h denote the respective cardinalities of these sets. The basic inequalities (4.3.7) and our assumptions on $\mathbf{v}$ imply that

$$\{1, \dots, n\} = E \amalg F \amalg G \amalg H \amalg E^* \amalg F^* \amalg G^*.$$

Counting cardinalities on both sides of the display, we get

$$n = 2e + 2f + 2g + h.$$

On the other hand, by (4.3.5)

$$n = \Sigma \mu_i^{\mathbf{v}} = 3e + f + 2g + h.$$

Hence $e = f$. $\square$

Lemma 4.4.3. Under the same assumptions as in the previous lemma,

$$\#\{j \mid \mu_i^{\mathbf{v}}(j) = 2\} = \#\{j \mid \mu_i^{\mathbf{v}}(j) = 0\}.$$

Proof. In the notation of (4.4.2), we have

$$\{j \mid \mu_i^\vee(j) = 2\} = E \amalg G \quad \text{and} \quad \{j \mid \mu_i^\vee(j) = 0\} = F \amalg G^*.$$

Now use (4.4.1). $\square$

5. IWAHORI-WEYL GROUPS

In this section we review some generalities on Iwahori-Weyl groups and the $\{\mu\}$ -admissible set, and we make these generalities explicit in the case of our group G . We then show that the Iwahori-Weyl group for G , equipped with its Bruhat order, is isomorphic to the Iwahori-Weyl group for split GO_{2m+1} . The significance of the Iwahori-Weyl group (and certain parahoric variants of it) is that, in the function field case, it indexes the Schubert varieties in the affine flag variety, and the Bruhat order gives the inclusion relations between the Schubert varieties. We take [PRS, §4] as a general reference. We write $i^* := 2m + 1 - i$.

5.1. Iwahori-Weyl groups. Let L be a complete, discretely valued, strictly Henselian field with algebraic closure $\overline{L}$ and Galois group $\mathcal{I} := \text{Gal}(\overline{L}/L)$. Let $\mathcal{G}$ be a connected reductive group over L . Let $\mathcal{S}$ be a maximal split torus in $\mathcal{G}$, and let $\mathcal{T}$ and $\mathcal{N}$ be its respective normalizer and centralizer in $\mathcal{G}$. Then $\mathcal{T}$ is a maximal torus since by Steinberg's theorem $\mathcal{G}$ is quasi-split. The *Iwahori-Weyl group* of $\mathcal{G}$ with respect to $\mathcal{S}$ is the group

$$\widetilde{W} := \mathcal{N}(L)/\mathcal{T}(L)_1,$$

where $\mathcal{T}(L)_1$ is the kernel of the Kottwitz homomorphism

$$\kappa_{\mathcal{T}}: \mathcal{T}(L) \twoheadrightarrow X_*(\mathcal{T})_{\mathcal{G}}.$$

The evident exact sequence

$$1 \longrightarrow \mathcal{T}(L)/\mathcal{T}(L)_1 \longrightarrow \widetilde{W} \longrightarrow \mathcal{N}(L)/\mathcal{T}(L) \longrightarrow 1$$

splits by Haines–Rapoport [HR, Prop. 13]. Hence, via the Kottwitz homomorphism, $\widetilde{W}$ is expressible as a semidirect product

$$\widetilde{W} \simeq X_*(\mathcal{T})_{\mathcal{G}} \rtimes W,$$

where $W := \mathcal{N}(L)/\mathcal{T}(L)$ is the relative Weyl group of $\mathcal{S}$ in $\mathcal{G}$. Given $\mu \in X_*(\mathcal{T})_{\mathcal{G}}$, we usually write t_μ when we wish to regard μ as an element of $\widetilde{W}$, and we refer to such elements in $\widetilde{W}$ as *translation elements*.

If $\mathcal{R} = (X^*, X_*, \Phi, \Phi^\vee)$ is a root datum, then we define its *extended affine Weyl group* $\widetilde{W}_{\mathcal{R}} := X_* \rtimes W_{\mathcal{R}}$, where $W_{\mathcal{R}}$ denotes the Weyl group of $\mathcal{R}$. If $\mathcal{G}$ is split with root datum $\mathcal{R}$ attached to $\mathcal{S}$, then we obtain a canonical isomorphism $\widetilde{W} \cong \widetilde{W}_{\mathcal{R}}$.

To consider local models with general parahoric, not just Iwahori, level structure, it is necessary to consider certain double coset variants of $\widetilde{W}$. Let $\mathcal{A} := X_*(\mathcal{S}) \otimes \mathbb{R}$ denote the apartment attached to $\mathcal{S}$. The group $\widetilde{W}$ acts on $\mathcal{A}$ by affine transformations in a way that is prescribed uniquely up to isomorphism in [T, §1.2]. Having chosen such an action, we may then consider the affine root system Φ_{aff} and the attendant affine root hyperplanes. Fix an alcove $\mathbf{a}$ in $\mathcal{A}$, and let $\Pi_{\mathbf{a}}$ denote the set of reflections across the walls of $\mathbf{a}$. The *affine Weyl group* W_{aff} is the group of affine transformations on $\mathcal{A}$ generated by $\Pi_{\mathbf{a}}$; it lifts canonically to a normal subgroup

of $\widetilde{W}$. In fact the Kottwitz homomorphism $\kappa_{\mathcal{G}}$, restricted to $\mathcal{N}(L)$, induces an isomorphism

$$\widetilde{W}/W_{\text{aff}} \xrightarrow[\sim]{\kappa_{\mathcal{G}}} \pi_1(\mathcal{G})_{\mathcal{I}},$$

as is explained in [HR, p. 196] or [PRS, §4.2]. For $\mathbf{f}$ a subfacet of $\mathbf{a}$, we denote by $W_{\mathbf{f}}$ the common stabilizer and pointwise fixer of $\mathbf{f}$ in $\widetilde{W}_{\text{aff}}$. Then $W_{\mathbf{f}}$ is Coxeter group, generated by the reflections across the walls of $\mathbf{a}$ containing $\mathbf{f}$ [B, V §3.3 Prop. 2]. If P is the parahoric subgroup of $\mathcal{G}(L)$ corresponding to $\mathbf{f}$, then

$$W_{\mathbf{f}} = (\mathcal{N}(L) \cap P) / \mathcal{T}(L)_1;$$

see [PRS, §4.2]. The double coset variants of $\widetilde{W}$ that we shall be interested in are those of the form $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ for $\mathbf{f}_1, \mathbf{f}_2 \preceq \mathbf{a}$.

5.2. Bruhat order. We continue with the notation of the previous subsection. Let $\Omega_{\mathbf{a}}$ denote the stabilizer of $\mathbf{a}$ in $\widetilde{W}$. The affine Weyl group W_{aff} acts simply transitively on the alcoves in $\mathcal{A}$, and therefore $\widetilde{W}$ decomposes as a semidirect product

$$\widetilde{W} = W_{\text{aff}} \rtimes \Omega_{\mathbf{a}}.$$

The pair $(W_{\text{aff}}, \Pi_{\mathbf{a}})$ is a Coxeter system. Hence W_{aff} is endowed with a length function ℓ and Bruhat order $\leq$, which extend to $\widetilde{W}$ via the rules $\ell(w\tau) = \ell(w)$ and $w\tau \leq w'\tau'$ if $\tau = \tau'$ and $w \leq w'$, where $w, w' \in W_{\text{aff}}$ and $\tau, \tau' \in \Omega_{\mathbf{a}}$.

For subfacets $\mathbf{f}_1, \mathbf{f}_2 \preceq \mathbf{a}$, the Bruhat order extends to $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ as follows. Each double coset $w \in W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ is known to contain a unique element ${}^{\mathbf{f}_1}w^{\mathbf{f}_2} \in \widetilde{W}$ of minimal length, and one then defines $w \leq x$ in $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ if ${}^{\mathbf{f}_1}w^{\mathbf{f}_2} \leq {}^{\mathbf{f}_1}x^{\mathbf{f}_2}$ in $\widetilde{W}$. The Bruhat order has the property that if $\tilde{w} \leq \tilde{x}$ in $\widetilde{W}$, then $W_{\mathbf{f}_1} \tilde{w} W_{\mathbf{f}_2} \leq W_{\mathbf{f}_1} \tilde{x} W_{\mathbf{f}_2}$.

5.3. Admissible and permissible sets. We now recall the key notion of $\{\mu\}$ -admissibility, and the related notion of $\{\mu\}$ -permissibility. We continue with the notation of the previous two subsections.

Let $\{\mu\}$ denote a $\mathcal{G}(\overline{L})$ -conjugacy class of cocharacters of $\mathcal{G}$, or what amounts to the same, a W^{abs} -conjugacy class in $X_*(\mathcal{T})$, where $W^{\text{abs}} := \mathcal{N}(\overline{L}) / \mathcal{T}(\overline{L})$ is the absolute Weyl group of $\mathcal{T}$ in $\mathcal{G}$. Let $\tilde{\Lambda}_{\{\mu\}} \subset \{\mu\}$ denote the subset of cocharacters contained in the closure of some absolute Weyl chamber corresponding to a Borel subgroup of $\mathcal{G}$ containing $\mathcal{T}$ and defined over L . Such Borel subgroups exist since $\mathcal{G}$ is quasi-split, and the elements of $\tilde{\Lambda}_{\{\mu\}}$ form a single W -orbit, since all such Borel subgroups are W -conjugate. Let $\Lambda_{\{\mu\}}$ denote the image of $\tilde{\Lambda}_{\{\mu\}}$ in the coinvariants $X_*(\mathcal{T})_{\mathcal{I}}$. Finally, let $\mathbf{f}_1, \mathbf{f}_2 \preceq \mathbf{a}$.

Definition 5.3.1 (Kottwitz–Rapoport [KR, Intro.]; Rapoport [R, disp. 3.6]). An element $w \in W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ is $\{\mu\}$ -*admissible* if there exists $\lambda \in \Lambda_{\{\mu\}}$ such that $w \leq W_{\mathbf{f}_1} t_{\lambda} W_{\mathbf{f}_2}$. We write $\text{Adm}_{\mathbf{f}_1, \mathbf{f}_2}(\{\mu\})$, or $\text{Adm}_{\mathbf{f}_1, \mathbf{f}_2, \mathcal{G}}(\{\mu\})$ when we wish to emphasize $\mathcal{G}$, for the set of $\{\mu\}$ -admissible elements in $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$. We abbreviate this to $\text{Adm}_{\mathcal{G}}(\{\mu\}) = \text{Adm}(\{\mu\}) \subset \widetilde{W}$ when $\mathbf{f}_1 = \mathbf{f}_2 = \mathbf{a}$.

In the special case that $\mathcal{G}$ is split, we have $\tilde{\Lambda}_{\{\mu\}} = \{\mu\} = W^{\text{abs}}\mu = W\mu$ for any $\mu \in \{\mu\}$. We then typically speak of μ -admissibility in place of $\{\mu\}$ -admissibility, and we write $\text{Adm}_{\mathbf{f}_1, \mathbf{f}_2}(\mu)$. If $\mathcal{R}$ is a root datum, then we analogously define $\text{Adm}_{\mathbf{f}_1, \mathbf{f}_2, \mathcal{R}}(\mu) \subset W_{\mathbf{f}_1} \backslash \widetilde{W}_{\mathcal{R}} / W_{\mathbf{f}_2}$.

Let $\{\bar{\mu}\}$ denote the image of the conjugacy class $\{\mu\}$ in $X_*(\mathcal{T})_{\mathcal{G}}$. It is conjectured in [PRS, Conj. 4.5.3] that $\Lambda_{\{\mu\}}$ consists of exactly the subset of elements in $\{\bar{\mu}\}$ that are maximal for the Bruhat order. Note that if this conjecture holds true, then the notion of $\{\mu\}$ -admissibility is just that $w \leq W_{\mathbf{f}_1} t_{\bar{\mu}} W_{\mathbf{f}_2}$ for some $\bar{\mu} \in \{\bar{\mu}\}$. We shall see explicitly in §5.8 that the conjecture holds for the pair $(G, \{\mu_{r,s}\})$, and it is not hard to see that furthermore the conjecture holds for any geometric conjugacy class for G .

Now let $\text{Conv}(\Lambda_{\{\mu\}})$ denote the convex hull of the image of $\Lambda_{\{\mu\}}$ in $X_*(\mathcal{T})_{\mathcal{G}} \otimes \mathbb{R} \cong \mathcal{A}$, and recall from [R, Lem. 3.1] that all elements of $\Lambda_{\{\mu\}}$, regarded as elements of $\widetilde{W}$, are congruent mod W_{aff} . Suppose furthermore that $\mathbf{f}_2 \preceq \mathbf{f}_1 \preceq \mathbf{a}$.

Definition 5.3.2 (Kottwitz-Rapoport [KR, Intro.]; Rapoport [R, disp. 3.10]). An element $w \in W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_1}$ is $\{\mu\}$ -*permissible* if $w \equiv t_{\lambda} \text{ mod } W_{\text{aff}}$ for one, hence any, $\lambda \in \Lambda_{\{\mu\}}$ and $wv - v \in \text{Conv}(\Lambda_{\{\mu\}})$ for all $v \in \mathbf{f}_2$. We write $\text{Perm}_{\mathbf{f}_1, \mathbf{f}_2}(\{\mu\})$, or $\text{Perm}_{\mathbf{f}_1, \mathbf{f}_2, \mathcal{G}}(\{\mu\})$ when we wish to emphasize $\mathcal{G}$, for the set of $\{\mu\}$ -permissible elements in $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_1}$. We abbreviate this to $\text{Perm}_{\mathcal{G}}(\{\mu\}) = \text{Perm}(\{\mu\}) \subset \widetilde{W}$ when $\mathbf{f}_1 = \mathbf{f}_2 = \mathbf{a}$.

It is clear that $\{\mu\}$ -permissibility is well-defined on right $W_{\mathbf{f}_2}$ -cosets. That it is also well-defined on left $W_{\mathbf{f}_1}$ -cosets is shown immediately after display 3.10 in [R]. By convexity, the condition

$$wv - v \in \text{Conv}(\Lambda_{\{\mu\}}) \text{ for all } v \in \mathbf{f}_2$$

is equivalent to the condition

$$\text{for all minimal subfacets } \mathbf{f} \preceq \mathbf{f}_2, wv - v \in \text{Conv}(\Lambda_{\{\mu\}}) \text{ for one, hence any, } v \in \mathbf{f}.$$

In the split or root datum case, we typically speak of μ -permissibility, for any $\mu \in \{\mu\}$, in place of $\{\mu\}$ -permissibility, and we write $\text{Perm}_{\mathbf{f}_1, \mathbf{f}_2}(\mu)$.

5.4. Case of GU_{2m} . We now begin to apply the generalities of the three previous subsections to our even unitary similitude group G over K_0 and its maximal split torus S from §3. We shall assume throughout the rest of §5 that the residue field of K_0 is algebraically closed.

We first compute the Galois coinvariants of $X_*(T)$. Since G splits over K , the action of the Galois group on $X_*(T)$ factors through Γ . In terms of our identification $X_*(T) \simeq \mathbb{Z}^{2m} \times \mathbb{Z}$ from §3.2, the nontrivial element in Γ acts on $X_*(T)$ by sending

$$(x_1, \dots, x_{2m}, y) \mapsto (y - x_{2m}, \dots, y - x_1, y).$$

Let

$$X := \{v \in \mathbb{Z}^{2m} \mid v + v^* = \mathbf{d} \text{ for some } d \in 2\mathbb{Z}\}.$$

It follows that the surjective map

$$(5.4.1) \quad \begin{array}{ccc} X_*(T) & \xrightarrow{\hspace{1.5cm}} & X \\ (x_1, \dots, x_{2m}, y) & \longmapsto & (x_1 - x_{2m} + y, x_2 - x_{2m+1} + y, \dots, x_{2m} - x_1 + y) \end{array}$$

identifies

$$(5.4.2) \quad X_*(T)_{\Gamma} \xrightarrow{\sim} X.$$

In this way the Kottwitz map for T is given by

$$\begin{aligned} \kappa_T: \quad T(K_0) &\longrightarrow X \\ \text{diag}(a_1, \dots, a_{2m}) &\longmapsto (\text{ord}_u(a_1), \dots, \text{ord}_u(a_{2m})). \end{aligned}$$

Now consider the apartment $\mathcal{A}_G = X_*(S) \otimes \mathbb{R} \cong X_*(T)_\Gamma \otimes \mathbb{R}$, which we identify with $X \otimes \mathbb{R}$ via (5.4.2). We declare that $\widetilde{W}_G$ acts on $\mathcal{A}_G$ via the semidirect product decomposition $N(K_0) = T(K_0) \rtimes S_{2m}^*$ from §3.5, with $T(K_0)$ acting by translations on $X_*(T)_\Gamma$ via the Kottwitz homomorphism, and the permutation matrices S_{2m}^* acting via their natural linear action on $X_*(T)_\Gamma$.

Note that the map (5.4.1) identifies $X_*(S)$ with its image in X ,

$$(5.4.3) \quad X_*(S) \xrightarrow{\sim} \{v \in X \mid \text{the entries of } v \text{ are even}\}.$$

These coordinates on $X_*(S)$ are not the ones given earlier in (3.2.1). From now on we shall work with respect to these new coordinates (5.4.3), and therefore we must re-express the affine roots (3.5.1) in terms of them. We get that the affine roots consist of the functions on $X \otimes \mathbb{R}$

$$(5.4.4) \quad \frac{1}{2}\alpha_{i,j} + \frac{1}{2}\mathbb{Z} \quad \text{for } j \neq i, i^* \quad \text{and} \quad \frac{1}{2}\alpha_{i,i^*} + \frac{1}{2} + \mathbb{Z} \quad \text{for } 1 \leq i \leq 2m.$$

Regarding $X \otimes \mathbb{R}$ as a subspace of $\mathbb{R}^{2m}$, we then take as our base alcove

$$A_G := \left\{ (x_1, \dots, x_{2m}) \in \mathbb{R}^{2m} \mid \begin{array}{l} x_1 + x_{2m} = \dots = x_m + x_{m+1} \text{ and} \\ x_1 < x_2 < x_3 < \dots < x_{m-1} < x_m, x_{m+1} \end{array} \right\}.$$

Of course, this choice of base alcove endows $\widetilde{W}_G$ with a Bruhat order, as discussed in §5.2.

5.5. Faces of type I . The action of $\widetilde{W}_G$ on $\mathcal{A}_G$ described in the previous subsection identifies $\widetilde{W}_G$ with $X \rtimes S_{2m}^*$ inside the group of affine transformations $\text{Aff}(X \otimes \mathbb{R})$. In this way we may regard $\widetilde{W}_G$ as a subgroup of $\widetilde{W}_{2m}$ (4.1.2), and we attach to each $w \in \widetilde{W}_G$ a face $\mathbf{v}$ of type $(2m, \{0, \dots, m\})$ by letting w act on the standard face $\omega_{\{0, \dots, m\}}$, as in §4.2. We write μ_i^w for the vector $\mu_i^{\mathbf{v}}$ attached to this face,

$$(5.5.1) \quad \mu_i^w := w\omega_i - \omega_i \quad \text{for } i \in \mathbb{Z}.$$

For each $\mathcal{O}_K$ -lattice λ in K^{2m} of the form $\sum_{i=1}^{2m} u^{j_i} \mathcal{O}_K e_i$, let

$$T(\lambda) := (j_1, \dots, j_{2m}) \in \mathbb{Z}^{2m}.$$

Then T is equivariant in the sense that for $n \in N(K_0)$,

$$T(n\lambda) = \bar{n} \cdot T(\lambda),$$

where $\bar{n}$ denotes the image of n in $\widetilde{W}_G$, which acts via the action of $\widetilde{W}_{2m}$ on $\mathbb{Z}^{2m}$.

Recalling the lattice λ_i from (3.4.1), we have

$$T(\lambda_i) = \omega_i \quad \text{for all } i \in \mathbb{Z}.$$

The points $\frac{\omega_i + \omega_{-i}}{2}$ are contained in $\overline{A}_G$ for all i , and we conclude that $P_{\{0, \dots, m\}}$ is precisely the Iwahori subgroup of G fixing the alcove A_G in the building. For nonempty $I \subset \{0, \dots, m\}$ satisfying (1.1), we define

$$W_{I,G} := (N(K_0) \cap P_I^\circ) / T(K_0)_1 \subset \widetilde{W}_G.$$

Then, as discussed in §5.1, $W_{I,G}$ is a subgroup of the affine Weyl group inside $\widetilde{W}_G$. It is generated by the reflections across the walls of A_G containing the points $\frac{\omega_i + \omega_{-i}}{2}$ for all $i \in I$.

By definition of P_I° , elements of $W_{I,G}$ fix ω_i for all $i \in 2m\mathbb{Z} \pm I$. Thus to each $w \in \widetilde{W}_G/W_{I,G}$ we get a well-defined face $w \cdot \omega_I$ of type $(2m, I)$ and vector $\mu_i^w = w\omega_i - \omega_i$ for $i \in 2m\mathbb{Z} \pm I$.

5.6. Kottwitz homomorphism on $\widetilde{W}_G$. Let $W_{\text{aff},G}$ denote the affine Weyl group inside $\widetilde{W}_G$. Then the Kottwitz homomorphism κ_G (see §3.3) identifies

$$\widetilde{W}_G/W_{\text{aff},G} \xrightarrow{\sim} \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z}).$$

Explicitly, for $w \in \widetilde{W}_G$, the Kottwitz homomorphism sends

$$(5.6.1) \quad w \mapsto (d, \sum_{i=1}^m \mu_m^w(i) \bmod 2),$$

where w defines a d -face of type $\{0, \dots, m\}$.

5.7. Relation to type B_m . As noted in the Introduction, upon choosing a special vertex in $\mathcal{A}_G$ as origin, there exists a unique reduced root system Σ on $\mathcal{A}_G$ such that the affine root hyperplanes for G are precisely the affine root hyperplanes for Σ , that is, the vanishing loci of the functions $\alpha + \mathbb{Z}$ for $\alpha \in \Sigma$. In this subsection we shall show that Σ is of type B_m . In fact, we shall construct an explicit isomorphism between the triple $(\widetilde{W}_G, \mathcal{A}_G, \mathcal{H}_G)$, where $\mathcal{H}_G$ denotes the system of affine root hyperplanes in $\mathcal{A}_G$, and the analogous triple attached to split GO_{2m+1} .

The point $\omega_m \in X \otimes \mathbb{R}$ is a special vertex for A_G , and we shall begin by transporting it to the origin. Rather than simply translating by $-\omega_m$, let

$$\delta := (1, m+1)(2, m+2) \cdots (m, 2m) \in S_{2m}^*,$$

where (i, j) denotes the transposition interchanging i and j and fixing the other elements of $\{1, \dots, 2m\}$. Of course $\delta^{-1} = \delta$. Let

$$z := \delta t_{-\omega_m} \in \text{Aff}(X \otimes \mathbb{R}).$$

Then

$$(5.7.1) \quad X \otimes \mathbb{R} \xrightarrow{z} X \otimes \mathbb{R}$$

sends $\omega_m \mapsto \mathbf{0}$ and our base alcove A_G to the set

$$(5.7.2) \quad \left\{ (x_1, \dots, x_{2m}) \in \mathbb{R}^{2m} \mid \begin{array}{l} x_1 + x_{2m} = \cdots = x_m + x_{m+1} \text{ and} \\ x_1, x_{2m} - 1 < x_2 < x_3 < \cdots < x_m < x_{m+1} \end{array} \right\}.$$

Each affine root $\tilde{\alpha}$, viewed as a function on the source in (5.7.1), is carried by z to the function on the target

$$z_* \tilde{\alpha} = \tilde{\alpha} \circ z^{-1} = \tilde{\alpha} \circ (t_{\omega_m} \delta).$$

Recalling the explicit form (5.4.4) for the affine roots, we then have

$$z_* \left(\frac{1}{2} \alpha_{i,j} + \frac{d}{2} \right) = \frac{1}{2} \alpha_{\delta(i), \delta(j)} + \frac{\alpha_{i,j}(\omega_m) + d}{2} \quad (j \neq i, i^*, d \in \mathbb{Z})$$

and

$$z_* \left(\frac{1}{2} \alpha_{i,i^*} + \frac{1}{2} + d \right) = \frac{1}{2} \alpha_{\delta(i), \delta(i^*)} + \frac{\alpha_{i,i^*}(\omega_m) + 1}{2} + d \quad (d \in \mathbb{Z}).$$

Since $\alpha_{i,i^*}(\omega_m) = \pm 1$, we conclude that z carries the affine root hyperplanes in the source of (5.7.1) to the zero loci of the functions

$$(5.7.3) \quad \alpha_{i,j} + \mathbb{Z} \quad \text{for } j \neq i, i^* \quad \text{and} \quad \frac{1}{2}\alpha_{i,i^*} + \mathbb{Z} \quad \text{for } 1 \leq i \leq 2m.$$

The functions (5.7.3) are the affine roots for the root system

$$\Sigma := \left\{ \alpha_{i,j}, \frac{1}{2}\alpha_{i,i^*} \mid 1 \leq i, j \leq 2m, j \neq i, i^* \right\}.$$

Now consider the isomorphism

$$\begin{aligned} f: \quad X &\xrightarrow{\sim} X_{*2m+1} \\ (x_1, \dots, x_{2m}) &\longmapsto (x_1, \dots, x_m, c/2, x_{m+1}, \dots, x_{2m}), \end{aligned}$$

where c denotes the common integer $x_1 + x_{2m} = \dots = x_m + x_{m+1}$. The group X_{*2m+1} is the cocharacter group for the root datum $\mathcal{R}_{B_m}$ defined later in §9.2, which is the root datum for split GO_{2m+1} . It is immediate that the roots Φ_{B_m} (9.2.1) for $\mathcal{R}_{B_m}$ pull back under f to the functions Σ on X . Furthermore f induces an isomorphism $S_{2m}^* \xrightarrow{\sim} S_{2m+1}^*$ between groups acting faithfully on source and target, respectfully. Hence f induces an isomorphism, which we still denote f ,

$$f: X \rtimes S_{2m}^* \xrightarrow{\sim} X_{*2m+1} \rtimes S_{2m+1}^*;$$

here the target is the extended affine Weyl group $\widetilde{W}_{B_m}$ of the root datum $\mathcal{R}_{B_m}$. Since $f \otimes \mathbb{R}$ plainly identifies the set (5.7.2) with the alcove A_{B_m} (9.2.2), and since $X \rtimes S_{2m}^*$ is easily seen to be stable under conjugation by z inside $\text{Aff}(X \otimes \mathbb{R})$, we conclude the following.

Lemma 5.7.4. *The composite*

$$X \otimes \mathbb{R} \xrightarrow[\sim]{z} X \otimes \mathbb{R} \xrightarrow[\sim]{f \otimes \mathbb{R}} X_{*2m+1} \otimes \mathbb{R}$$

induces a composite isomorphism

$$\widetilde{W}_G \cong X \rtimes S_{2m}^* \xrightarrow[\sim]{\text{int}(z)} X \rtimes S_{2m}^* \xrightarrow[\sim]{f} X_{*2m+1} \rtimes S_{2m+1}^* = \widetilde{W}_{B_m}$$

which identifies the A_G -Bruhat order on $\widetilde{W}_G$ with the A_{B_m} -Bruhat order on $\widetilde{W}_{B_m}$. Here $\text{int}(z)$ is the conjugation map $w \mapsto zwz^{-1}$. $\square$

The lemma immediately allows us to transfer all questions we shall have about $\widetilde{W}_G$ to $\widetilde{W}_{B_m}$. To do so in an effective way, we shall need to explicitly address the intervention of $\text{int}(z)$ in the displayed isomorphism. The following will take care of everything we need.

Lemma 5.7.5.

- (i) $z\omega_i = \omega_{i-m}$ for all $i \in \mathbb{Z}$.
- (ii) $\mu_i^{zwz^{-1}} = \delta\mu_{i+m}^w$ for all $i \in \mathbb{Z}$ and $w \in \widetilde{W}_G$.

Proof. The first assertion is an exercise, and the second is an easy consequence of the first. $\square$

It follows easily from the first part of the lemma that, for any nonempty $I \subset \{0, \dots, m\}$ satisfying (1.1), the composite isomorphism $\widetilde{W}_G \xrightarrow{\sim} \widetilde{W}_{B_m}$ in (5.7.4) sends the subgroup $W_{I,G}$ isomorphically to the subgroup W_{m-I, B_m} defined in (9.4.1).

5.8. The $\{\mu_{r,s}\}$ -admissible set. Recalling our identification $X_*(T) \simeq \mathbb{Z}^{2m} \times \mathbb{Z}$ from §3.2, let

$$\mu_{r,s} := (1^{(s)}, 0^{(2m-s)}, 1) \in X_*(T),$$

and let

$$\{\mu_{r,s}\} := S_{2m} \cdot \mu_{r,s}$$

denote the geometric conjugacy class of $\mu_{r,s}$, where S_{2m} acts by permuting the first $2m$ entries. The image of $\{\mu_{r,s}\}$ in X under the map (5.4.1) is the set

$$\{\bar{\mu}_{r,s}\} = S_{2m}^* \cdot (2^{(s)}, 1^{(2m-2s)}, 0^{(s)}) \amalg S_{2m}^* \cdot (2^{(s-2)}, 1^{(2m-2s+4)}, 0^{(s-2)}) \amalg \dots,$$

where the union terminates with $\{1\}$ or $S_{2m}^* \cdot (2, 1^{(2m-1)}, 0)$ according as s is even or odd. Regarding $\{\bar{\mu}_{r,s}\}$ as a subset of the translation subgroup of $\widetilde{W}_G$, the image of $\{\bar{\mu}_{r,s}\}$ in $\widetilde{W}_{B_m}$ under the isomorphism in (5.7.4) consists of the translation elements

$$S_{2m+1}^* \cdot (2^{(s)}, 1^{(2m-2s+1)}, 0^{(s)}) \amalg S_{2m+1}^* \cdot (2^{(s-2)}, 1^{(2m-2s+5)}, 0^{(s-2)}) \amalg \dots$$

These elements are all congruent modulo coroots, and the maximal elements among them in the Bruhat order are the elements $S_{2m+1}^* \cdot (2^{(s)}, 1^{(2m-2s+1)}, 0^{(s)})$. Hence the elements $S_{2m}^* \cdot (2^{(s)}, 1^{(2m-2s)}, 0^{(s)})$ are maximal in $\{\bar{\mu}_{r,s}\}$. This confirms, for G and the conjugacy class $\{\mu_{r,s}\}$, the conjecture [PRS, Conj. 4.5.3] we discussed in §5.3, since in this case the set $\Lambda_{\{\mu_{r,s}\}}$ is by definition the image of $S_{2m}^* \cdot \mu_{r,s}$ in the coinvariants.

Now let $\text{Adm}_{J,I,G}(\{\mu_{r,s}\})$ denote the $\{\mu_{r,s}\}$ -admissible set in $W_{J,G} \backslash \widetilde{W}_G / W_{I,G}$, and for any cocharacter $\mu \in X_{*2m+1}$, let $\text{Adm}_{J,I,B_m}(\mu)$ denote the μ -admissible set in $W_{J,B_m} \backslash \widetilde{W}_{B_m} / W_{I,B_m}$. We have shown the following.

Lemma 5.8.1. *For any nonempty subsets $I, J \subset \{0, \dots, m\}$ satisfying property (1.1), the isomorphism $W_{J,G} \backslash \widetilde{W}_G / W_{I,G} \xrightarrow{\sim} \widetilde{W}_{m-J,B_m} \backslash \widetilde{W}_{B_m} / W_{m-I,B_m}$ induced by (5.7.4) identifies*

$$\text{Adm}_{J,I,G}(\{\mu_{r,s}\}) \xrightarrow{\sim} \text{Adm}_{m-J,m-I,B_m}(\mu),$$

where μ is the cocharacter $(2^{(s)}, 1^{(2m-2s+1)}, 0^{(s)}) \in X_{*2m+1}$. $\square$

6. AFFINE FLAG VARIETY

From now on we take $K_0 = k((t))$, $\mathcal{O}_{K_0} = k[[t]]$, $K = k((u))$, and $\mathcal{O}_K = k[[u]]$, where we recall that k is an algebraic closure of the common residue field of F_0 and F . In this section we review the affine flag variety attached to a parahoric subgroup of G , and the embedding of the geometric special fiber of the naive local model into it, from [PR4, §3]. Let $I \subset \{0, \dots, m\}$ be a nonempty subset satisfying property (1.1).

6.1. Affine flag variety. Let P be a parahoric subgroup of $G(K_0)$. Then Bruhat-Tits theory attaches to P a smooth affine $\mathcal{O}_{K_0}$ -group scheme whose generic fiber identifies with G , whose special fiber is connected, and whose group of $\mathcal{O}_{K_0}$ -points identifies with P ; abusing notation, we denote this group scheme again by P . The *affine flag variety $\mathcal{F}_P$ relative to P* is the fpqc quotient of functors on the category of k -algebras,

$$\mathcal{F}_P := LG / L^+P,$$

where LG is the *loop group* $LG: R \mapsto G(R((t)))$ and L^+P is the *positive loop group* $L^+P: R \mapsto G(R[[t]])$. See [PR3]. The affine flag variety is an ind- k -scheme of ind-finite type.

6.2. Lattice-theoretic description. In this subsection we give a slight variant of (and make a minor correction to) the description in [PR4, §3.2] of the affine flag variety in terms of lattice chains in K^{2m} .

Let R be a k -algebra. Recall that an $R[[u]]$ -lattice in $R((u))^{2m}$ is an $R[[u]]$ -submodule $L \subset R((u))^{2m}$ which is free as an $R[[u]]$ -module Zariski-locally on $\text{Spec } R$, and such that the natural arrow $L \otimes_{R[[u]]} R((u)) \rightarrow R((u))^{2m}$ is an isomorphism. Borrowing notation from (2.1.3), given an $R[[u]]$ -lattice L , we write $\widehat{L}$ for the dual lattice

$$\widehat{L} := \{ x \in R((u))^{2m} \mid h_{R((u))}(L, x) \subset R[[u]] \},$$

where $h_{R((u))} := h \otimes_K R((u))$ is the induced form on $R((u))^{2m}$. A collection $\{L_i\}_i$ of $R[[u]]$ -lattices in $R((u))^{2m}$ is a *chain* if it is totally ordered under inclusion and all successive quotients are locally free R -modules (necessarily of finite rank). A lattice chain is *periodic* if uL is in the chain for every lattice L in the chain. We write $\mathcal{L}(R((u))^{2m})$ for the category whose objects are the $R[[u]]$ -lattices in $R((u))^{2m}$ and whose morphisms are the natural inclusions of lattices. Of course, any $R[[u]]$ -lattice chain may be regarded as a full subcategory of $\mathcal{L}(R((u))^{2m})$.

We define $\mathcal{F}_I$ to be the functor on k -algebras that assigns to each R the set of all functors $L: \lambda_I \rightarrow \mathcal{L}(R((u))^{2m})$ such that

- (C) (chain) the image $L(\lambda_I)$ is a lattice chain in $R((u))^{2m}$;
- (P) (periodicity) $L(u\lambda_i) = uL(\lambda_i)$ for all $i \in 2m\mathbb{Z} \pm I$, so that the chain $L(\lambda_I)$ is periodic;
- (R) (rank) $\dim_k \lambda_i / \lambda_j = \text{rank}_R L(\lambda_i) / L(\lambda_j)$ for all $j < i$; and
- (D) (duality) Zariski-locally on $\text{Spec } R$, there exists $\alpha \in R((t))^\times \subset R((u))^\times$ such that $\widehat{L(\lambda_i)} = \alpha L(\widehat{\lambda_i})$ for all $i \in 2m\mathbb{Z} \pm I$.

If $L \in \mathcal{F}_I(R)$ globally admits a scalar α as in (D), then α is well-defined modulo the group

$$\{ a \in R((t))^\times \mid aL(\lambda_{-i}) = L(\lambda_{-i}) \} = R[[t]]^\times$$

(independent of i). It is then an easy exercise to check that the map specified locally by

$$L \longmapsto \left((L(\lambda_i))_{i \in I}, \alpha \bmod R[[t]]^\times \right)$$

defines an isomorphism from the functor $\mathcal{F}_I$ as we've defined it to the functor $\mathcal{F}_I$ as defined in [PR4, §3.2], except that the functor in [PR4] should only require that $\alpha \bmod R[[t]]^\times$ be given Zariski-locally.

Now recall the subgroup $P_I \subset G(K_0)$, which is defined to be the lattice-wise stabilizer of the chain λ_I ; it equals the parahoric subgroup P_I° when $m \in I$, contains P_I° as a subgroup of index 2 when $m \notin I$, and in all cases is the full stabilizer in $G(K_0)$ of the facet corresponding to P_I° . As for parahoric subgroups, Bruhat-Tits theory attaches to P_I a canonical smooth affine group scheme over $\mathcal{O}_{K_0}$ with generic fiber G and whose $\mathcal{O}_{K_0}$ points identify with P_I ; abusing notation as in §6.1, we continue to denote this group scheme by P_I . We may then consider the positive loop group L^+P_I , which assigns each k -algebra R to the lattice-wise stabilizer in $G(R((t)))$ of the lattice chain $\lambda_I \otimes_{\mathcal{O}_K} R[[u]]$.

The loop group LG acts on $\mathcal{F}_I$ via the natural representation of $G(R((t)))$ on $R((u))^{2m}$, and it follows that the LG -equivariant map $LG \rightarrow \mathcal{F}_I$ specified by taking the tautological inclusion $(\lambda_I \hookrightarrow \mathcal{L}(K^{2m})) \in \mathcal{F}_I(k)$ as basepoint defines an LG -equivariant isomorphism

$$(6.2.1) \quad LG/L^+P_I \xrightarrow{\sim} \mathcal{F}_I.$$

When $m \in I$, this identifies $\mathcal{F}_I$ with the affine flag variety $\mathcal{F}_{P_I^\circ}$. When $m \notin I$, we recall that the Kottwitz homomorphism

$$\kappa_G: G(K_0) \longrightarrow \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})$$

extends to a homomorphism of functors

$$\kappa_G: LG \longrightarrow \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z});$$

here the target even classifies the connected components of LG [PR3, Th. 0.1]. Let H denote the kernel of the map to the second factor,

$$H := \ker[LG \longrightarrow \mathbb{Z}/2\mathbb{Z}],$$

and let τ denote the block diagonal matrix in $LG(k)$

$$(6.2.2) \quad \tau := \begin{pmatrix} \text{Id}_{m-1} & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & \text{Id}_{m-1} \end{pmatrix}.$$

Then $LG = H \amalg \tau H$, and

$$(6.2.3) \quad \mathcal{F}_{P_I^\circ} = (H/L^+P_I^\circ) \amalg (\tau H/L^+P_I^\circ) \xrightarrow{\sim} LG/L^+P_I \amalg LG/L^+P_I \xrightarrow{\sim} \mathcal{F}_I \amalg \mathcal{F}_I.$$

6.3. Schubert cells and varieties. Consider the parahoric subgroup scheme P_I° over $\mathcal{O}_{K_0}$ and its associated affine flag variety $\mathcal{F}_{P_I^\circ}$. For $n \in N(K_0)$, the associated *Schubert cell* is the reduced k -subscheme

$$L^+P_I^\circ \cdot n \subset \mathcal{F}_{P_I^\circ}.$$

The Schubert cell depends only on the image w of n in $W_{I,G} \backslash \widetilde{W}_G / W_{I,G}$, and we denote the Schubert cell by C_w . By Haines–Rapoport [HR, Prop. 8], the inclusion $N(K_0) \subset G(K_0)$ induces a bijection

$$W_{I,G} \backslash \widetilde{W}_G / W_{I,G} \xrightarrow{\sim} P_I^\circ(\mathcal{O}_{K_0}) \backslash G(K_0) / P_I^\circ(\mathcal{O}_{K_0}),$$

so that the Schubert cells are indexed by precisely the elements of $W_{I,G} \backslash \widetilde{W}_G / W_{I,G}$ and give a stratification of all of $\mathcal{F}_{P_I^\circ}$. Note that in the special case $I = \{0, \dots, m\}$, P_I° is an Iwahori subgroup, the group $W_{I,G}$ is trivial, and the Schubert cells are indexed by $\widetilde{W}_G$ itself.

For $w \in W_{I,G} \backslash \widetilde{W}_G / W_{I,G}$, the associated *Schubert variety* S_w is the reduced closure of C_w in $\mathcal{F}_{P_I^\circ}$. The closure relations between Schubert cells are given by the Bruhat order: for $w, w' \in W_{I,G} \backslash \widetilde{W}_G / W_{I,G}$, we have $S_w \subset S_{w'}$ in $\mathcal{F}_{P_I^\circ} \iff w \leq w'$ in $W_{I,G} \backslash \widetilde{W}_G / W_{I,G}$. See Richarz [R, Prop. 2.8].

6.4. Embedding the geometric special fiber. We now recall from [PR4, §3.3] the embedding of the geometric special fiber $M_{I,k}^{\text{naive}} := M_I^{\text{naive}} \otimes_{\mathcal{O}_E} k$ of the naive local model into the affine flag variety $\mathcal{F}_{P_I^\circ}$.

The embedding makes use of the lattice-theoretic description of the affine flag variety in terms of $\mathcal{F}_I$ from §6.2. First note that the $\mathcal{O}_K$ -lattice chain λ_I admits a trivialization in obvious analogy with the trivialization of Λ_I specified by (2.2.2), where λ_i replaces Λ_i , $\mathcal{O}_K$ replaces $\mathcal{O}_F$, and u replaces π . Upon identifying $\mathcal{O}_K \otimes_{\mathcal{O}_{K_0}} k \xrightarrow{\sim} \mathcal{O}_F \otimes_{\mathcal{O}_{F_0}} k$ by sending the k -basis elements $1 \otimes 1 \mapsto 1 \otimes 1$ and $u \otimes 1 \mapsto \pi \otimes 1$, our trivializations then yield an identification of chains of k -vector spaces

$$(6.4.1) \quad \lambda_i \otimes_{\mathcal{O}_{K_0}} k \simeq \Lambda_i \otimes_{\mathcal{O}_{F_0}} k;$$

this is even an isomorphism of $k[u]/(u^2)$ -modules, where u acts on the right-hand side as multiplication by $\pi \otimes 1$.

Now let R be a k -algebra. Given an R -point $\{\mathcal{F}_i\}_i$ in $M_{I,k}^{\text{naive}}$, for each i , let $L_i \subset \lambda_i \otimes_{\mathcal{O}_{K_0}} R[[t]]$ denote the inverse image of

$$\mathcal{F}_i \subset \Lambda_i \otimes_{\mathcal{O}_{F_0}} R \simeq \lambda_i \otimes_{\mathcal{O}_{K_0}} R$$

under the reduction-mod- t -map

$$\lambda_i \otimes_{\mathcal{O}_{K_0}} R[[t]] \twoheadrightarrow \lambda_i \otimes_{\mathcal{O}_{K_0}} R.$$

Then L_i is naturally a lattice in $R((u))^{2m}$, and the functor $\lambda_I \rightarrow \mathcal{L}(R((u))^{2m})$ sending $\lambda_i \mapsto L_i$ determines a point in $\mathcal{F}_I(R)$. In this way we get a monomorphism

$$(6.4.2) \quad \begin{array}{ccc} M_{I,k}^{\text{naive}} & \hookrightarrow & \mathcal{F}_I \\ \{\mathcal{F}_i\}_i & \longmapsto & (\lambda_i \mapsto L_i) \end{array}.$$

Since $M_{I,k}^{\text{naive}}$ is proper, (6.4.2) is a closed immersion of ind-schemes. When $m \in I$, this embeds $M_{I,k}^{\text{naive}}$ in $\mathcal{F}_{P_I^\circ} = \mathcal{F}_{P_I}$ via the isomorphism (6.2.1). When $m \notin I$, $\mathcal{F}_{P_I^\circ}$ is a disjoint union of two copies of $\mathcal{F}_I$ via (6.2.3), and we use (6.4.2) to embed $M_{I,k}^{\text{naive}}$ in the copy marked by $t_{\mu_{r,s}} \in \widetilde{W}_G$. From now on we shall often identify $M_{I,k}^{\text{naive}}$ with its image in $\mathcal{F}_{P_I^\circ}$.

The embedding $M_{I,k}^{\text{naive}} \hookrightarrow \mathcal{F}_{P_I^\circ}$ is $L^+P_I^\circ$ -equivariant with respect to the following left $L^+P_I^\circ$ -actions on source and target; compare [PR1, §3; PR2, §6, §11; PR3, §11; PR4, §3.3; Sm3, §4.4]. On $\mathcal{F}_{P_I^\circ}$ we just take the natural left $L^+P_I^\circ$ -action. For $M_{I,k}^{\text{naive}}$, $L^+P_I^\circ$ acts naturally on $\lambda_I \otimes_{\mathcal{O}_{K_0}} k$ via the tautological action of L^+P_I on λ_I . The chain isomorphism (6.4.1) then induces a homomorphism

$$L^+P_I^\circ \longrightarrow \underline{\text{Aut}}(\Lambda_I)_{\mathcal{O}_E} \otimes_{\mathcal{O}_E} k,$$

where we recall the $\mathcal{O}_E$ -group scheme $\underline{\text{Aut}}(\Lambda_I)_{\mathcal{O}_E}$ from §2.5. The $\underline{\text{Aut}}(\Lambda_I)_{\mathcal{O}_E}$ -action on M_I^{naive} now furnishes the desired $L^+P_I^\circ$ -action on $M_{I,k}^{\text{naive}}$. In this way $L^+P_I^\circ$ also acts on $M_{I,k}^\wedge$, $M_{I,k}^{\text{spin}}$, and $M_{I,k}^{\text{loc}}$.

For nonempty $J \subset I \subset \{0, \dots, m\}$ satisfying (1.1), we get a diagram

$$\begin{array}{ccc} M_{I,k}^{\text{naive}} & \hookrightarrow & \mathcal{F}_{P_I^\circ} \\ \downarrow & & \downarrow \\ M_{J,k}^{\text{naive}} & \hookrightarrow & \mathcal{F}_{P_J^\circ}, \end{array}$$

where the vertical arrows are the natural projections and the horizontal arrows are the embeddings just defined. If $m \in J$ or $m \notin I$, then this diagram commutes, but if $m \in I \setminus J$, then it does not, as will follow from our description of the Schubert varieties in $M_{I,k}^{\text{naive}}$ and $M_{J,k}^{\text{naive}}$ in the next section. On the other hand, if we replace the geometric special fiber of the naive local model with that of the spin local model everywhere, then the diagram does commute for all nonempty $J \subset I$ satisfying (1.1).

7. SCHUBERT CELLS IN THE GEOMETRIC SPECIAL FIBER

We continue to take I to be a nonempty subset of $\{0, \dots, m\}$ satisfying property (1.1), and to identify $M_{I,k}^{\text{naive}}$ with its image in the affine flag variety $\mathcal{F}_{P_I^\circ}$ under the embedding described in §6.4. It follows from $L^+P_I^\circ$ -equivariance that the underlying topological spaces of $M_{I,k}^{\text{naive}}$, $M_{I,k}^\wedge$, $M_{I,k}^{\text{spin}}$, and $M_{I,k}^{\text{loc}}$ are all unions of Schubert varieties in $\mathcal{F}_{P_I^\circ}$. In this subsection we shall describe the Schubert varieties that are contained in each, and we shall prove Theorems 1.3 and 1.4 modulo the main combinatorial results of the next two sections. We write $i^* := 2m + 1 - i$.

7.1. The image of the geometric special fiber. Let R be a k -algebra. It is clear from the definition of the embedding $M_{I,k}^{\text{naive}} \hookrightarrow \mathcal{F}_I$ (6.4.2) and the various conditions in the definition of M_I^{naive} that the image of $M_{I,k}^{\text{naive}}(R)$ in $\mathcal{F}_I(R)$ consists precisely of the functors $\lambda_i \mapsto L_i$ in $\mathcal{F}_I(R)$ such that, for all $i \in 2m\mathbb{Z} \pm I$,

- (1) $\lambda_{i,R[[t]]} \supset L_i \supset t\lambda_{i,R[[t]]}$, where $\lambda_{i,R[[t]]} := \lambda_i \otimes_{\mathcal{O}_{K_0}} R[[t]]$;
- (2) the R -module $\lambda_{i,R[[t]]}/L_i$ is locally free of rank $2m$; and
- (3) $\hat{L}_i = t^{-1}L_{-i}$.

We remark that condition (3) is actually redundant; that is, for any functor $\lambda_I \rightarrow \mathcal{L}(R((u))^{2m})$ satisfying conditions (C), (P), (R), and (D) from §6.2 and conditions (1) and (2) above, the scalar α appearing in (D) must be congruent to $t^{-1} \bmod R[[t]]^\times$. Since we won't need to use this fact anywhere in the paper, we leave the details as an exercise to the reader.

7.2. Naive permissibility. Let $w \in W_{I,G} \backslash \widetilde{W}_G / W_{I,G}$. Then the condition that the Schubert variety S_w be contained in $M_{I,k}^{\text{naive}}$ (resp. $M_{I,k}^\wedge$, resp. $M_{I,k}^{\text{spin}}$) amounts to the condition that for one, hence any, representative $\tilde{w}$ of w in $\widetilde{W}_G / W_{I,G}$, the point $\tilde{w} \cdot \lambda_I \in \mathcal{F}_I(k)$ be contained in $M_{I,k}^{\text{naive}}(k)$ (resp. $M_{I,k}^\wedge(k)$, resp. $M_{I,k}^{\text{spin}}(k)$) and, in case $m \notin I$, that $\tilde{w}$ mark the same copy of $\mathcal{F}_I$ inside $\mathcal{F}_{P_I^\circ}$ as $t_{\mu_{r,s}}$ does. We shall find it convenient to express these conditions in terms of faces of type I , beginning in this subsection with containment in $M_{I,k}^{\text{naive}}$.

Definition 7.2.1. Let $\tilde{w} \in \widetilde{W}_G / W_{I,G}$. We say that $\tilde{w}$ is *naively permissible* if the Schubert variety in $\mathcal{F}_{P_I^\circ}$ attached to $\tilde{w}$ is contained in $M_{I,k}^{\text{naive}}$.

To translate the notion of naive permissibility into combinatorics, let $\tilde{w} \in \widetilde{W}_G / W_{I,G}$, and consider the face $(\tilde{w}\omega_i)_{i \in 2m\mathbb{Z} \pm I}$ of type $(2m, I)$ attached to $\tilde{w}$, as in §5.5. Then it is clear from the definitions and from §7.1 that $\tilde{w}$ is naively permissible $\iff$

- (1) $\omega_i \leq \tilde{w}\omega_i \leq \omega_i + \mathbf{2}$ for all $i \in 2m\mathbb{Z} \pm I$;
- (2) $\Sigma(\tilde{w}\omega_i) = 2m - i$ for one, hence every, $i \in 2m\mathbb{Z} \pm I$; and

- (3) if $m \notin I$, then $\tilde{w}$ and $t_{\mu_{r,s}} W_{I,G}$ have the same image under the composite map

$$\widetilde{W}_G/W_{I,G} \xrightarrow{\kappa_G} \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z}) \xrightarrow{\text{pr}_{\mathbb{Z}/2\mathbb{Z}}} \mathbb{Z}/2\mathbb{Z}.$$

Recasting this in terms of the vector $\mu_i^{\tilde{w}} = \tilde{w}\omega_i - \omega_i$ (5.5.1), we obtain the following.

Proposition 7.2.2. $\tilde{w} \in \widetilde{W}_G/W_{I,G}$ is naively permissible $\iff$

- (P1) $\mu_i^{\tilde{w}} + (\mu_{-i}^{\tilde{w}})^* = \mathbf{2}$ and $\mathbf{0} \leq \mu_i^{\tilde{w}} \leq \mathbf{2}$ for all $i \in I$, and if $m \notin I$, then furthermore $\tilde{w} \equiv t_{\mu_{r,s}} \pmod{W_{\text{aff},G}}$.

Proof. Note that condition (2) above just says that $\tilde{w}$ defines a 2-face. In light of the explicit description of the Kottwitz homomorphism (5.6.1), the implication $\implies$ is then clear, and the implication $\impliedby$ is clear from (4.3.2) and (4.3.4). $\square$

Given a naively permissible $\tilde{w}$, the point $\tilde{w} \cdot \lambda_I$ in $\mathcal{F}_I(k)$ corresponds to a point $(\mathcal{F}_i \subset \Lambda_i \otimes_{\mathcal{O}_{F_0}} k)_{i \in 2m\mathbb{Z} \pm I} \in M_{I,k}^{\text{naive}}(k)$ of a rather special sort. Namely, identifying $\Lambda_i \otimes_{\mathcal{O}_{F_0}} k$ with $\mathcal{O}_F^{2m} \otimes_{\mathcal{O}_{F_0}} k$ via (2.2.2), we have

- (S) for all i , $\mathcal{F}_i$, regarded as a subspace in $\mathcal{O}_F^{2m} \otimes_{\mathcal{O}_{F_0}} k$, is k -spanned by $2m$ of the elements $\epsilon_1 \otimes 1, \dots, \epsilon_{2m} \otimes 1, \pi\epsilon_1 \otimes 1, \dots, \pi\epsilon_{2m} \otimes 1$,

where we recall that $\epsilon_1, \dots, \epsilon_{2m}$ denotes the standard $\mathcal{O}_F$ -basis in $\mathcal{O}_F^n$. On the other hand, for any point $(\mathcal{F}_i)_i$ in $M_{I,k}^{\text{naive}}(k)$, let us say that $(\mathcal{F}_i)_i$ is an S -fixed point if it satisfies (S); it is easy to check that the S -fixed points are exactly the points in $M_k^{\text{naive}}(k)$ fixed by $L^+S(k)$. In this way, we get a bijection between the naively permissible $\tilde{w} \in \widetilde{W}_G/W_{I,G}$ and the S -fixed points in $M_{I,k}^{\text{naive}}(k)$, which we denote by $\tilde{w} \mapsto (\mathcal{F}_i^{\tilde{w}})_i$. Explicitly, for naively permissible $\tilde{w}$, in terms of the vector $\mu_i^{\tilde{w}}$, we have

$$(7.2.3) \quad \mathcal{F}_i^{\tilde{w}} = \sum_{\mu_i^{\tilde{w}}(j)=0} k \cdot (\epsilon_j \otimes 1) + \sum_{\mu_i^{\tilde{w}}(j)=0,1} k \cdot (\pi\epsilon_j \otimes 1) \subset \mathcal{O}_F^{2m} \otimes_{\mathcal{O}_{F_0}} k.$$

7.3. Wedge-permissibility. We continue with the notation of the previous subsection.

Definition 7.3.1. We say that $\tilde{w} \in \widetilde{W}_G/W_{I,G}$ is *wedge-permissible* if the Schubert variety in $\mathcal{F}_{P_I^\circ}$ attached to $\tilde{w}$ is contained in $M_{I,k}^\wedge$.

Of course wedge-permissibility depends only on the image of $\tilde{w}$ in $W_{I,G} \backslash \widetilde{W}_G/W_{I,G}$. By definition, for naively permissible $\tilde{w}$, we have

$$(\mathcal{F}_i^{\tilde{w}})_i \in M_{I,k}^\wedge(k) \iff \begin{aligned} &\text{for all } i \in 2m\mathbb{Z} \pm I, \bigwedge_k^{s+1}(\pi \otimes 1 \mid \mathcal{F}_i^{\tilde{w}}) = 0 \\ &\text{and } \bigwedge_k^{r+1}(\pi \otimes 1 \mid \mathcal{F}_i^{\tilde{w}}) = 0, \end{aligned}$$

where we recall our fixed partition $2m = s + r$ with $s < r$. For fixed i , the second equality on the right-hand side of the display is implied by the first. We then have the following.

Proposition 7.3.2. $\tilde{w} \in \widetilde{W}_G/W_{I,G}$ is wedge-permissible $\iff \tilde{w}$ is naively permissible and

- (P2) for all $i \in I$, $\#\{j \mid \mu_i^{\tilde{w}}(j) = 0\} \leq s$. $\square$

Proof. The implication $\implies$ is immediate from the description of $\mathcal{F}_i^{\tilde{w}}$ (7.2.3), and the implication $\impliedby$ follows from this, (4.3.2), (4.3.4), and (4.4.3). $\square$

7.4. Spin-permissibility. We continue with the notation of the previous subsections. We also freely use the notation of §2.4, and we recall the sets A_i and B_i (taken with $n = 2m$) from (4.3.6).

Definition 7.4.1. We say that $\tilde{w} \in \widetilde{W}/W_{I,G}$ is *spin-permissible* if the Schubert variety in $\mathcal{F}_{P_I^\circ}$ attached to $\tilde{w}$ is contained in $M_{I,k}^{\text{spin}}$.

As for naive and wedge-permissibility, spin-permissibility depends only on the image of $\tilde{w}$ in $W_{I,G} \backslash \widetilde{W}_G/W_{I,G}$. Our aim in this subsection is to characterize the set of spin-permissible $\tilde{w}$ in terms of the vector $\mu_i^{\tilde{w}}$, for $i \in 2m\mathbb{Z} \pm I$.

Let $i \in I$, and suppose that $\tilde{w}$ is wedge-permissible. Continuing to identify $\mathcal{O}_F^{2m}$ with Λ_i (resp. Λ_{2m-i}) via (2.2.2), the $2m$ elements in $\mathcal{O}_F^{2m}$

$$(7.4.2) \quad \begin{aligned} \epsilon_j & \text{ for } \mu_i^{\tilde{w}}(j) = 0 \quad (\text{resp. } \mu_{2m-i}^{\tilde{w}}(j) = 0); \quad \text{and} \\ \pi\epsilon_j & \text{ for } \mu_i^{\tilde{w}}(j) = 0, 1 \quad (\text{resp. } \mu_{2m-i}^{\tilde{w}}(j) = 0, 1) \end{aligned}$$

span an $\mathcal{O}_{F_0}$ -submodule whose image under the reduction-mod- π_0 map

$$\mathcal{O}_F^{2m} \twoheadrightarrow \mathcal{O}_F^{2m} \otimes_{\mathcal{O}_{F_0}} k$$

is $\mathcal{F}_i^{\tilde{w}}$ (resp. $\mathcal{F}_{2m-i}^{\tilde{w}}$). Take the wedge product (in any order) of the elements (7.4.2) in $\bigwedge_{\mathcal{O}_{F_0}}^{2m} \mathcal{O}_F^{2m}$, and let $g_i \in \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_i$ (resp. $g_{2m-i} \in \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_{2m-i}$) denote the image of this element under the isomorphism $\bigwedge_{\mathcal{O}_{F_0}}^{2m} \mathcal{O}_F^{2m} \xrightarrow{\sim} \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_i$ (resp. $\bigwedge_{\mathcal{O}_{F_0}}^{2m} \mathcal{O}_F^{2m} \xrightarrow{\sim} \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_{2m-i}$) induced by (2.2.2). Then, up to sign, and in terms of the notation (2.4.1), g_i equals

$$\pi_0^{a_i} f_{E_i} \in \bigwedge_{F_0}^{2m} V,$$

where $E_i \subset \{1, \dots, 4m\}$ is the subset of cardinality $2m$

$$(7.4.3) \quad \begin{aligned} E_i &:= \{j \in \{1, \dots, i\} \mid \mu_i^{\tilde{w}}(j) = 0\} \\ &\quad \amalg \{j \in \{i+1, \dots, m\} \mid \mu_i^{\tilde{w}}(j) = 0, 1\} \\ &\quad \amalg \{j \in \{m+1, \dots, 2m\} \mid \mu_i^{\tilde{w}}(j) = 0\} \\ &\quad \amalg \{2m+j \in \{2m+1, \dots, 2m+i\} \mid \mu_i^{\tilde{w}}(j) = 0, 1\} \\ &\quad \amalg \{2m+j \in \{2m+i+1, \dots, 3m\} \mid \mu_i^{\tilde{w}}(j) = 0\} \\ &\quad \amalg \{2m+j \in \{3m+1, \dots, 4m\} \mid \mu_i^{\tilde{w}}(j) = 0, 1\}, \end{aligned}$$

and

$$a_i := \#(E_i \cap \{i+1, \dots, m\}) = \#\{j \in \{i+1, \dots, m\} \mid \mu_i^{\tilde{w}}(j) = 0, 1\};$$

and again up to sign, g_{2m-i} equals

$$\pi_0^{a_{2m-i}} f_{E_{2m-i}} \in \bigwedge_{F_0}^{2m} V,$$

where $E_{2m-i} \subset \{1, \dots, 4m\}$ is the subset of cardinality $2m$

$$\begin{aligned} E_{2m-i} &:= \{j \in \{1, \dots, m\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0\} \\ &\quad \amalg \{j \in \{m+1, \dots, 2m-i\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0, 1\} \\ &\quad \amalg \{j \in \{i^*, \dots, 2m\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0\} \\ &\quad \amalg \{2m+j \in \{2m+1, \dots, 3m\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0, 1\} \end{aligned}$$

$$\begin{aligned} & \Pi \{ 2m + j \in \{3m + 1, \dots, 4m - i\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0 \} \\ & \Pi \{ 2m + j \in \{2m + i^*, \dots, 4m\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0, 1 \}, \end{aligned}$$

and

$$\begin{aligned} a_{2m-i} &:= -\#(E_{2m-i} \cap \{3m + 1, \dots, 4m - i\}) \\ &= -\#\{j \in \{m + 1, \dots, 2m - i\} \mid \mu_{2m-i}^{\tilde{w}}(j) = 0\}. \end{aligned}$$

To study the spin condition for $\mathcal{F}_i^{\tilde{w}}$, we shall also need the set $E_i^\perp = (4m + 1 - E_i)^c$, which is given by

$$\begin{aligned} E_i^\perp &= \{j \in \{1, \dots, m\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0, 1\} \\ &\quad \Pi \{j \in \{m + 1, \dots, 2m - i\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0\} \\ &\quad \Pi \{j \in \{i^*, \dots, 2m\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0, 1\} \\ &\quad \Pi \{2m + j \in \{2m + 1, \dots, 3m\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0\} \\ &\quad \Pi \{2m + j \in \{3m + 1, \dots, 4m - i\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0, 1\} \\ &\quad \Pi \{2m + j \in \{2m + i^*, \dots, 4m\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0\}. \end{aligned}$$

We then have the following.

Lemma 7.4.4.

- (i) $E_i^\perp = E_{2m-i}$.
- (ii) Let $a_i^\perp$ (resp. $a_{2m-i}^\perp$) denote the unique integer such that

$$\pi_0^{a_i^\perp} f_{E_i^\perp} \in \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_i \setminus \pi_0 \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_i$$

(resp.

$$\pi_0^{a_{2m-i}^\perp} f_{E_{2m-i}^\perp} \in \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_{2m-i} \setminus \pi_0 \bigwedge_{\mathcal{O}_{F_0}}^{2m} \Lambda_{2m-i}).$$

Then

$$\begin{aligned} a_i^\perp &= \#(E_i^\perp \cap \{i + 1, \dots, m\}) \\ &= \#\{j \in \{i + 1, \dots, m\} \mid \mu_i^{\tilde{w}}(j^*) \neq 0, 1\} \end{aligned}$$

and

$$\begin{aligned} a_{2m-i}^\perp &= -\#(E_{2m-i}^\perp \cap \{3m + 1, \dots, 4m - i\}) \\ &= -\#\{j \in \{m + 1, \dots, 2m - i\} \mid \mu_{2m-i}^{\tilde{w}}(j^*) \neq 0\}. \end{aligned}$$

- (iii) $a_i \geq a_i^\perp$ and $a_{2m-i} \geq a_{2m-i}^\perp$.
- (iv) $a_i - a_{2m-i} = a_i^\perp - a_{2m-i}^\perp = m - i$.

Proof. Assertion (i) follows immediately from our explicit expressions for E_{2m-i} and $E_i^\perp$, using that $\mu_i^{\tilde{w}} + (\mu_{2m-i}^{\tilde{w}})^* = \mathbf{2}$ and $\mathbf{0} \leq \mu_i^{\tilde{w}} \leq \mathbf{2}$ by naive permissibility. Assertion (ii) is clear. Assertion (iii) follows from our explicit expressions for the quantities involved, naive permissibility, and the basic inequalities (4.3.7). To prove (iv), by naive permissibility we have

$$a_{2m-i} = -\#\{j \in \{i + 1, \dots, m\} \mid \mu_i^{\tilde{w}}(j) = 2\}.$$

Hence

$$a_i - a_{2m-i} = \#\{i + 1, \dots, m\} = m - i.$$

The proof that $a_i^\perp - a_{2m-i}^\perp = m - i$ is similar. $\square$

As we shall see in a moment, one of the upshots of (7.4.4) is that $\mathcal{F}_i^{\tilde{w}}$ satisfies the spin condition $\iff \mathcal{F}_{2m-i}^{\tilde{w}}$ does. Before explaining this, let us first prove another lemma.

Lemma 7.4.5. *Consider the following conditions.*

- (i) $E_i = E_i^\perp$.
- (ii) $a_i = a_i^\perp$.
- (iii) $\mu_i^{\tilde{w}}(j) \in \{0, 2\}$ for all $j \in B_i$.
- (iv) $\mu_i^{\tilde{w}}$ is self-dual.

Then (i) $\iff$ (ii) $\iff$ (iii) $\implies$ (iv).

Proof. The implication (iii) $\implies$ (iv) is given by (4.3.10).

We now show (ii) $\iff$ (iii). Suppose that $a_i = a_i^\perp$, or in other words, since $\tilde{w}$ is naively permissible, that

$$\#\{j \in \{i+1, \dots, m\} \mid \mu_i^{\tilde{w}}(j) = 0, 1\} = \#\{j \in \{i+1, \dots, m\} \mid \mu_i^{\tilde{w}}(j^*) = 2\}.$$

For j contained in the set on the right-hand side, the basic inequalities (4.3.7) imply that $\mu_i^{\tilde{w}}(j) = 0$. Hence j is contained in the set on the left-hand side. Since the two sets have the same cardinality, they must then be equal. The basic inequalities then imply (iii). Conversely, it is clear that (iii) implies (ii).

We finally show (i) $\iff$ (iii). Suppose that (iii) holds. If $j \in A_i$, then

$$\begin{aligned} j \in E_i &\iff \mu_i^{\tilde{w}}(j) = 0 \\ &\iff \mu_i^{\tilde{w}}(j^*) = 2 \quad (\text{by (iv)}) \\ &\iff 2m + j^* \notin E_i. \end{aligned}$$

If $j \in B_i$, then

$$\begin{aligned} j \in E_i &\iff \mu_i^{\tilde{w}}(j) = 0 \quad (\text{by (iii)}) \\ &\iff \mu_i^{\tilde{w}}(j^*) = 2 \quad (\text{by (iv)}) \\ &\iff 2m + j^* \notin E_i \quad (\text{by (iii)}). \end{aligned}$$

Hence $E_i = E_i^\perp$. Conversely, suppose that there exists $j \in B_i$ such that $\mu_i^{\tilde{w}}(j) = 1$. Then $\mu_i^{\tilde{w}}(j^*) \in \{0, 1\}$ by the basic inequalities. Let $l := \min\{j, j^*\}$. Then $l, 2m + l^* \in E_i$. Hence $E_i \neq E_i^\perp$. $\square$

We are now ready to translate the spin condition for $\mathcal{F}_i^{\tilde{w}}$ and $\mathcal{F}_{2m-i}^{\tilde{w}}$ into combinatorics. Following (7.4.4)(iii), we consider separately the cases $a_i > a_i^\perp$ and $a_i = a_i^\perp$.

First suppose that $a_i > a_i^\perp$, or equivalently, by (7.4.4)(iv), that $a_{2m-i} > a_{2m-i}^\perp$. Consider the elements

$$h_i := \pi_0^{a_i} f_{E_i} + (-1)^s \pi_0^{a_i - a_i^\perp} \pi_0^{a_i^\perp} \operatorname{sgn}(\sigma_{E_i}) f_{E_i^\perp} \in \left(\bigwedge_{\mathcal{O}_F}^{2m} \Lambda_i \right)_{(-1)^s}$$

and

$$\begin{aligned} h_{2m-i} &:= \pi_0^{a_{2m-i}} f_{E_{2m-i}} + (-1)^s \pi_0^{a_{2m-i} - a_{2m-i}^\perp} \pi_0^{a_{2m-i}^\perp} \operatorname{sgn}(\sigma_{E_{2m-i}}) f_{E_{2m-i}^\perp} \\ &\in \left(\bigwedge_{\mathcal{O}_F}^{2m} \Lambda_{2m-i} \right)_{(-1)^s}. \end{aligned}$$

Then the image of h_i in $\bigwedge_k^{2m}(\Lambda_i \otimes_{\mathcal{O}_{F_0}} k)$ spans $\bigwedge_k^{2m} \mathcal{F}_i^{\tilde{w}}$, and analogously with i replaced by $2m-i$. Hence $\mathcal{F}_i^{\tilde{w}}$ and $\mathcal{F}_{2m-i}^{\tilde{w}}$ satisfy the spin condition.

Now suppose that $a_i = a_i^\perp$, or equivalently by (7.4.4)(iv), that $a_{2m-i} = a_{2m-i}^\perp$. Then $E_i = E_i^\perp = E_{2m-i} = E_{2m-i}^\perp$ by (7.4.4) and (7.4.5). Thus we see easily that $\mathcal{F}_i^{\tilde{w}}$ and $\mathcal{F}_{2m-i}^{\tilde{w}}$ simultaneously satisfy or fail the spin condition according as

$$f_{E_i} \in \left(\bigwedge_{F_0}^{2m} V \right)_{(-1)^s} \quad \text{or} \quad f_{E_i} \in \left(\bigwedge_{F_0}^{2m} V \right)_{(-1)^{s+1}},$$

or equivalently, according as

$$\text{sgn}(\sigma_{E_i}) = (-1)^s \quad \text{or} \quad \text{sgn}(\sigma_{E_i}) = (-1)^{s+1}.$$

Lemma 7.4.6. *Let $E \subset \{1, \dots, 4m\}$ be a subset of cardinality $2m$ such that $E = E^\perp$. Then $\sigma_E \in S_{4m}^*$, that is, $\sigma_E(4m+1-j) = 4m+1-\sigma_E(j)$ for all $j \in \{1, \dots, 4m\}$.*

Proof. Obvious from the definition of σ_E . $\square$

It follows from the lemma that σ_{E_i} is even or odd according as the number of elements $j \in \{1, \dots, 2m\}$ such that $\sigma_{E_i}(j) > 2m$ is even or odd. By definition of σ_{E_i} , this is in turn equal to the parity of the cardinality of $E_i \cap \{2m+1, \dots, 4m\}$. By (7.4.5), if $\mu_i^{\tilde{w}}(j) = 1$, then $j \in A_i$. So we see from our explicit description of E_i (7.4.3) that

$$\#(E_i \cap \{2m+1, \dots, 4m\}) = \#\{j \mid \mu_i^{\tilde{w}}(j) = 0\} + \#\{j \mid \mu_i^{\tilde{w}}(j) = 1\}.$$

Since $\mu_i^{\tilde{w}}$ is self-dual, the second term on the right-hand side is even. We conclude that under our assumption $a_i = a_i^\perp$,

$$\mathcal{F}_i^{\tilde{w}} \text{ and } \mathcal{F}_{2m-i}^{\tilde{w}} \text{ satisfy the spin condition} \iff \#\{j \mid \mu_i^{\tilde{w}}(j) = 0\} \equiv s \pmod{2}.$$

Rephrasing slightly, we have now proved the following.

Proposition 7.4.7. $\tilde{w} \in \widetilde{W}_G/W_{I,G}$ is spin-permissible $\iff \tilde{w}$ is wedge-permissible and

(P3) for all $i \in I$, if $\mu_i^{\tilde{w}}$ is self-dual, then $\#\{j \mid \mu_i^{\tilde{w}}(j) = 0\} \equiv s \pmod{2}$ or there exists $j \in B_i$ such that $\mu_i^{\tilde{w}}(j) = 1$. $\square$

Remark 7.4.8. Suppose that $\tilde{w}$ is spin-permissible and $m \in I$. Then $\mu_m^{\tilde{w}}$ is self-dual and $B_m = \emptyset$. Hence (P3) and the condition $\mu_m^{\tilde{w}} + (\mu_m^{\tilde{w}})^* = \mathbf{2}$ in (P1) imply that $\tilde{w} \equiv t_{\mu_r, s} \pmod{W_{\text{aff}, G}}$. On the other hand, it really is necessary to impose the condition $\tilde{w} \equiv t_{\mu_r, s} \pmod{W_{\text{aff}, G}}$ when $m \notin I$. Indeed, for such I and any $\tilde{w} \in \widetilde{W}_G/W_{I,G}$, we have $\mu_i^{\tilde{w}} = \mu_i^{\tilde{w}\tau}$ for all $i \in 2m \pm I$, where τ is the matrix (6.2.2), regarded as an element of the image of S_{2m}^* in $\widetilde{W}_G/W_{I,G}$. Here τ has nontrivial Kottwitz invariant, so that $\tilde{w} \not\equiv \tilde{w}\tau \pmod{W_{\text{aff}, G}}$.

Remark 7.4.9. Suppose that $\tilde{w}$ is wedge-permissible and $0 \in I$. Then we claim that $\mathcal{F}_0^{\tilde{w}}$ automatically satisfies the spin condition. Indeed, $B_0 = \{1, \dots, 2m\}$. So if $\mu_0^{\tilde{w}}(j) \neq 1$ for all $j \in B_0$, then by wedge-permissibility we must have $\mu_0^{\tilde{w}}(j) \in \{0, 2\}$ for all j and $s = m$. Hence $\#\{j \mid \mu_0^{\tilde{w}}(j) = 0\} = s$ on the nose. For $m \geq 2$, it follows that $M_{\{0\}}^\wedge$ and $M_{\{0\}}^{\text{spin}}$ coincide as topological spaces. Once we complete the proof in the next subsection that M_I^{spin} is topologically flat, it will furthermore follow that $M_{\{0\}}^\wedge$ is topologically flat, in support of Pappas's conjecture [P, §4 p. 594] that $M_{\{0\}}^\wedge$ is flat over $\mathcal{O}_E$.

7.5. Topological flatness of M_I^{spin} . We now come to the main algebro-geometric results of the paper. The key combinatorial fact we shall need to prove that M_I^{spin} is topologically flat is the following.

Theorem 7.5.1. *Let $\tilde{w} \in \widetilde{W}_G/W_{I,G}$. Then $\tilde{w}$ is spin-permissible $\iff \tilde{w}$ is $\{\mu_{r,s}\}$ -admissible.*

Proof. We exploit the isomorphism

$$\widetilde{W}_G/W_{I,G} \xrightarrow{\sim} \widetilde{W}_{B_m}/W_{m-I,B_m}$$

induced by the composite isomorphism $\widetilde{W}_G \xrightarrow{\sim} \widetilde{W}_{B_m}$ in (5.7.4). By (5.8.1) this identifies the $\{\mu_{r,s}\}$ -admissible set in $\widetilde{W}_G/W_{I,G}$ with the μ -admissible set in $\widetilde{W}_{B_m}/W_{m-I,B_m}$, where μ denotes the cocharacter $(2^{(s)}, 1^{(2m-2s+1)}, 0^{(s)})$. On the other hand, it follows easily from (5.7.5)(ii) that the displayed isomorphism identifies the set of spin-permissible elements in $\widetilde{W}_G/W_{I,G}$ with the set of μ -spin-permissible elements in $\widetilde{W}_{B_m}/W_{m-I,B_m}$ defined in (9.5.1). So the result follows from the equivalence of μ -admissibility and μ -spin-permissibility in $\widetilde{W}_{B_m}/W_{m-I,B_m}$, which we shall prove later in (9.6.1). $\square$

As corollaries, we deduce Theorems 1.3 and 1.4 from the Introduction.

Proof of (1.4). By definition, the set of spin-permissible elements in $\widetilde{W}_G/W_{I,G}$ surjects onto the set of $w \in W_{I,G} \backslash \widetilde{W}_G/W_{I,G}$ such that the associated Schubert variety S_w is contained in $M_{I,k}^{\text{spin}}$; and the set of $\{\mu_{r,s}\}$ -admissible elements in $\widetilde{W}_G/W_{I,G}$ surjects onto the set of $\{\mu_{r,s}\}$ -admissible elements in $W_{I,G} \backslash \widetilde{W}_G/W_{I,G}$. So the theorem is immediate from (7.5.1). $\square$

Proof of (1.3). Without loss of generality, we may assume that $\mathcal{O}_E$ has algebraically closed residue field. Then, since the set of $\{\mu_{r,s}\}$ -admissible elements in $\widetilde{W}_G$ surjects onto the set of $\{\mu_{r,s}\}$ -admissible elements in $W_{I,G} \backslash \widetilde{W}_G/W_{I,G}$, [PR4, Prop. 3.1] asserts exactly that the Schubert varieties S_w indexed by $\{\mu_{r,s}\}$ -admissible w are contained in $M_{I,k}^{\text{loc}}$. Now use (1.4). $\square$

8. TYPE D COMBINATORICS

In this section we work through the paper's combinatorics in type D . Our main aim is to prove Theorem 1.5 from the Introduction and its parahoric generalization (8.14.5). We shall also prove results, for cocharacters of the form μ below, on permissible sets (see §8.6 and §8.16) and vertexwise admissibility (see §8.15). For most of the section we shall work just in the Iwahori case; beginning in §8.13 we shall turn to the parahoric case. Aside from the preliminaries in §4 (which we shall always apply with $n = 2m$), this section is independent of everything else in the paper.

For convenience, we shall make some notational changes from earlier in the paper. We fix an integer $m \geq 2$. Except where noted to the contrary, throughout this section we denote by μ the cocharacter

$$(8.0.1) \quad \mu := (2^{(q)}, 1^{(2m-2q)}, 0^{(q)}), \quad 0 \leq q \leq m-1.$$

For $i \in \{1, \dots, 2m\}$, we write $i^* := 2m+1-i$. For any nonempty $E \subset \{1, \dots, 2m\}$, we put $E^* := 2m+1-E$. We denote by $e_1, \dots, e_{2m}$ the standard basis in $\mathbb{Z}^{2m}$. We

no longer use the symbol k to denote an algebraic closure of the common residue field of F_0 and F ; instead we shall use k to denote integers. By a *half-integer* we shall mean an element of $\frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$.

8.1. The root datum $\mathcal{R}_{D_m}$. In this subsection we define the root datum $\mathcal{R}_{D_m}$. Let

$$X_{*D_m} := X_{*2m} = \left\{ (x_1, \dots, x_{2m}) \in \mathbb{Z}^{2m} \mid x_1 + x_{2m} = x_2 + x_{2m-1} = \dots = x_m + x_{m+1} \right\},$$

where we recall X_{*2m} from (4.1.1), and

$$X_{D_m}^* := \mathbb{Z}^{2m} / \left\{ (x_1, \dots, x_{m-1}, -\sum_{i=1}^{m-1} x_i, -\sum_{i=1}^{m-1} x_i, x_{m-1}, \dots, x_1) \right\}.$$

Then the standard dot product on $\mathbb{Z}^{2m}$ induces a perfect pairing $X_{D_m}^* \times X_{*D_m} \rightarrow \mathbb{Z}$. For $1 \leq i, j \leq 2m$ with $j \neq i, i^*$, let

$$(8.1.1) \quad \begin{array}{ccc} \alpha_{i,j} : & X_{*D_m} & \longrightarrow \mathbb{Z} \\ & (x_1, \dots, x_{2m}) & \longmapsto x_i - x_j \end{array}$$

and

$$(8.1.2) \quad \alpha_{i,j}^\vee := e_i - e_j + e_{j^*} - e_{i^*} \in X_{*D_m}.$$

Then we may regard $\alpha_{i,j}$ as an element of $X_{D_m}^* \cong \text{Hom}(X_{*D_m}, \mathbb{Z})$, and we let

$$\Phi_{D_m} := \{ \alpha_{i,j} \}_{j \neq i, i^*} \subset X_{D_m}^* \quad \text{and} \quad \Phi_{D_m}^\vee := \{ \alpha_{i,j}^\vee \}_{j \neq i, i^*} \subset X_{*D_m}.$$

The objects so defined form a root datum

$$\mathcal{R}_{D_m} := (X_{D_m}^*, X_{*D_m}, \Phi_{D_m}, \Phi_{D_m}^\vee),$$

which is the root datum for split GO_{2m} . We take as simple roots

$$(8.1.3) \quad \alpha_{1,2}, \alpha_{2,3}, \dots, \alpha_{m-1,m}, \alpha_{m-1,m+1}.$$

The Weyl group of $\mathcal{R}_{D_m}$ identifies canonically with S_{2m}° , as defined at the end of the Introduction. As discussed in §5.1, we then have the extended affine Weyl group

$$\widetilde{W}_{D_m} := X_{*D_m} \rtimes S_{2m}^\circ.$$

The affine Weyl group W_{aff, D_m} is the subgroup $Q_{D_m}^\vee \rtimes S_{2m}^\circ \subset \widetilde{W}_{D_m}$, where $Q_{D_m}^\vee \subset X_{*D_m}$ is the coroot lattice. Explicitly,

$$Q_{D_m}^\vee = \left\{ (x_1, \dots, x_{2m}) \in X_{*D_m} \mid \begin{array}{l} x_1 + x_{2m} = \dots = x_m + x_{m+1} = 0 \\ \text{and } x_1 + \dots + x_m \text{ is even} \end{array} \right\}.$$

Let

$$\mathcal{A}_{D_m} := X_{*D_m} \otimes_{\mathbb{Z}} \mathbb{R} \subset \mathbb{R}^{2m}.$$

We take as positive Weyl chamber the chamber in $\mathcal{A}_{D_m}$ on which the simple roots are all positive, and we say that a cocharacter is dominant if it is contained in the closure of this chamber. We take as our base alcove

$$A_{D_m} := \left\{ (x_1, \dots, x_{2m}) \in \mathbb{R}^{2m} \mid \begin{array}{l} x_1 + x_{2m} = \dots = x_m + x_{m+1} \text{ and} \\ x_1, x_{2m} - 1 < x_2 < x_3 < \dots < x_{m-1} < x_m, x_{m+1} \end{array} \right\};$$

note that this is the unique alcove contained in the Weyl chamber *opposite* the positive chamber and whose closure contains the origin. The minimal facets of A_{D_m} are the lines

$$a + \mathbb{R} \cdot (1, \dots, 1)$$

for a one of the points

$$(8.1.4) \quad \begin{aligned} a_k &:= \left(\left(-\frac{1}{2}\right)^{(k)}, 0^{(2m+1-2k)}, \left(\frac{1}{2}\right)^{(k)} \right) \quad \text{for } k = 0, 2, 3, \dots, m-2, m, \\ a_{0'} &:= (-1, 0^{2m-1}, 1), \\ a_{m'} &:= \left(\left(-\frac{1}{2}\right)^{(m-1)}, \frac{1}{2}, -\frac{1}{2}, \left(\frac{1}{2}\right)^{(m-1)} \right). \end{aligned}$$

As discussed in §5.2, our choice of A_{D_m} endows W_{aff, D_m} and $\widetilde{W}_{D_m}$ with Bruhat orders.

The extended affine Weyl group $\widetilde{W}_{D_m}$ is naturally a subgroup of $\widetilde{W}_{2m}$ (4.1.2). Thus for each $w \in \widetilde{W}_{D_m}$ we get a face $\mathbf{v}$ of type $(2m, \{0, \dots, m\})$ by letting w act on the standard face $\omega_{\{0, \dots, m\}}$, as in §4.2. We write μ_k^w for the vector $\mu_k^{\mathbf{v}}$ attached to this face,

$$(8.1.5) \quad \mu_k^w := w\omega_k - \omega_k \quad \text{for } k \in \mathbb{Z}.$$

The rule $w \mapsto w \cdot \omega_{\{0, \dots, m\}}$ identifies $\widetilde{W}_{D_m}$ with the faces $\mathbf{v}$ of type $(2m, \{0, \dots, m\})$ such that $\mu_0^{\mathbf{v}} \equiv \mu_m^{\mathbf{v}} \pmod{Q_{D_m}^\vee}$.

8.2. μ -spin-permissibility. In this subsection we define μ -spin-permissibility in $\widetilde{W}_{D_m}$, for μ the cocharacter (8.0.1), and we formulate its equivalence with μ -admissibility.

Definition 8.2.1. Let $\mu \in X_{*D_m}$ be the cocharacter (8.0.1). We say that $w \in \widetilde{W}_{D_m}$ is μ -spin-permissible if it satisfies the following conditions for all $0 \leq k \leq m$.

- (SP1) $\mu_k^w + (\mu_{-k}^w)^* = \mathbf{2}$ and $\mathbf{0} \leq \mu_k^w \leq \mathbf{2}$.
- (SP2) $\#\{j \mid \mu_k^w(j) = 2\} \leq q$.
- (SP3) (spin condition) If μ_k^w is self-dual and $\mu_k^w \not\equiv \mu \pmod{Q_{D_m}^\vee}$, then there exist elements $j_1 \in A_k$ and $j_2 \in B_k$ such that $\mu_k^w(j_1) = \mu_k^w(j_2) = 1$, with the sets A_k and B_k as in (4.3.6).

For fixed k with $0 \leq k \leq m$, we say that μ_k^w is μ -spin-permissible if it satisfies (SP1), (SP2), and (SP3). We say that w is *naively μ -permissible* if it satisfies (SP1) for all $0 \leq k \leq m$.

Note that the condition $\mu_k^w + (\mu_{-k}^w)^* = \mathbf{2}$ in (SP1) holds for all k as soon as it holds for a single k , and says just that w defines a 2-face. By (4.4.3), when (SP1) is satisfied, the quantity $\#\{j \mid \mu_k^w(j) = 2\}$ appearing in (SP2) is also equal to $\#\{j \mid \mu_k^w(j) = 0\}$; we shall study it in more detail in §8.4. Also note that the $k = 0$ and $k = m$ cases of (SP3) require respectively that $\mu_0^w \equiv \mu$ and $\mu_m^w \equiv \mu \pmod{Q_{D_m}^\vee}$, since μ_0^w and μ_m^w are always self-dual and $A_0 = B_m = \emptyset$. Of course, the condition that μ_k^w be self-dual is exactly the condition that it be contained in X_{*D_m} .

The key result we shall prove in §8 is Theorem 1.5 from the Introduction, which we formulate in our present notation as follows. Recall from §5.3 that an element $w \in \widetilde{W}_{D_m}$ is μ -admissible if $w \leq t_\lambda$ for some $\lambda \in S_{2m}^\circ \cdot \mu$.

Theorem 8.2.2. *Let μ be the cocharacter (8.0.1). Then $w \in \widetilde{W}_{D_m}$ is μ -admissible $\iff w$ is μ -spin-permissible.*

We shall generalize the theorem to the general parahoric case in (8.14.5). We emphasize that the theorem does not include the case $\mu = (2^{(m)}, 0^{(m)})$; we shall not consider in this paper how to characterize the admissible set for this cocharacter. We shall prove the implication $\implies$ in §8.10 and the implication $\impliedby$ in §8.12; the

intervening sections shall large serve to lay the groundwork. The implication $\Leftarrow$ is by far the harder of the two. The strategy we shall use to prove it is, in essence, that of Kottwitz–Rapoport [KR], which is also the strategy used in [Sm1, Sm2]. To explain it in the present situation, let us first prove a baby case of the theorem.

Lemma 8.2.3. *Let $w = t_\nu$ be a translation element in $\widetilde{W}_{D_m}$, and suppose that w is μ -spin-permissible. Then w is μ -admissible.*

Proof. Let ν^+ denote the dominant Weyl conjugate of ν . Then it is immediate from the definition of μ -spin-permissibility that $\nu^+ \preceq \mu$ in the dominance order. Hence w is μ -admissible. $\square$

To now explain our strategy to prove the implication $\Leftarrow$ in (8.2.2), suppose that $w \in \widetilde{W}_{D_m}$ is μ -spin-permissible. If w is a translation element, then w is μ -admissible by the lemma. If w is not a translation element, then our task will be to find an affine root $\tilde{\alpha}$ for $\mathcal{R}_{D_m}$ such that, for the associated reflection $s_{\tilde{\alpha}} \in W_{\text{aff}, D_m}$, $s_{\tilde{\alpha}}w$ is again μ -spin-permissible and $w < s_{\tilde{\alpha}}w$ in the Bruhat order. For then, repeating the argument as needed, we obtain a chain $w < s_{\tilde{\alpha}}w < \cdots$ of μ -spin-permissible elements which must terminate in a translation element, since the set of μ -spin-permissible elements is manifestly finite.

8.3. Proper elements. Let $w = t_{\mu_0^w} \sigma \in \widetilde{W}_{D_m}$, with $\mu_0^w \in X_{*D_m}$ and $\sigma \in S_{2m}^\circ$.

Definition 8.3.1. We say the element $j \in \{1, \dots, 2m\}$ is *proper* if $\sigma(j) \neq j$.

Of course j is proper $\iff j^*$ is proper. The formula (8.1.5) for μ_k^w gives the obvious recursion relation

$$(8.3.2) \quad \mu_k^w = \mu_{k-1}^w + e_k - e_{\sigma(k)}, \quad 1 \leq k \leq 2m.$$

When j is proper, it follows from this that $\mu_k^w(j)$ takes exactly two values as k varies between 0 and $2m$ and j remains fixed, and that these two values differ by 1.

Definition 8.3.3. For $j \in \{1, \dots, 2m\}$ proper, we define the *upper value*

$$u(j) := \max\{\mu_k^w(j)\}_{0 \leq k \leq 2m}.$$

Of the course the upper value depends on w as well as j , but to avoid clutter, we do not embed w in the notation. If j is proper and w defines a d -face, then it is clear from the formula (4.3.4) that $u(j) + u(j^*) = d + 1$. In particular, when w satisfies (SP1), one of every pair j, j^* of proper elements has upper value 2, and the other has upper value 1.

8.4. The integer c_k^w . We continue with $w = t_{\mu_0^w} \sigma \in \widetilde{W}_{D_m}$. In this subsection we study the quantity $\#\{j \mid \mu_k^w(j) = 2\}$ appearing in condition (SP2). In light of (4.4.3), we make the following definition.

Definition 8.4.1. Let $0 \leq k \leq m$, and suppose that μ_k^w satisfies (SP1). We define c_k^w to be the common integer

$$c_k^w := \#\{j \mid \mu_k^w(j) = 2\} = \#\{j \mid \mu_k^w(j) = 0\}.$$

Thus we write $c_k^w \leq q$ for condition (SP2) in the definition of μ -spin-permissible; it will be useful to have both interpretations of c_k^w in the display. Another use for c_k^w is the following, which is relevant for the spin condition.

Lemma 8.4.2. *Suppose that μ_k^w satisfies (SP1) and is self-dual. Then $\mu_k^w \equiv \mu \pmod{Q_{D_m}^\vee} \iff c_k^w \equiv q \pmod{2}$.*

Proof. The map $X_{*D_m} \rightarrow \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ sending

$$(x_1, \dots, x_{2m}) \mapsto (d, x_1 + x_2 + \dots + x_m \pmod{2}),$$

where d denotes the common integer $x_1 + x_{2m} = \dots = x_m + x_{m+1}$, induces an isomorphism $X_{*D_m}/Q_{D_m}^\vee \xrightarrow{\sim} \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$. This renders the lemma transparent. $\square$

We conclude the subsection with a couple of simple lemmas.

Lemma 8.4.3. *Let $1 \leq k \leq m$, and suppose that μ_k^w and μ_{k-1}^w satisfy (SP1). Then $c_k^w = c_{k-1}^w + u(k) - u(\sigma(k))$.*

Proof. This is easily checked using the recursion relation (8.3.2). $\square$

Our second lemma will prove to be a very useful tool later on in the proof of the key proposition (8.12.1).

Lemma 8.4.4. *Let $0 \leq i < p \leq m$, and suppose that μ_i^w and μ_p^w satisfy (SP1).*

- (i) *If $c_i^w < c_p^w$, then there exists $j \in \{i+1, \dots, p\}$ such that $\mu_i^w(j) = 1$ and $\mu_p^w(j) = 2$.*
- (ii) *If $c_i^w > c_p^w$, then there exists $j \in \{i+1, \dots, p\}$ such that $\mu_i^w(j) = 0$ and $\mu_p^w(j) = 1$.*

Proof. The hypothesis in (i) implies that there exists a j such that $\mu_p^w(j) = 2$ and $\mu_i^w(j) = 1$. The recursion relation (8.3.2) then forces $j \in \{i+1, \dots, p\}$. Assertion (ii) is proved in a similar way, using that c_i^w also equals the number of entries of μ_i^w equal to 0. $\square$

8.5. The vector ν_k^w . We continue with our element $w = t_{\mu_0^w} \sigma \in \widetilde{W}_{D_m}$. In this subsection we introduce the vector ν_k^w , which for many purposes is better to work with than μ_k^w .

In addition to the vertices (8.1.4), let us define the points in $\mathcal{A}_{D_m}$

$$(8.5.1) \quad a_1 := \left(-\frac{1}{2}, 0^{(2m-2)}, \frac{1}{2}\right) \quad \text{and} \quad a_{m-1} := \left(\left(-\frac{1}{2}\right)^{(m-1)}, 0, 0, \left(\frac{1}{2}\right)^{(m-1)}\right).$$

Then $a_1, a_{m-1} \in \overline{A_{D_m}}$, but neither is a vertex. Instead the points $a_0, a_1, \dots, a_m$ are vertices for an alcove for the symplectic group. Since this symplectic alcove is contained in A_{D_m} , we shall find the a_k 's quite suitable for our purposes.

Definition 8.5.2. For $0 \leq k \leq m$, we define

$$\nu_k^w := wa_k - a_k = \frac{\mu_k^w + \mu_{-k}^w}{2} = \frac{\mu_k^w + \mu_{2m-k}^w}{2}.$$

Note that $\nu_k^w = \mu_k^w$ whenever μ_k^w is self-dual; in particular $\nu_0^w = \mu_0^w$ and $\nu_m^w = \mu_m^w$. If w defines a d -face, then (4.3.4) gives

$$(8.5.3) \quad \nu_k^w = \frac{\mu_k^w - (\mu_k^w)^* + \mathbf{d}}{2}.$$

Since μ_k^w has only integer entries, the basic inequalities (4.3.7) then allow us to recover μ_k^w uniquely from ν_k^w . Using the recursion relation (8.3.2) for μ_k^w , we get

the recursion relation for ν_k^w , valid for $1 \leq k \leq m$,

$$(8.5.4) \quad \begin{aligned} \nu_k^w &= \frac{\mu_{k-1}^w + e_k - e_{\sigma(k)} + e_{\sigma(k^*)} - e_{k^*} + \mu_{k^*}^w}{2} \\ &= \nu_{k-1}^w + \begin{cases} 0, & \sigma(k) = k \\ e_k - e_{k^*}, & \sigma(k) = k^* \\ \frac{1}{2}\alpha_{k, \sigma(k)}^\vee, & \sigma(k) \neq k, k^*. \end{cases} \end{aligned}$$

It is clear from the definition of ν_k^w that

$$\nu_k^w(j) \in \{u(j) - 1, u(j) - \frac{1}{2}, u(j)\} \quad \text{for all } 0 \leq k \leq m \text{ and } 1 \leq j \leq 2m.$$

We shall devote the rest of the subsection to expressing conditions (SP1)–(SP3) in terms of the vector ν_k^w .

Lemma 8.5.5. *Let $0 \leq k \leq m$. The following are equivalent.*

- (i) μ_k^w is self-dual.
- (ii) $\nu_k^w = \mu_k^w$.
- (iii) $\nu_k^w \in \mathbb{Z}^{2m}$ (as opposed to $\frac{1}{2}\mathbb{Z}^{2m}$).

Proof. The implications (i) $\implies$ (ii) and (ii) $\implies$ (iii) are trivial. Now assume (iii). To deduce (i), we must show that $\mu_k^w = \mu_{-k}^w$, or equivalently, if w defines a d -face, that $\mu_k^w = \mathbf{d} - (\mu_k^w)^*$. This is a straightforward consequence of the formula (8.5.3) and the basic inequalities (4.3.7). $\square$

Note that if $\nu_k^w \in \mathbb{Z}^{2m}$, then $\nu_k^w = \mu_k^w \in X_{*D_m}$.

Lemma 8.5.6. *Let $0 \leq k \leq m$, and suppose that μ_k^w satisfies (SP1). Then*

$$\begin{aligned} c_k^w &= \#\{j \mid \nu_k^w(j) = 2\} + \frac{\#\{j \mid \nu_k^w(j) \notin \mathbb{Z}\}}{4} \\ &= \#\{j \mid \nu_k^w(j) = 0\} + \frac{\#\{j \mid \nu_k^w(j) \notin \mathbb{Z}\}}{4}. \end{aligned}$$

Proof. Since μ_k^w satisfies (SP1), we have $\nu_k^w + (\nu_k^w)^* = \mathbf{2}$. This implies the second equality. To prove the first, consider the sets defined in (4.4.2), where we replace μ_i^y with μ_k^w . We have

$$\{j \mid \nu_k^w(j) = 2\} = G \quad \text{and} \quad \{j \mid \nu_k^w(j) \notin \mathbb{Z}\} = E \amalg E^* \amalg F \amalg F^*.$$

By (4.4.1) $\#E = \#F$, and the lemma follows. $\square$

We now introduce the following conditions on ν_k^w for $0 \leq k \leq m$.

- (SP1') $\nu_k^w + (\nu_k^w)^* = \mathbf{2}$ and $\mathbf{0} \leq \nu_k^w \leq \mathbf{2}$.
- (SP2') $\#\{j \mid \nu_k^w(j) = 2\} + \#\{j \mid \nu_k^w(j) \notin \mathbb{Z}\}/4 \leq q$.
- (SP3') (spin condition) If $\nu_k^w \in \mathbb{Z}^{2m}$ and $\nu_k^w \not\equiv \mu \pmod{Q_{D_m}^\vee}$, then there exist elements $j_1 \in A_k$ and $j_2 \in B_k$ such that $\nu_k^w(j_1) = \nu_k^w(j_2) = 1$.

Lemma 8.5.7. *Let $0 \leq k \leq m$.*

- (i) μ_k^w satisfies (SP1) $\iff \nu_k^w$ satisfies (SP1').
- (ii) Suppose that the equivalent conditions in (i) hold. Then μ_k^w satisfies (SP2) $\iff \nu_k^w$ satisfies (SP2').
- (iii) μ_k^w satisfies (SP3) $\iff \nu_k^w$ satisfies (SP3').

Thus w is μ -spin-permissible $\iff$ (SP1')–(SP3') hold for all $0 \leq k \leq m$. We shall find this formulation of μ -spin-permissibility more convenient to work with than the original one (8.2.1). We remind the reader that when the equivalent conditions (SP1) and (SP1') hold and $\nu_k^w \in \mathbb{Z}^{2m}$, the condition $\nu_k^w \not\equiv \mu \pmod{Q_{D_m}^\vee}$ is equivalent to $c_k^w \not\equiv q \pmod{2}$ by (8.4.2). We shall usually work with this latter formulation of the spin condition in practice.

Proof of (8.5.7). Assertion (ii) is proved by (8.5.6), and assertion (iii) is proved by (8.5.5). For (i), suppose that w defines a d -face. Then $\mu_k^w + (\mu_{-k}^w)^* = \mathbf{d}$, and

$$\nu_k^w + (\nu_k^w)^* = \frac{\mu_k^w + \mu_{-k}^w + (\mu_k^w)^* + (\mu_{-k}^w)^*}{2} = \mathbf{d}.$$

So the conditions $\mu_k^w + (\mu_{-k}^w)^* = \mathbf{2}$ and $\nu_k^w + (\nu_k^w)^* = \mathbf{2}$ are equivalent. Assume that they hold. If $\mathbf{0} \leq \mu_k^w \leq \mathbf{2}$, then this assumption implies that $\mathbf{0} \leq \mu_{-k}^w \leq \mathbf{2}$. Hence $\mathbf{0} \leq \nu_k^w \leq \mathbf{2}$. Conversely, if $\mathbf{0} \leq \nu_k^w \leq \mathbf{2}$, then writing ν_k^w as in (8.5.3) and combining with the basic inequalities (4.3.7) (applied with $d = 2$), we easily get $\mathbf{0} \leq \mu_k^w \leq \mathbf{2}$. $\square$

8.6. μ -permissibility. We continue with our element $w = t_{\nu_0^w} \sigma \in \widetilde{W}_{D_m}$. In this subsection we shall characterize the μ -permissible set of Kottwitz–Rapoport (see §5.3) inside $\widetilde{W}_{D_m}$ in terms of a subset of the conditions that define the μ -spin-permissible set. We begin with a lemma which gives a fairly direct geometric interpretation of conditions (SP1') and (SP2'). Let $\text{Conv}(S_{2m}^\circ \mu)$ denote the convex hull in $\mathcal{A}_{D_m}$ of the Weyl group orbit $S_{2m}^\circ \mu$.

Lemma 8.6.1. *Let $0 \leq k \leq m$. Then ν_k^w satisfies (SP1') and (SP2') $\iff \nu_k^w \in \text{Conv}(S_{2m}^\circ \mu)$.*

Proof. This is proved in the proof of [Sm3, Prop. 6.6.2], but for convenience we shall make the argument explicit here. Recall the group S_{2m}^* from the Introduction. Since $q < m$, we have $S_{2m}^\circ \mu = S_{2m}^* \mu$. For $1 \leq i \leq m$, let $\lambda_i = (1^{(i)}, 0^{(2m-i)})$. Then the convex hull $\text{Conv}(S_{2m}^\circ \mu) = \text{Conv}(S_{2m}^* \mu)$ admits the description

$$(8.6.2) \quad \text{Conv}(S_{2m}^\circ \mu) = \left\{ v \in \mathcal{A}_{D_m} \mid \begin{array}{l} v + v^* = \mathbf{2} \text{ and } \lambda \cdot v \leq \lambda_i \cdot \mu \\ \text{for all } 1 \leq i \leq m \text{ and all } \lambda \in S_{2m}^* \lambda_i \end{array} \right\},$$

where we use the standard dot product on $\mathbb{R}^{2m}$; see for example [Sm3, Lem. 6.6.1].

Now suppose that ν_k^w satisfies (SP1') and (SP2'). Then, using (8.5.7), μ_k^w and μ_{-k}^w each have all entries contained in $\{0, 1, 2\}$ and at most q entries equal to 2. Hence

$$\lambda \cdot \mu_k^w, \lambda \cdot \mu_{-k}^w \leq \lambda_i \cdot \mu \quad \text{for all } 1 \leq i \leq m \text{ and } \lambda \in S_{2m}^* \lambda_i.$$

Hence

$$\lambda \cdot \nu_k^w = \lambda \cdot \left(\frac{\mu_k^w + \mu_{-k}^w}{2} \right) \leq \lambda_i \cdot \mu \quad \text{for all } 1 \leq i \leq m \text{ and } \lambda \in S_{2m}^* \lambda_i.$$

Hence $\nu_k^w \in \text{Conv}(S_{2m}^\circ \mu)$.

Conversely, suppose that $\nu_k^w \in \text{Conv}(S_{2m}^\circ \mu)$. Then ν_k^w clearly satisfies (SP1'). Next recall the sets E , F , G , and H from (4.4.2), where μ_k^w replaces μ_i^v , and again denote their respective cardinalities by e , f , g , and h . By (4.4.1) $e = f$, and by

definition $c_k^w = e + g = f + g$. Let $l := e + f + g$. If $l < q$, then trivially $c_k^w < q$. If $l \geq q$, then

$$\begin{aligned} \sum_{j \in E \amalg F \amalg G} \nu_k^w(j) &= \frac{3}{2}e + \frac{3}{2}f + 2g = 3e + 2g \\ &\leq \lambda_l \cdot \mu = 2q + (l - q) = q + 2e + g, \end{aligned}$$

where the inequality in the middle holds because $\nu_k^w \in \text{Conv}(S_{2m}^\circ \mu)$. Hence

$$c_k^w = e + g \leq q,$$

as desired. $\square$

Now recall the vertices $a_{0'}$ and $a_{m'}$ from (8.1.4), and in analogy with (8.5.2), define

$$\nu_{0'}^w := wa_{0'} - a_{0'} \quad \text{and} \quad \nu_{m'}^w := wa_{m'} - a_{m'}.$$

Lemma 8.6.3.

- (i) ν_0^w and ν_1^w are μ -spin-permissible $\iff w \equiv t_\mu \pmod{W_{\text{aff}, D_m}}$ and $\nu_0^w, \nu_{0'}^w \in \text{Conv}(S_{2m}^\circ \mu)$.
- (ii) ν_{m-1}^w and ν_m^w are μ -spin-permissible $\iff w \equiv t_\mu \pmod{W_{\text{aff}, D_m}}$ and $\nu_m^w, \nu_{m'}^w \in \text{Conv}(S_{2m}^\circ \mu)$.

Proof. We shall just prove (i); (ii) is proved in a completely analogous way, with ν_{m-1}^w in the role of ν_1^w , ν_m^w in the role of ν_0^w , and $\nu_{m'}^w$ in the role of $\nu_{0'}^w$. Let

$$\varepsilon := e_1 - e_{\sigma(1)} + e_{\sigma(2m)} - e_{2m}.$$

Then

$$\nu_{0'}^w = \nu_0^w + \varepsilon \quad \text{and} \quad \nu_1^w = \nu_0^w + \frac{1}{2}\varepsilon.$$

The congruence $w \equiv t_\mu \pmod{W_{\text{aff}, D_m}}$ is equivalent to $\nu_0^w \equiv \mu \pmod{Q_{D_m}^\vee}$, and we shall assume that both hold throughout the proof.

Let us first prove the implication $\Leftarrow$. Lemma 8.6.1 gives us what we need to conclude that ν_0^w is μ -spin-permissible. Since ν_1^w is the midpoint of ν_0^w and $\nu_{0'}^w$, we have $\nu_1^w \in \text{Conv}(S_{2m}^\circ \mu)$, and we conclude from (8.6.1) that ν_1^w satisfies (SP1') and (SP2').

To see that ν_1^w satisfies the spin condition, suppose that $\nu_1^w \in \mathbb{Z}^{2m}$. Then, since $\nu_0^w \in \mathbb{Z}^{2m}$, we must have $\varepsilon = 0$ or $\varepsilon = 2e_1 - 2e_{2m}$. In the former case $\nu_1^w = \nu_0^w$ certainly satisfies (SP3'). In the latter case, since $\nu_0^w, \nu_{0'}^w \in \text{Conv}(S_{2m}^\circ \mu)$, we must have $\nu_0^w(1) = 0$ and $\nu_{0'}^w(1) = 2$. Hence $\nu_1^w(1) = 1$. Since $q < m$, there also exists a j such that $\nu_0^w(j) = 1$. Evidently $j \neq 1, 2m$, whence $\nu_1^w(j) = 1$. Since $1 \in A_1$ and $j \in B_1$, we conclude that ν_1^w satisfies (SP3').

We now prove the implications $\Rightarrow$. Lemma 8.6.1 immediately gives $\nu_0^w \in \text{Conv}(S_{2m}^\circ \mu)$. It remains to show that $\nu_{0'}^w \in \text{Conv}(S_{2m}^\circ \mu)$. For this, since plainly $\nu_{0'}^w + (\nu_{0'}^w)^* = \mathbf{2}$ and $\nu_{0'}^w \in \mathbb{Z}^{2m}$, it is immediate from the description of the convex hull (8.6.2) that it suffices to show that $\mathbf{0} \leq \nu_{0'}^w \leq \mathbf{2}$ and $c_{0'}^w \leq q$, where

$$c_{0'}^w := \{j \mid \nu_{0'}^w(j) = 2\}.$$

To do so, we shall enter into a case analysis based on the possibilities for $\sigma(1)$.

If $\sigma(1) = 1$, then $\varepsilon = 0$ and $\nu_{0'}^w = \nu_0^w \in \text{Conv}(S_{2m}^\circ \mu)$. If $\sigma(1) = 2m$, then $\varepsilon = 2e_1 - 2e_{2m}$. Since $\nu_1^w = \nu_0^w + e_1 - e_{2m}$ satisfies (SP1'), we must have $\nu_0^w(1) \neq 2$. We claim that furthermore $\nu_0^w(1) \neq 1$. For evidently $\nu_1^w \in \mathbb{Z}^{2m}$, and $c_1^w = c_0^w \pm 1 \neq q \pmod{2}$ by (8.4.2). If $\nu_0^w(1) = 1$, then $\nu_1^w(1) = 2$ and $\nu_1^w(2m) = 0$, so that $\nu_1^w(j) \neq 1$

for all $j \in A_1$, in violation of the spin condition. Thus the only possibility is that $\nu_0^w(1) = 0$, and we see by inspection that $\mathbf{0} \leq \nu_{0'}^w \leq \mathbf{2}$ and $c_{0'}^w = c_0^w \leq q$.

Finally suppose that $\sigma(1) \neq 1, 2m$. Since ν_1^w satisfies (SP1'), we have $\nu_0^w(1) \in \{0, 1\}$ and $\nu_0^w(\sigma(1)) \in \{1, 2\}$. Hence $\mathbf{0} \leq \nu_{0'}^w \leq \mathbf{2}$. Moreover, by (8.4.3) $c_1^w \in \{c_0^w - 1, c_0^w, c_0^w + 1\}$, and one sees easily that $c_{0'}^w = c_0^w + 2(c_1^w - c_0^w)$. So if $c_1^w \leq c_0^w$, then certainly $c_{0'}^w \leq c_0^w \leq q$. If $c_1^w = c_0^w + 1$, then the conditions $c_1^w \leq q$ and $c_0^w \equiv q \pmod{2}$ give us $c_0^w + 1 = c_1^w < q$. Hence $c_{0'}^w = c_0^w + 2 \leq q$, as desired. $\square$

Our characterization of the μ -permissible set in $\widetilde{W}_{D_m}$ is as follows.

Proposition 8.6.4. *Let μ be the cocharacter (8.0.1). Then w is μ -permissible $\iff$ conditions (SP1') and (SP2') hold for all $0 \leq k \leq m$, and condition (SP3') holds for $k = 0, 1, m-1, m$.*

Proof. Since each minimal facet of A_{D_m} contains exactly one of the vertices $a_0, a_{0'}, a_2, a_3, \dots, a_{m-2}, a_m, a_{m'}$, this is obvious from (8.6.1) and (8.6.3). $\square$

Example 8.6.5. The proposition affords us a ready supply of elements in $\widetilde{W}_{D_m}$ that are μ -permissible but not μ -spin-permissible. Once we have shown later in (8.10.4) that μ -admissibility implies μ -spin-permissibility, this will give us examples of elements that are μ -permissible but not μ -admissible. For example, take $m = 4$ and $\mu = (2^{(3)}, 1, 1, 0^{(3)})$, and let $w = t_{(2,1^{(6)},0)}(27)(36)$, where we use cycle notation to denote elements in the Weyl group. Then one readily verifies, using (8.6.4), that w is μ -permissible. However

$$\nu_2^w = (2, 2, 1, 1, 1, 1, 0, 0) \not\equiv \mu \pmod{Q_{D_m}^\vee},$$

in violation of the spin condition. This example generalizes in an obvious way to show that the μ -admissible and μ -permissible sets in $\widetilde{W}_{D_m}$ differ whenever $m \geq 4$ and $q \geq 3$ (still with $q < m$, of course).

Note that for $m \geq 4$,

$$(2^{(m-1)}, 1, 1, 0^{(m-1)}) = (1^{(m)}, 0^{(m)}) + (1^{(m-1)}, 0, 1, 0^{(m-1)})$$

is a sum of two dominant minuscule cocharacters. Thus we get a counterexample to Rapoport's conjecture [R, §3, p. 283] that μ -admissibility and μ -permissibility are equivalent whenever μ is a sum of dominant minuscule cocharacters.

Two further remarks on this point are in order. First, our counterexample aside, Rapoport's conjecture is known to hold true when μ *itself* is minuscule and the root datum involves only types A, B, C and D ; see [PRS, Prop. 4.4.5] or [Sm2, Th. 2.2.1] for an overview. Moreover, no counterexamples to the conjecture in other types are known when μ is minuscule. Second, something of a replacement for Rapoport's conjecture in general is proposed in [PRS, Conj. 4.5.3] in terms of the notion of *vertexwise admissibility*. See the discussion in [PRS, §4.5]. We shall show in (8.15.3) that μ -admissibility is equivalent to μ -vertexwise admissibility for the cocharacter μ (8.0.1).

As a counterpoint to the example just given, we have the following result.

Proposition 8.6.6. *Let μ be the cocharacter (8.0.1), and suppose that $q \leq 2$. Then w is μ -admissible $\iff w$ is μ -permissible.*

Proof. By Kottwitz–Rapoport [KR, 11.2], we need only prove the implication $\Leftarrow$. We shall do so by assuming the equivalence of μ -admissibility and μ -spin-permissibility asserted in (8.2.2); the proof of this equivalence will be completed in §8.12

and will make no use of the result we are proving now. So suppose that w is μ -permissible. If $q = 0$, then the μ -admissible and μ -permissible sets each consist of a single element, namely $t_{(1^{(2m)})}$, and the result is trivial. If q equals 1 or 2, then we must show that ν_k^w satisfies the spin condition for all $2 \leq k \leq m-2$. Fix such a k . Since ν_k^w satisfies (SP1') and (SP2'), we immediately reduce to considering the case that $\nu_k^w \in \mathbb{Z}^{2m}$ and $c_k^w < q$, i.e. c_k^w equals 0 or 1. But our restriction on k means that A_k and B_k each have cardinality at least 4. The constraint on c_k^w then forces each to contain an element j such that $\nu_k^w(j) = 1$. $\square$

Remark 8.6.7. For $m = 2, 3$, we conclude that μ -permissibility and μ -admissibility in $\widetilde{W}_{D_m}$ are equivalent. Here the root system Φ_{D_m} is of type $A_1 \times A_1$ and A_3 , respectively. Thus this equivalence is a special case of the general result of Haines–Ngô [HN, Th. 1] that μ -admissibility and μ -permissibility are equivalent for all cocharacters μ in root data of type A .

8.7. Intervals. We continue with our element $w = t_{\nu_0^w} \sigma \in \widetilde{W}_{D_m}$. By the periodicity relation (4.3.2) (applied with $n = 2m$), we may regard the vector μ_k^w as indexed by $k \in \mathbb{Z}/2m\mathbb{Z}$.

Definition 8.7.1. For $j \in \{1, \dots, 2m\}$ proper, we define K_j to be the set of all $k \in \mathbb{Z}/2m\mathbb{Z}$ for which $\mu_k^w(j)$ takes its lower value,² i.e. the value $u(j) - 1$.

For j proper, the set K_j is an example of an *interval* in $\mathbb{Z}/2m\mathbb{Z}$, in the terminology of Kottwitz–Rapoport [KR, 5.4]. Recall from their paper that if a and b are distinct in $\mathbb{Z}/2m\mathbb{Z}$, then the interval $[a, b) \subset \mathbb{Z}/2m\mathbb{Z}$ is defined in the following way. Let $\tilde{a}$ denote any representative of a in $\mathbb{Z}$, and let $\tilde{b}$ denote the unique representative of b in $\mathbb{Z}$ satisfying $\tilde{a} < \tilde{b} < \tilde{a} + 2m$. Then $[a, b)$ is the image of $\{k \in \mathbb{Z} \mid \tilde{a} \leq k < \tilde{b}\}$ in $\mathbb{Z}/2m\mathbb{Z}$. For j proper, it is immediate from the recursion relation (8.3.2) that K_j equals the interval $[\sigma^{-1}(j), j)$ in $\mathbb{Z}/2m\mathbb{Z}$; here and elsewhere we use the same symbols to denote integers and their residues mod $2m\mathbb{Z}$. Of course $K_{j^*} = [\sigma^{-1}(j^*), j^*) = [\sigma^{-1}(j)^*, j^*)$.

Note that if $i, i' \in \mathbb{Z}/2m\mathbb{Z}$ are distinct from j , then one of the intervals $[i, j)$ and $[i', j)$ is always contained in the other.

Let us extend the definition of ν_k^w to all $k \in \mathbb{Z}/2m\mathbb{Z}$ in the obvious way, by setting $\nu_k^w = \frac{1}{2}(\mu_k^w + \mu_{-k}^w)$. Then for proper fixed j , the sets of k for which $\nu_k^w(j)$ takes its various values are easily described in terms of intervals. Indeed, the formula (8.5.3) makes clear that

$$\begin{aligned} \{k \in \mathbb{Z}/2m\mathbb{Z} \mid \nu_k^w(j) = u(j) - 1\} &= K_j \cap K_{j^*}^c = [\sigma^{-1}(j), j) \cap [j^*, \sigma^{-1}(j)^*), \\ \{k \in \mathbb{Z}/2m\mathbb{Z} \mid \nu_k^w(j) = u(j)\} &= K_j^c \cap K_{j^*} = [j, \sigma^{-1}(j)) \cap [\sigma^{-1}(j)^*, j^*), \end{aligned}$$

²This definition is the analog of the set denoted K_j in [KR, 5.2], but, in terms of the terminology we've introduced, Kottwitz and Rapoport define K_j to be the set of all k where $\mu_k^w(j)$ takes its *upper* value. This discrepancy is due to differing choices of base alcove: our a_k 's are located in the closure of the Weyl chamber for GL_{2m} *opposite* the one containing Kottwitz and Rapoport's base alcove. On the other hand, our definition of K_j is in obvious analogy with the definition of K_j in [Sm1, §8.3].

and

$$\begin{aligned} \left\{ k \in \mathbb{Z}/2m\mathbb{Z} \mid \nu_k^w(j) = u(j) - \frac{1}{2} \right\} &= (K_j \cup K_{j^*}^c) \setminus (K_j \cap K_{j^*}^c) \\ &= (K_j^c \cup K_{j^*}) \setminus (K_j^c \cap K_{j^*}), \end{aligned}$$

where we use a superscript c to denote set complements in $\mathbb{Z}/2m\mathbb{Z}$. Of course, some of these sets may be empty.

In practice, rather than having to consider $\nu_k^w(j)$ for all $k \in \mathbb{Z}/2m\mathbb{Z}$, we shall wish to restrict just to k in the range $0 \leq k \leq m$. The following is an immediate consequence of our above discussion; it also follow easily from the recursion relation (8.5.4).

Lemma 8.7.2. *Let $j \in \{1, \dots, 2m\}$. Then exactly one of the following three possibilities holds.*

- (1) j is not proper, and $\nu_k^w(j)$ is constant as k varies in the range $0 \leq k \leq m$.
- (2) $\sigma(j) = j^*$, and $\nu_k^w(j)$ takes exactly the two values $u(j)$ and $u(j) - 1$ as k varies in the range $0 \leq k \leq m$. There is exactly one k in the range $1 \leq k \leq m$ with the property that $\nu_k^w(j) \neq \nu_{k-1}^w(j)$, namely $k = \min\{j, j^*\}$.
- (3) $\sigma(j) \neq j, j^*$, and $\nu_k^w(j)$ takes the value $u(j) - \frac{1}{2}$ for some k in the range $0 \leq k \leq m$. There are exactly two k in the range $1 \leq k \leq m$ with the property that $\nu_k^w(j) \neq \nu_{k-1}^w(j)$, namely $k = \min\{j, j^*\}$ and $k = \min\{\sigma^{-1}(j), \sigma^{-1}(j)^*\}$. For each of these two k , the values $\nu_{k-1}^w(j)$ and $\nu_k^w(j)$ differ by $\pm \frac{1}{2}$. $\square$

8.8. Reflections. The affine roots in $\mathcal{R}_{D_m}$ are the functions

$$(8.8.1) \quad \begin{aligned} \tilde{\alpha}_{i,j;d}: \quad X_{*D_m} &\longrightarrow \mathbb{Z} \\ (x_1, \dots, x_{2m}) &\longmapsto x_i - x_j - d \end{aligned}$$

for $1 \leq i, j \leq 2m$ with $j \neq i, i^*$, and $d \in \mathbb{Z}$. Plainly $\tilde{\alpha}_{i,j;d} = \tilde{\alpha}_{j^*,i^*;d}$. Attached to the affine root $\tilde{\alpha} = \tilde{\alpha}_{i,j;d}$ is its linear part $\alpha := \alpha_{i,j}$ (8.1.1); the coroot $\alpha^\vee := \alpha_{i,j}^\vee$ (8.1.2); and the reflection $s_{\tilde{\alpha}} \in W_{\text{aff}, D_m}$ sending

$$v \longmapsto v - \langle \tilde{\alpha}, v \rangle \alpha^\vee, \quad v \in \mathcal{A}_{D_m};$$

here of course $\langle \tilde{\alpha}, v \rangle = \langle \alpha, v \rangle - d$, where $\langle \alpha, v \rangle$ is defined by $\mathbb{R}$ -linearly extending α from a function on X_{*D_m} to $\mathcal{A}_{D_m}$. We also write $s_{i,j;d} := s_{\tilde{\alpha}}$, and we define $s_{i,j} := s_{i,j;0} = s_\alpha$. Note that $s_{i,j}$ acts on vectors as the permutation $(ij)(i^*j^*)$ (expressed in cycle notation) on entries.

For $w \in \widetilde{W}_{D_m}$, it will be important for us to relate the vectors ν_k^w and $\nu_k^{s_{\tilde{\alpha}}w}$. In general, for any $v \in \mathcal{A}_{D_m}$, we have

$$(8.8.2) \quad s_{\tilde{\alpha}}wv - v = wv - v - \langle \tilde{\alpha}, wv \rangle \alpha^\vee.$$

Hence

$$(8.8.3) \quad \begin{aligned} s_\alpha(s_{\tilde{\alpha}}wv - v) &= wv - v - \langle \tilde{\alpha}, wv \rangle \alpha^\vee - (\langle \alpha, wv \rangle - \langle \alpha, v \rangle - 2\langle \tilde{\alpha}, wv \rangle) \alpha^\vee \\ &= wv - v + \langle \tilde{\alpha}, v \rangle \alpha^\vee. \end{aligned}$$

Taking $v = a_k$, we conclude

$$(8.8.4) \quad \nu_k^{s_{\tilde{\alpha}}w} = s_\alpha(\nu_k^w + \langle \tilde{\alpha}, a_k \rangle \alpha^\vee), \quad 0 \leq k \leq m.$$

8.9. Reflections and the Bruhat order. We continue with our element $w \in \widetilde{W}_{D_m}$ from previous subsections. Let $\tilde{\alpha}$ be an affine root with linear part α . In this subsection we give a characterization of when $w < s_{\tilde{\alpha}}w$ in the Bruhat order. We shall simply recall the result from [Sm2], noting that that argument given there is completely general. Recall that A_{D_m} denotes our base alcove in $\mathcal{A}_{D_m}$.

Lemma 8.9.1 ([Sm2, Lem. 5.3.2]). *Let S be any subset of $\overline{A_{D_m}}$, and suppose that*

- $|\langle \alpha, wv - v \rangle| < |\langle \alpha, wv - v + \langle \tilde{\alpha}, v \rangle \alpha^\vee \rangle|$ for some $v \in S$; or
- *there exists $v \in S$ such that $\langle \tilde{\alpha}, v \rangle = 0$ and $\langle \alpha, wv - v \rangle$ is positive or negative according as $\tilde{\alpha}$ is positive or negative on A_{D_m} .*

Then $w < s_{\tilde{\alpha}}w$. The converse holds if S is not contained in a single proper subfacet of A_{D_m} . In particular, $w < s_{\tilde{\alpha}}w \iff$ for some $0 \leq k \leq m$,

- (1) $|\langle \alpha, \nu_k^w \rangle| < |\langle \alpha, \nu_k^w + \langle \tilde{\alpha}, a_k \rangle \alpha^\vee \rangle|$; or
- (2) $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\langle \alpha, \nu_k^w \rangle$ is positive or negative according as $\tilde{\alpha}$ is positive or negative on A_{D_m} . $\square$

8.10. μ -admissibility implies μ -spin-permissibility. It is a general result of Kottwitz–Rapoport [KR, 11.2] that for any cocharacter μ in any root datum, μ -admissibility implies μ -permissibility for elements in the extended affine Weyl group. In this subsection we shall show that for the cocharacter μ (8.0.1) in $\mathcal{R}_{D_m}$, we have the stronger result that μ -admissibility implies μ -spin-permissibility in $\widetilde{W}_{D_m}$. This will prove the “easy” implication in (8.2.2).

We first prove that the set of μ -spin-permissible elements is closed in the Bruhat order.

Proposition 8.10.1. *Let $w, x \in \widetilde{W}_{D_m}$. Suppose that $w \leq x$ in the Bruhat order and that x is μ -spin-permissible. Then w is μ -spin-permissible.*

Proof. Since the μ -permissible set is closed in the Bruhat order by [KR, Lem. 11.3], our characterization of μ -permissibility (8.6.4) immediately reduces us, via an obvious induction argument, to proving the following lemma. $\square$

Lemma 8.10.2. *Let $0 \leq k \leq m$, and let $\tilde{\alpha} := \tilde{\alpha}_{i,j;d}$ for $i \neq j$, j^* be an affine root such that $w < s_{\tilde{\alpha}}w$. Suppose that ν_k^w satisfies (SP1') and that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') and the spin condition. Then ν_k^w satisfies the spin condition.*

Proof. Suppose that $\nu_k^w \in \mathbb{Z}^{2m}$. We must show that if $\nu_k^w(l) \in \{0, 2\}$ for all $l \in A_k$ or for all $l \in B_k$, then $\nu_k^w \equiv \mu \pmod{Q_{D_m}^\vee}$. Let $x := s_{\tilde{\alpha}}w$. By [KR, Lem. 11.4], since $w < x$, the vector ν_k^w lies on the line segment between ν_k^x and $s_{i,j}\nu_k^x$. Hence for any l ,

$$(8.10.3) \quad \nu_k^x(l) \leq \nu_k^w(l) \leq (s_{i,j}\nu_k^x)(l) \quad \text{or} \quad \nu_k^x(l) \geq \nu_k^w(l) \geq (s_{i,j}\nu_k^x)(l).$$

To continue with the proof, let us suppose that $\nu_k^w(l) \in \{0, 2\}$ for all $l \in A_k$; the case that B_k replaces A_k is entirely similar. We consider subcases based on the containment of i and j in A_k . First suppose that $i \in A_k$. Then $\nu_k^w(i) \in \{0, 2\}$. Since ν_k^x satisfies (SP1'), the inequalities (8.10.3) then imply that $\nu_k^w(i) = \nu_k^x(i)$ or $\nu_k^w(i) = (s_{i,j}\nu_k^x)(i)$. If $\nu_k^w(i) = \nu_k^x(i)$, then by examining the i th entries on both sides of the equality

$$\nu_k^w - \nu_k^x = \langle \tilde{\alpha}, wa_k \rangle \alpha_{i,j}^\vee,$$

which is obtained from (8.8.2), we see that $\langle \tilde{\alpha}, wa_k \rangle = 0$. This and the spin condition for ν_k^x then yield $\nu_k^w = \nu_k^x \equiv \mu \pmod{Q_{D_m}^\vee}$, as desired. If $\nu_k^w(i) = (s_{i,j}\nu_k^x)(i)$,

then by examining the i th entries on both sides of

$$s_{i,j}\nu_k^x - \nu_k^w = \langle \tilde{\alpha}, a_k \rangle \alpha_{i,j}^\vee,$$

which is obtained from (8.8.3), we see that $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^w = s_{i,j}\nu_k^x$. Since $\langle \tilde{\alpha}, a_k \rangle \in \mathbb{Z}$, we must have $j \in A_k$ as well. Thus A_k is $s_{i,j}$ -stable and $\nu_k^x(l) = (s_{i,j}\nu_k^w)(l) \in \{0, 2\}$ for all $l \in A_k$. Since ν_k^x satisfies the spin condition, we conclude $\nu_k^w = s_{i,j}\nu_k^x \equiv \nu_k^x \equiv \mu \pmod{Q_{D_m}^\vee}$, as desired.

If $j \in A_k$ then one proves the lemma in a completely similar way.

Finally suppose that $i, j \in B_k$. Then $\langle \tilde{\alpha}, wa_k \rangle \in \mathbb{Z}$, so that $\nu_k^w \equiv \nu_k^x \pmod{Q_{D_m}^\vee}$; and for $l \in A_k$, the inequalities (8.10.3) are all equalities, so that $\nu_k^x(l) = \nu_k^w(l) \in \{0, 2\}$. Hence $\nu_k^x \equiv \mu \pmod{Q_{D_m}^\vee}$ by the spin condition, and we're done. $\square$

Corollary 8.10.4. *Let $w \in \widetilde{W}_{D_m}$. Then w is μ -admissible $\implies w$ is μ -spin-permissible.*

Proof. Since the translation elements t_λ for $\lambda \in S_{2m}^\circ \mu$ are evidently μ -spin-permissible, this follows from (8.10.1). $\square$

8.11. Lemmas on reflections and μ -spin-permissibility. In this subsection we collect a number of technical lemmas for use in the next subsection when we prove that μ -spin-permissibility implies μ -admissibility. We continue with our element $w = t_{\nu_0^w} \sigma \in \widetilde{W}_{D_m}$.

Lemma 8.11.1. *Let $\tilde{\alpha} = \tilde{\alpha}_{i,j;d}$ for some $j \neq i$, i^* be an affine root. Let $0 \leq k \leq m$, and suppose that ν_k^w and $\nu_k^{s_{\tilde{\alpha}} w}$ satisfy (SP1').*

(i) *If $\nu_k^w(i)$, $\nu_k^w(j)$, and $\langle \tilde{\alpha}, a_k \rangle$ are all integers, then*

$$c_k^{s_{\tilde{\alpha}} w} \in \{c_k^w - 2, c_k^w, c_k^w + 2\}.$$

(ii) *If any one of $\nu_k^w(i)$, $\nu_k^w(j)$, or $\langle \tilde{\alpha}, a_k \rangle$ is a half-integer, then*

$$c_k^{s_{\tilde{\alpha}} w} \in \{c_k^w - 1, c_k^w, c_k^w + 1\}.$$

Proof. This is easily verified by explicit case analysis using the formula $\nu_k^{s_{\tilde{\alpha}} w} = s_{i,j}(\nu_k^w + \langle \tilde{\alpha}, a_k \rangle \alpha_{i,j}^\vee)$ from (8.8.4) and our various equivalent expressions for c_k^w in §§8.4–8.5. $\square$

The following lemma will be a key tool for us in the next subsection. It is an analog of [Sm1, Lem. 8.5.2], but rather than taking the time to build up to it systematically, as is done to some extent in [Sm1], we shall give a narrow statement and proof tailored just to what we need later on. Recall our discussion of intervals in §8.7.

Lemma 8.11.2. *Suppose that w is naively μ -permissible, and let $i \in \{1, \dots, 2m\}$ be such that $i < \sigma(i) < i^*$.*

- *Suppose $u(i) = u(\sigma(i))$. If $K_i \subset [\sigma(i), i)$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); 0}$. If $[\sigma(i), i) \subset K_i$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); -1}$.*
- *Suppose $u(i) = 2$ and $u(\sigma(i)) = 1$. If $K_i \subset [\sigma(i), i)$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); 1}$. If $[\sigma(i), i) \subset K_i$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); -1}$.*
- *Suppose $u(i) = 1$ and $u(\sigma(i)) = 2$. If $K_i \subset [\sigma(i), i)$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); 0}$. If $[\sigma(i), i) \subset K_i$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); -2}$.*

Then in every case, $s_{\tilde{\alpha}} w$ is naively μ -permissible and $w < s_{\tilde{\alpha}} w$ in the Bruhat order. If $u(i) = u(\sigma(i))$, then moreover $c_k^{s_{\tilde{\alpha}} w} = c_k^w$ for all k .

Note that one of the inclusions $K_i \subset [\sigma(i), i]$ or $[\sigma(i), i] \subset K_i$ always holds. So for any naively μ -permissible w and element i such that $i < \sigma(i) < i^*$, the lemma gives us an $\tilde{\alpha}$ such that $s_{\tilde{\alpha}}w$ is naively μ -permissible and $w < s_{\tilde{\alpha}}w$. To be clear, when there is an *equality* $K_i = [\sigma(i), i]$, the conclusion of the lemma holds for both choices of $\tilde{\alpha}$.

Proof. We shall proceed by a fairly explicit case analysis, making repeated use of the formula $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}(\nu_k^w + \langle \tilde{\alpha}, a_k \rangle \alpha_{i, \sigma(i)}^\vee)$ (8.8.4). Let $n := \min\{\sigma(i), \sigma(i)^*\}$. It is natural to consider $\nu_k^{s_{\tilde{\alpha}}w}$ and ν_k^w for k separately in the ranges $0 \leq k < i$, $i \leq k < n$, $n \leq k \leq m$ with $n = \sigma(i)$, and $n \leq k \leq m$ with $n = \sigma(i)^*$. In all cases, we have

$$\begin{aligned} \nu_k^w(\sigma(i)) &= u(\sigma(i)) && \text{for } 0 \leq k < i; \\ \nu_k^w(\sigma(i)) &= u(\sigma(i)) - \frac{1}{2} && \text{for } i \leq k < n; \\ \nu_k^w(\sigma(i)) &= u(\sigma(i)) && \text{for } \sigma(i) \leq k \leq m; \text{ and} \\ \nu_k^w(\sigma(i)) &= u(\sigma(i)) - 1 && \text{for } \sigma(i)^* \leq k \leq m. \end{aligned}$$

First suppose that $K_i \subset [\sigma(i), i]$. Then for $0 \leq k < i$ or $\sigma(i) \leq k \leq m$, there is no condition on $\nu_k^w(i)$, i.e. $\nu_k^w(i) \in \{u(i) - 1, u(i) - \frac{1}{2}, u(i)\}$. For $i \leq k < n$, this assumption implies that $\mu_k^w(i) = u(i)$. Hence $\nu_k^w(i) \in \{u(i) - \frac{1}{2}, u(i)\}$. And for $\sigma(i)^* \leq k \leq m$, this assumption implies that $\mu_k^w(i) = \mu_{2m-k}^w(i) = u(i)$. Hence $\nu_k^w(i) = u(i)$.

If $u(i) = u(\sigma(i))$, or if $u(i) = 1$ and $u(\sigma(i)) = 2$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); 0}$. Then for $0 \leq k < i$ or $\sigma(i) \leq k \leq m$, we have $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}\nu_k^w$. Hence $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') because ν_k^w does, and $c_k^{s_{\tilde{\alpha}}w} = c_k^w$. Moreover, by the recursion relation (8.5.4) $\nu_{i-1}^w(i) = \nu_i^w(i) - \frac{1}{2} \leq u(i) - \frac{1}{2}$. Since $\nu_k^w(\sigma(i)) = u(\sigma(i))$, we conclude that $\nu_{i-1}^w(i) < \nu_{i-1}^w(\sigma(i))$. Hence $w < s_{\tilde{\alpha}}w$ by taking $k = i - 1$ in criterion (2) in (8.9.1). For $i \leq k < n$, we have $\langle \tilde{\alpha}, a_k \rangle = -\frac{1}{2}$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}(\nu_k^w - \frac{1}{2}\alpha_{i, \sigma(i)}^\vee)$. Since $\nu_k^w(i) \in \{u(i) - \frac{1}{2}, u(i)\}$ and $\nu_k^w(\sigma(i)) = u(\sigma(i)) - \frac{1}{2}$, we conclude that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). If moreover $u(i) = u(\sigma(i))$, then we see by inspection that $c_k^{s_{\tilde{\alpha}}w} = c_k^w$. For $\sigma(i)^* \leq k \leq m$, we have $\nu_k^w(i) = u(i)$, $\nu_k^w(\sigma(i)) = u(\sigma(i)) - 1$, $\langle \tilde{\alpha}, a_k \rangle = -1$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}(\nu_k^w - \alpha_{i, \sigma(i)}^\vee)$. So $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') by inspection. And if moreover $u(i) = u(\sigma(i))$, then $\nu_k^{s_{\tilde{\alpha}}w} = \nu_k^w$, so that certainly $c_k^{s_{\tilde{\alpha}}w} = c_k^w$.

If $u(i) = 2$ and $u(\sigma(i)) = 1$, then let $\tilde{\alpha} := \tilde{\alpha}_{i, \sigma(i); 1}$. Then for $0 \leq k < i$ or $\sigma(i) \leq k \leq m$, we have $\langle \tilde{\alpha}, a_k \rangle = -1$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}(\nu_k^w - \alpha_{i, \sigma(i)}^\vee)$. So by our assumption on u -values, $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). Moreover, $\nu_{i-1}^w(\sigma(i)) = 1$ and, by the same reasoning as in the previous paragraph, $\nu_{i-1}^w(i) \in \{1, \frac{3}{2}\}$. Hence $w < s_{\tilde{\alpha}}w$ by taking $k = i - 1$ in criterion (1) in (8.9.1). For $i \leq k < n$, we have $\langle \tilde{\alpha}, a_k \rangle = -\frac{3}{2}$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}(\nu_k^w - \frac{3}{2}\alpha_{i, \sigma(i)}^\vee)$. Since $\nu_k^w(i) \in \{\frac{3}{2}, 2\}$ and $\nu_k^w(\sigma(i)) = \frac{1}{2}$, we conclude that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). For $\sigma(i)^* \leq k \leq m$, we have $\nu_k^w(i) = 2$, $\nu_k^w(\sigma(i)) = 0$, $\langle \tilde{\alpha}, a_k \rangle = -2$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma(i)}(\nu_k^w - 2\alpha_{i, \sigma(i)}^\vee)$. Hence $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') by inspection. This completes the proof when $K_i \subset [\sigma(i), i]$.

Now suppose that $[\sigma(i), i] \subset K_i$. Then for $0 \leq k < i$ or $\sigma(i) \leq k \leq m$, this assumption implies that $\mu_k^w(i) = \mu_{2m-k}^w(i) = u(i) - 1$. Hence $\nu_k^w(i) = u(i) - 1$.

For $i \leq k < n$, our assumption implies that $\mu_{2m-k}^w(i) = u(i) - 1$. Hence $\nu_k^w(i) \in \{u(i) - 1, u(i) - \frac{1}{2}\}$. For $\sigma(i)^* \leq k \leq m$, there is no constraint on $\nu_k^w(i)$.

If $u(i) = u(\sigma(i))$, or if $u(i) = 2$ and $u(\sigma(i)) = 1$, then let $\tilde{\alpha} := \tilde{\alpha}_{i,\sigma(i);-1}$. Then for $0 \leq k < i$ or $\sigma(i) \leq k \leq m$, we have $\nu_k^w(i) = u(i) - 1$, $\nu_k^w(\sigma(i)) = u(\sigma(i))$, $\langle \tilde{\alpha}, a_k \rangle = 1$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,\sigma(i)}(\nu_k^w + \alpha_{i,\sigma(i)}^\vee)$. So $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') by inspection. And if moreover $u(i) = u(\sigma(i))$, then $\nu_k^{s_{\tilde{\alpha}}w} = \nu_k^w$, so that certainly $c_k^{s_{\tilde{\alpha}}w} = c_k^w$. For $i \leq k < n$, we have $\langle \tilde{\alpha}, a_k \rangle = \frac{1}{2}$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,\sigma(i)}(\nu_k^w + \frac{1}{2}\alpha_{i,\sigma(i)}^\vee)$. Since $\nu_k^w(i) \in \{u(i) - 1, u(i) - \frac{1}{2}\}$ and $\nu_k^w(\sigma(i)) = u(\sigma(i)) - \frac{1}{2}$, we conclude that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). Moreover, since $\nu_{i-1}^w(i) = u(i) - 1$, we have $\nu_i^w(i) = u(i) - \frac{1}{2}$ by (8.5.4). Hence $w < s_{\tilde{\alpha}}w$ by taking $k = i$ in criterion (1) in (8.9.1). If furthermore $u(i) = u(\sigma(i))$, then we also see by inspection that $c_k^{s_{\tilde{\alpha}}w} = c_k^w$. For $\sigma(i)^* \leq k \leq m$, we have $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,\sigma(i)}\nu_k^w$. Hence $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') because ν_k^w does, and $c_k^{s_{\tilde{\alpha}}w} = c_k^w$.

If $u(i) = 1$ and $u(\sigma(i)) = 2$, then let $\tilde{\alpha} := \tilde{\alpha}_{i,\sigma(i);-2}$. Then for $0 \leq k < i$ or $\sigma(i) \leq k \leq m$, we have $\nu_k^w(i) = 0$, $\nu_k^w(\sigma(i)) = 2$, $\langle \tilde{\alpha}, a_k \rangle = 2$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,\sigma(i)}(\nu_k^w + 2\alpha_{i,\sigma(i)}^\vee)$. Hence $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). For $i \leq k < n$, we have $\langle \tilde{\alpha}, a_k \rangle = \frac{3}{2}$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,\sigma(i)}(\nu_k^w + \frac{3}{2}\alpha_{i,\sigma(i)}^\vee)$. Since $\nu_k^w(i) \in \{0, \frac{1}{2}\}$ and $\nu_k^w(\sigma(i)) = \frac{3}{2}$, we conclude that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). Moreover, by the same reasoning as in the previous paragraph, $\nu_i^w(i) = \frac{1}{2}$. Hence $w < s_{\tilde{\alpha}}w$ by taking $k = i$ in criterion (1) in (8.9.1). Finally, for $\sigma(i)^* \leq k \leq m$ we have $\langle \tilde{\alpha}, a_k \rangle = 1$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,\sigma(i)}(\nu_k^w + \alpha_{i,\sigma(i)}^\vee)$. So $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') by our assumption on u -values. This exhausts all the cases we have to consider and completes the proof of the lemma. $\square$

Lemma 8.11.3. *Suppose that w is naively μ -permissible, and let i be as in the preceding lemma. Let $\tilde{\alpha}$ be an affine root prescribed by the preceding lemma such that $s_{\tilde{\alpha}}w$ is naively μ -permissible and $w < s_{\tilde{\alpha}}w$. Suppose that $u(i) \neq u(\sigma(i))$ and that $\nu_k^w(i)$ and $\nu_k^w(\sigma(i))$ are half-integers for some k . Then $c_k^{s_{\tilde{\alpha}}w} = c_k^w + 1$.*

Proof. Our assumption $u(i) \neq u(\sigma(i))$ requires that for one element $a \in \{i, \sigma(i)\}$ we have $\nu_i^w(a) = \frac{3}{2}$, and for the other element $b \in \{i, \sigma(i)\}$ we have $\nu_i^w(b) = \frac{1}{2}$. Since $\nu_k(\sigma(i))$ is a half-integer we moreover have $i \leq k < \min\{\sigma(i), \sigma(i)^*\}$. Thus we see from the formula (8.8.4) and the explicit possibilities for $\tilde{\alpha}$ in (8.11.2) that

$$\nu_k^{s_{\tilde{\alpha}}w} = s_{a,b}(\nu_k^w + \varepsilon \alpha_{a,b}^\vee) \quad \text{for } \varepsilon \in \left\{-\frac{3}{2}, \frac{1}{2}\right\}.$$

The conclusion now follows by inspection. $\square$

Lemma 8.11.4. *Let $0 \leq k \leq m$, and suppose that ν_k^w satisfies (SP1') and (SP3'). For $j \neq i, i^*$, let $\tilde{\alpha} = \tilde{\alpha}_{i,j;d}$ be an affine root such that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'). Suppose that $\nu_k^{s_{\tilde{\alpha}}w}$ fails (SP3') and $\langle \tilde{\alpha}, a_k \rangle \in \mathbb{Z}$. Then*

$$\nu_k^w \in \mathbb{Z}^{2m}, \quad \langle \tilde{\alpha}, a_k \rangle = \pm 1, \quad \nu_k^w(i) = \nu_k^w(j) = 1, \quad \text{and} \quad c_k^w \not\equiv q \pmod{2}.$$

Proof. For $\nu_k^{s_{\tilde{\alpha}}w}$ to fail the spin condition, we must have $\nu_k^{s_{\tilde{\alpha}}w} \in \mathbb{Z}^{2m}$ and $c_k^{s_{\tilde{\alpha}}w} \not\equiv q \pmod{2}$. Since $\langle \tilde{\alpha}, a_k \rangle \in \mathbb{Z}$, the formula (8.8.3) makes clear that $\nu_k^w \in \mathbb{Z}^{2m}$, and moreover that

$$\nu_k^w \equiv s_{i,j}\nu_k^{s_{\tilde{\alpha}}w} \equiv \nu_k^{s_{\tilde{\alpha}}w} \pmod{Q_{D_m}^\vee}.$$

Hence $c_k^w \not\equiv q \pmod{2}$ by (8.4.2). Since ν_k^w satisfies the spin condition, there then exist $l_1 \in A_k$ and $l_2 \in B_k$ such that $\nu_k^w(l_1) = \nu_k^w(l_2) = 1$. Since $\langle \tilde{\alpha}, a_k \rangle \in \mathbb{Z}$, we

also see that i and j are both in A_k or both in B_k . Without loss of generality, let us assume the former. Then $\nu_k^{s_{\tilde{\alpha}}w}(j_2) = 1$, since ν_k^w and $\nu_k^{s_{\tilde{\alpha}}w}$ can differ only in their i, i^*, j , and j^* entries. Since $\nu_k^{s_{\tilde{\alpha}}w}$ fails the spin condition, we conclude that $\nu_k^{s_{\tilde{\alpha}}w}(l) \in \{0, 2\}$ for all $l \in A_k$. Hence $l_1 \in \{i, i^*, j, j^*\}$ and $\langle \tilde{\alpha}, a_k \rangle \neq 0$. Without loss of generality, let us assume $l_1 = i$. In view of (8.8.4), the fact that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') forces $\langle \tilde{\alpha}, a_k \rangle = \pm 1$. And since $\nu_k^{s_{\tilde{\alpha}}w}(j) \in \{0, 2\}$, we conclude from this that $\nu_k^w(j) = 1$ as well, which completes the proof. $\square$

Lemma 8.11.5. *Let $0 \leq k \leq m$, and suppose that ν_k^w is μ -spin-permissible. For $j \neq i, i^*$, let $\tilde{\alpha} := \tilde{\alpha}_{i,j;d}$ be an affine root such that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') but not (SP2') or (SP3'). Then*

$$\nu_k^w(i), \nu_k^w(j) \in \left\{ \frac{1}{2}, 1, \frac{3}{2} \right\}.$$

Proof. We shall show that if $\nu_k^w(i) \in \{0, 2\}$ or $\nu_k^w(j) \in \{0, 2\}$, then $\nu_k^{s_{\tilde{\alpha}}w}$ is μ -spin-permissible. If $\langle \tilde{\alpha}, a_k \rangle \in \mathbb{Z}$, then the previous lemma shows that ν_k^w satisfies (SP3'). And if $\langle \tilde{\alpha}, a_k \rangle$ is a half-integer, then (8.8.4) shows that $\nu_k^{s_{\tilde{\alpha}}w} \notin \mathbb{Z}^{2m}$ so that it trivially satisfies (SP3'). So it remains to show that ν_k^w satisfies (SP2'). But under our present hypotheses, inspection of (8.8.4) yields easily that $c_k^{s_{\tilde{\alpha}}w} \leq c_k^w$, which is $\leq q$ by assumption. $\square$

Lemma 8.11.6. *Assume that ν_k^w satisfies (SP1') and (SP2') for all $0 \leq k \leq m$. Let $i < j < j^* < i^*$ be elements in $\{1, \dots, 2m\}$ such that for all $i \leq k < j$,*

- (1) $\nu_k^w(i) \leq 1$ and $\nu_k^w(j) \geq 1$;
- (2) $\nu_k^w(i) \geq \frac{1}{2}$ and $\nu_k^w(j) \leq \frac{3}{2}$; and
- (3) $c_k^w < q$.

Let $\tilde{\alpha} := \tilde{\alpha}_{i,j;0}$. Then $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') and (SP2') for all $0 \leq k \leq m$, and $w < s_{\tilde{\alpha}}w$. If moreover ν_k^w satisfies (SP3') for k in the range $0 \leq k < i$ or $j \leq k \leq m$, then so does $\nu_k^{s_{\tilde{\alpha}}w}$.

Note that if i (resp. j) is proper, then the inequality $\nu_k^w(i) \leq 1$ (resp. $\nu_k^w(j) \geq 1$) in condition (1) is implied by $u(i) = 1$ (resp. $u(j) = 2$).

Proof. We shall appeal to the formula (8.8.4) for $\nu_k^{s_{\tilde{\alpha}}w}$. If $0 \leq k < i$ or $j \leq k \leq m$, then $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j}\nu_k^w$, which satisfies (SP1') and (SP2') because ν_k^w does. If ν_k^w satisfies (SP3'), then this also implies that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP3') by (8.11.4). If $i \leq k < j$, then $\langle \tilde{\alpha}, a_k \rangle = -\frac{1}{2}$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j}(\nu_k^w - \frac{1}{2}\alpha_{i,j}^\vee)$. So we see immediately from hypothesis (2) that ν_k^w satisfies (SP1'). And since $\langle \tilde{\alpha}, a_k \rangle$ is a half-integer, (8.11.1)(ii) and hypothesis (3) together imply $c_k^{s_{\tilde{\alpha}}w} \leq q$, so that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). To finish, hypothesis (1) immediately implies that criterion (1) in (8.9.1) is satisfied for every k in the range $i \leq k < j$. So $w < s_{\tilde{\alpha}}w$. $\square$

8.12. μ -spin-permissibility implies μ -admissibility. As discussed at the end of §8.2, the implication μ -spin-permissible $\implies \mu$ -admissible in (8.2.2) is an immediate consequence of part (i) of the following proposition, whose proof we are now ready to give.

Proposition 8.12.1. *Suppose that $w \in \widetilde{W}_{D_m}$ is μ -spin-permissible and not a trans-
lation element.*

- (i) *There exists an affine reflection $s_{\tilde{\alpha}}$ such that $s_{\tilde{\alpha}}w$ is μ -spin-permissible and $w < s_{\tilde{\alpha}}w$.*

- (ii) If moreover $\nu_0^w(m) = 1$, then the reflection $s_{\tilde{\alpha}}$ in (i) can be chosen such that $\nu_0^{s_{\tilde{\alpha}}w}(m) = 1$.

Note that part (ii) of the proposition is not needed for the proof of (8.2.2). We shall use it to deduce (9.6.1) below, which is the type B analog of (8.2.2).

Proof. Let us begin by reviewing notation. Attached to w are the vectors μ_k^w (8.1.5) for $0 \leq k \leq 2m$ and ν_k^w (8.5.2) for $0 \leq k \leq m$. We decompose w as a product $t_{\nu_0^w}\sigma$, where $t_{\nu_0^w}$ is the translation part of w on the left and σ is the linear part of w , which we regard as a permutation of $\{1, \dots, 2m\}$. We recall the sets A_k and B_k (4.3.6) for $0 \leq k \leq m$; the upper value $u(i)$ (8.3.3) for $1 \leq i \leq 2m$; the integer c_k^w (8.4.1) for $0 \leq k \leq m$; the point a_k for $0 \leq k \leq m$ (8.1.4, 8.5.1); and the affine root $\tilde{\alpha}_{i,j;d}$ (8.8.1) and coroot $\alpha_{i,j}^\vee$ (8.1.2) for $1 \leq i, j \leq 2m$ and $j \neq i, i^*$. Given an affine root $\tilde{\alpha}$ with linear part α , we shall make repeated use of the formula $\nu_k^{s_{\tilde{\alpha}}w} = s_{\alpha}(\nu_k^w + \langle \tilde{\alpha}, a_k \rangle \alpha^\vee)$ (8.8.4).

Since w is not a translation element, there exists a proper element in $\{1, \dots, 2m\}$. Let i denote the *minimal* proper element. Then

$$\nu_0^w = \dots = \nu_{i-1}^w \neq \nu_i^w.$$

Throughout the proof, let

$$n := \min\{\sigma(i), \sigma(i)^*\}.$$

Before continuing, we record a subsidiary lemma for later use.

Lemma 8.12.2. *The upper values $u(i)$ and $u(\sigma(i))$ are distinct $\iff c_i^w \not\equiv q \pmod{2}$. In particular, if these equivalent conditions hold then $c_i^w < q$.*

Proof. Everything follows from the facts (8.4.3), $c_{i-1}^w = c_0^w \equiv q \pmod{2}$ (SP3'), and $c_i^w \leq q$ (SP2'). $\square$

We return to the proof of (8.12.1). We shall first prove part (i). Of course either $\sigma(i) = i^*$ or $\sigma(i) \neq i^*$, and we shall divide the proof into these two cases.

Case I: $\sigma(i) = i^*$. Then $\nu_i^w = \nu_0^w + e_i - e_{i^*} \in \mathbb{Z}^{2m}$ by (8.5.4), and $c_i^w \not\equiv q \pmod{2}$ by (8.12.2). We shall consider the subcases $u(i) = 1$ and $u(i) = 2$. In each subcase we shall use in a crucial way that ν_i^w satisfies the spin condition.

First suppose that $u(i) = 1$. Then $\nu_k^w(i) = 0$ for $0 \leq k < i$ and $\nu_k^w(i) = 1$ for $i \leq k \leq m$. By the spin condition, there exists a minimal $j \in B_i$ such that $\nu_i^w(j) = 1$. Of course $n \neq j, j^*$, and therefore $\nu_k(j) = \nu_k(j^*) = 1$ for all $0 \leq k \leq i$. Take

$$\tilde{\alpha} := \tilde{\alpha}_{i,j^*;-1}.$$

We must show that $\nu_k^{s_{\tilde{\alpha}}w}$ is μ -spin-permissible for all $0 \leq k \leq m$. First let $0 \leq k < i$. Then $\nu_k^w(i) = 0$, $\nu_k^w(j^*) = 1$, $\langle \tilde{\alpha}, a_k \rangle = 1$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \alpha_{i,j^*}^\vee) = \nu_k^w$, which is μ -spin-permissible. Next let $i \leq k < j$. Then $\nu_k^w(i) = 1$, $\nu_k^w(j^*) \in \{\frac{1}{2}, 1, \frac{3}{2}\}$ by (8.7.2), $\langle \tilde{\alpha}, a_k \rangle = \frac{1}{2}$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \frac{1}{2}\alpha_{i,j^*}^\vee)$. So $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') by inspection, and $c_k^{s_{\tilde{\alpha}}w} \leq c_k^w + 1$ by (8.11.1)(ii). Now, since $k < j$ and j was taken to be *minimal*, it follows from (8.4.4)(i) that $c_i^w \geq c_k^w$. Hence

$$c_k^{s_{\tilde{\alpha}}w} \leq c_k^w + 1 \leq c_i^w + 1 \leq q,$$

where the last inequality uses (8.12.2). So $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). Moreover $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies the spin condition since, for example, $\nu_k^{s_{\tilde{\alpha}}w}(j^*) = \frac{3}{2}$ is a half-integer. In addition, since $\nu_i^w(i) = \nu_i^w(j^*) = 1$, we see by taking $k = i$ in criterion

(1) in (8.9.1) that $w < s_{\tilde{\alpha}}w$. Finally let $i \leq k \leq m$. Then $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*} \nu_k^w$, which satisfies (SP1') and (SP2') by inspection and (SP3') by (8.11.4).

Now suppose that $u(i) = 2$. Then $\nu_k^w(i) = 1$ for $0 \leq k < i$ and $\nu_k^w(i) = 2$ for $i \leq k \leq m$. By the spin condition, there exists an element $j \leq i$ such that $\nu_i^w(j) = 1$. Evidently $j \neq i$, and therefore $\nu_k(j) = 1$ for all k since i is the minimal proper element. Take

$$\tilde{\alpha} := \tilde{\alpha}_{j,i;0}.$$

We shall show that $w < s_{\tilde{\alpha}}w$ and that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') and (SP2') for all $0 \leq k \leq m$ by applying (8.11.6) (with the roles of i and j reversed). Indeed, for $j \leq k < i$ we have $\nu_k^w(j) = \nu_k^w(i) = 1$, so that conditions (1) and (2) in (8.11.6) are satisfied. And by (8.4.3) $c_k^w < c_k^w + 1 = c_i^w \leq q$, so that condition (3) is satisfied as well. It remains to show that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies the spin condition for all k . For $0 \leq k < j$ and $i \leq k \leq m$, this is done by (8.11.4). And for $j \leq k < i$, we have $\langle \tilde{\alpha}, a_k \rangle = -\frac{1}{2}$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j}(\nu_k^w - \frac{1}{2}\alpha_{i,j}^\vee)$. Hence $\nu_k^{s_{\tilde{\alpha}}w}(j) = \frac{1}{2}$ is a half-integer and $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies the spin condition.

This completes the proof in Case I. We now turn to Case II, which will be much more involved.

Case II: $\sigma(i) \neq i^*$. In this case, by minimality of i , we have $i < \sigma(i), \sigma(i)^* < i^*$ and

$$\nu_0^w(i) = \cdots = \nu_{i-1}^w(i) = u(i) - 1 \quad \text{and} \quad \nu_i^w(i) = u(i) - \frac{1}{2}.$$

Moreover $\nu_k^w(\sigma(i))$ takes the constant value $u(\sigma(i))$ equal to 1 or 2 for $0 \leq k < i$; the constant value $u(\sigma(i)) - \frac{1}{2}$ for $i \leq k < n$; and the constant value $u(\sigma(i))$ or $u(\sigma(i)) - 1$ for $n \leq k \leq m$ according as $n = \sigma(i)$ or $n = \sigma(i)^*$.

By (8.11.2), the element $w' := s_{\tilde{\alpha}'}w$ is at least naively μ -permissible and satisfies $w < w'$ for some $\tilde{\alpha}' := \tilde{\alpha}_{i,\sigma(i);d}$ with $d \in \{-1, 0\}$ if $u(i) = u(\sigma(i))$; $d \in \{-1, 1\}$ if $u(i) > u(\sigma(i))$; and $d \in \{-2, 0\}$ if $u(i) < u(\sigma(i))$. If w' is μ -spin-permissible then we are done. So let us suppose it is not. Our problem is to find another affine root $\tilde{\alpha}$ such that $s_{\tilde{\alpha}}w$ is μ -spin-permissible and $w < s_{\tilde{\alpha}}w$. Before continuing with the main proof in Case II, we shall pause to prove some subsidiary lemmas.

Lemma 8.12.3. $\nu_k^{w'}$ is μ -spin-permissible for all $0 \leq k \leq i$. If moreover $u(i) = u(\sigma(i))$, then $\nu_k^{w'}$ is μ -spin-permissible for all $n \leq k \leq m$.

Proof. First let $0 \leq k < i$. Then $\langle \tilde{\alpha}', a_k \rangle = -d \in \mathbb{Z}$ and $c_k^w = c_0^w \equiv q \pmod{2}$. So $\nu_k^{w'}$ satisfies the spin condition by (8.11.4). So it remains to show that $\nu_k^{w'}$ satisfies (SP2'), that is, that $c_k^{w'} \leq q$. We have

$$\nu_k^{w'} = s_{i,\sigma(i)}(\nu_k^w - d\alpha_{i,\sigma(i)}^\vee).$$

Since $\nu_k^w \in \mathbb{Z}^{2m}$, we have $c_k^{w'} \in \{c_k^w - 2, c_k^w, c_k^w + 2\}$ by (8.11.1)(i). Thus $c_k^{w'} \leq c_k^w \leq q$ unless $c_k^{w'} = c_k^w + 2$, which occurs when $d = \pm 1$ and $\nu_k^w(i) = \nu_k^w(\sigma(i)) = 1$. In this case $u(i) = 2$ and $u(\sigma(i)) = 1$, and $c_k^w + 2 = c_i^w + 1 \leq q$ by (8.4.3) and (8.12.2), as desired.

The case $k = i$ requires a separate argument. If $u(i) = u(\sigma(i))$, then $c_i^{w'} = c_i^w \equiv q \pmod{2}$ by (8.11.2) and (8.12.2). Hence $\nu_i^{w'}$ satisfies (SP2') and (SP3'). If $u(i) \neq u(\sigma(i))$, then $c_i^{w'} = c_i^w + 1$ by (8.11.3). It follows immediately from this and (8.12.2) that $c_k^{w'} \leq q$ and $c_k^{w'} \equiv q \pmod{2}$, so that $\nu_i^{w'}$ satisfies (SP2') and (SP3').

Finally suppose that $u(i) = u(\sigma(i))$, and let $n \leq k \leq m$. By (8.11.2), $c_k^{w'} = c_k^w \leq q$, so that $\nu_k^{w'}$ satisfies (SP2'). Moreover $\langle \tilde{\alpha}', a_k \rangle \in \mathbb{Z}$, and it follows from (8.11.4) that if $\nu_k^{w'}$ fails (SP3'), then $c_k^{w'} = c_k^w + 2$. Since $c_k^{w'} = c_k^w$, we conclude that (SP3') holds. $\square$

Lemma 8.12.4. *Suppose that $c_i^w = c_k^w$ for some $0 \leq k \leq m$, and that $u(i) = u(\sigma(i))$ or $k < n$. Then $\nu_k^{w'}$ is μ -spin-permissible.*

Proof. First suppose that $u(i) = u(\sigma(i))$. Then by (8.11.2) we have $c_k^{w'} = c_k^w = c_i^w$. This is $\leq q$ because ν_i^w satisfies (SP2'), and $\equiv q \pmod{2}$ by (8.12.2). Thus $\nu_k^{w'}$ is μ -spin-permissible.

Now suppose that $u(i) \neq u(\sigma(i))$ and $k < n$. For $k \leq i$ the conclusion is given by (8.12.3), so assume $i < k < n$. Then $\langle \tilde{\alpha}', a_k \rangle$ is a half-integer and $c_k^{w'} \in \{c_k^w - 1, c_k^w, c_k^w + 1\}$ by (8.11.1)(ii). Since $c_k^w = c_i^w < q$ by (8.12.2), we conclude that $\nu_k^{w'}$ satisfies (SP2'). To see that $\nu_k^{w'}$ also satisfies the spin condition, assume that it has only integer entries. Then, since $\langle \tilde{\alpha}', a_k \rangle$ is a half-integer, (8.8.4) requires that $\nu_k^w(i)$ and $\nu_k^w(\sigma(i))$ are half-integers. Hence $c_k^{w'} = c_k^w + 1$ by (8.11.3). But $c_k^w = c_i^w \not\equiv q \pmod{2}$ by (8.12.2), and therefore $c_k^{w'} \equiv q \pmod{2}$. So $\nu_k^{w'}$ satisfies (SP3'). $\square$

Lemma 8.12.5. *Suppose that $\nu_p^{w'}$ is not μ -spin-permissible for some $0 \leq p \leq m$.*

- (i) *If $c_i^w > c_p^w$, then $i < p < n$ and $\nu_p^{w'}$ satisfies (SP2') but not (SP3').*
- (ii) *If $c_i^w \leq c_p^w$, then $c_i^w < q$.*
- (iii) *If $p \geq n$, then $\nu_p^w(\sigma(i)) = 1$, and*

$$u(\sigma(i)) = 1 \text{ and } n = \sigma(i) \quad \text{or} \quad u(\sigma(i)) = 2 \text{ and } n = \sigma(i)^*.$$

If furthermore $c_i^w = c_p^w$, then $\nu_p^w(i) = 1$.

Proof. We first prove (i). To see that $\nu_p^{w'}$ satisfies (SP2'), note that if $u(i) = u(\sigma(i))$, then by (8.11.2) $c_p^{w'} = c_p^w \leq q$. And if $u(i) \neq u(\sigma(i))$, then $c_p^w < c_i^w < q$ by (8.12.2). So by (8.11.1) $c_p^{w'} \leq c_p^w + 2 \leq q$. So either way (SP2') is satisfied.

It remains to show that $i < p < n$. For this, we already know from (8.12.3) that $i < p$. So to finish the proof of (i), it suffices to show that if $n \leq k \leq m$ and $c_i^w > c_k^w$, then $\nu_k^{w'}$ is μ -spin-permissible. If $u(i) = u(\sigma(i))$, then this is done by (8.12.3). So assume $u(i) \neq u(\sigma(i))$. We just showed in the previous paragraph that $\nu_k^{w'}$ satisfies (SP2'). To see that $\nu_k^{w'}$ also satisfies the spin condition, note that since $n \leq k$, we have $\langle \tilde{\alpha}', a_k \rangle \in \mathbb{Z}$. So by (8.11.4), we might as well assume that

$$\nu_k^w \in \mathbb{Z}^{2m}, \quad \nu_k^w(i) = \nu_k^w(\sigma(i)) = 1, \quad \text{and} \quad c_k^w \not\equiv q \pmod{2}.$$

Since ν_k^w satisfies the spin condition and $i, i^*, \sigma(i), \sigma(i)^* \in A_k$, there must exist an element $j \in B_k$ such that $\nu_k^w(j) = 1$. Now, since $u(i) \neq u(\sigma(i))$, we also have $c_i^w \not\equiv q \pmod{2}$ by (8.12.2). Since $c_i^w > c_k^w$, we conclude that $c_i^w \geq c_k^w + 2$. Let us now compare the entries of μ_i^w and μ_k^w . If $u(i) = 2$ and $u(\sigma(i)) = 1$, then

$$\mu_i^w(i) = 2, \quad \mu_i^w(i^*) = 1, \quad \mu_i^w(\sigma(i)) = 0, \quad \text{and} \quad \mu_i^w(\sigma(i)^*) = 1.$$

If $u(i) = 1$ and $u(\sigma(i)) = 2$, then

$$\mu_i^w(i) = 1, \quad \mu_i^w(i^*) = 2, \quad \mu_i^w(\sigma(i)) = 1, \quad \text{and} \quad \mu_i^w(\sigma(i)^*) = 0.$$

Either way, since $\mu_k^w(i) = \mu_k^w(i^*) = \mu_k^w(\sigma(i)) = \mu_k^w(\sigma(i)^*) = 1$ and $c_i^w \geq c_k^w + 2$, there must exist another element $l \notin \{i, i^*, \sigma(i), \sigma(i)^*\}$ such that $\mu_i^w(l) = 0$ and $\mu_k^w(l) = 1$. By the recursion relation (8.3.2) $l \in \{i+1, \dots, k\} \subset A_k$. And since $\nu_k^w \in \mathbb{Z}^{2m}$, we must have $\nu_k^w(l) = \mu_k^w(l) = 1$. Since $\nu_k^{w'}(l) = \nu_k^w(l) = 1$ and $\nu_k^{w'}(j) = \nu_k^w(j) = 1$, we conclude that $\nu_k^{w'}$ satisfies (SP3'), which completes (i).

We next prove (ii). If $c_i^w < c_p^w$ then the conclusion is clear since $c_p^w \leq q$. And if $c_i^w = c_p^w$ then $u(i) \neq u(\sigma(i))$ by (8.12.4). Hence (8.12.2) gives the conclusion.

We finally prove (iii). For $p \geq n$ the $\sigma(i)$ -entry of ν_p^w is an integer, which, since $\nu_p^{w'}$ is not μ -spin-permissible, must equal 1 by (8.11.5). This immediately implies that $u(\sigma(i)) = 1$ and $n = \sigma(i)$, or $u(\sigma(i)) = 2$ and $n = \sigma(i)^*$.

To complete (iii), note that if $\nu_p^{w'}$ fails (SP3'), then (8.11.4) gives $\nu_p^w(i) = 1$ independently of the values of c_i^w and c_p^w . If moreover $c_i^w = c_p^w$, then $u(i) \neq u(\sigma(i))$ by (8.12.4). Hence $c_p^w \not\equiv q \pmod{2}$ by (8.12.2). If $\nu_p^{w'}$ fails (SP2'), then this, (8.11.1), and the inequality $c_p^w \leq q$ imply that $c_p^w = q - 1$ and $c_p^{w'} = q + 1$. By (8.11.1) and (8.11.5) this requires $\nu_p^w(i) = 1$. $\square$

Our subsidiary lemmas now dispensed with, we return to the main proof of the proposition in Case II. Since w' is naively μ -permissible but not μ -spin-permissible, there exists a minimal $p \in \{1, \dots, m\}$ such that $\nu_p^{w'}$ fails (SP2') or (SP3'). By (8.12.3) $p > i$. Of course $c_i^w \leq c_p^w$ or $c_i^w > c_p^w$, and we shall consider each of these possibilities separately.

First suppose $c_i^w \leq c_p^w$. We claim that there exists $j' \in \{i+1, \dots, p\}$ such that $\mu_i(j') = 1$ and $\mu_p(j') = 2$. If $c_i^w < c_p^w$ then (8.4.4)(i) furnishes exactly the claim. If $c_i^w = c_p^w$ then note that $\mu_i^w(i) = 2$ or $\mu_i^w(i^*) = 2$ according as $u(i) = 2$ or $u(i) = 1$. By (8.12.4) $p \geq n$, and then by (8.12.5)(iii) $\mu_p^w(i) = \mu_p^w(i^*) = 1$. Since $c_i^w = c_p^w$, there therefore exists a j' such that $\mu_i^w(j') = 1$ and $\mu_p^w(j') = 2$. By the recursion relation (8.3.2) such a j' must be contained in $\{i+1, \dots, p\}$, which proves the claim.

Now, j' is clearly an element of the set

$$(8.12.6) \quad S := \{j' \in \{i+1, \dots, p\} \mid \mu_i^w(j') = 1 \text{ and } \mu_{j'}^w(j') = 2\}.$$

Let j denote the *minimal* element of S . Evidently $u(j) = 2$, so that $u(j^*) = 1$. Since i is the minimal proper element and $j \neq n$, we have $\nu_k^w(j) = \nu_k^w(j^*) = 1$ for all $0 \leq k \leq i$.

We claim that $j \neq n$. To prove this we might as well assume $n \leq j$. Then $n \leq p$. Then by (8.12.5)(iii) $u(n) = 1$. But $u(j) = 2$, which proves the claim.

Still supposing $c_i^w \leq c_p^w$, we are now ready to prescribe our affine root $\tilde{\alpha}$, although to do so we will need to consider some further subcases. First suppose $j < n$ and $u(i) = 2$. Then we take

$$\tilde{\alpha} := \tilde{\alpha}_{i,j^*;-1}.$$

To see that $s_{\tilde{\alpha}}w$ is μ -spin-permissible, first let $0 \leq k < i$. Then $\nu_k^w(i) = \nu_k^w(j^*) = 1$, $\langle \tilde{\alpha}, a_k \rangle = 1$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \alpha_{i,j^*}^\vee)$. Thus by inspection $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') and $c_k^{s_{\tilde{\alpha}}w} = c_k^w + 2$, and of course $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies the spin condition by (8.11.4). To see that $\nu_k^{s_{\tilde{\alpha}}w}$ also satisfies (SP2'), consider the inequalities

$$c_k^w = c_{i-1}^w \leq c_i^w < q;$$

here the first inequality uses that $\mu_{i-1}^w(i) = 1$ and $\mu_i^w(i) = 2$, and the second is (8.12.5)(ii). Hence $c_k^{s_{\tilde{\alpha}}w} = c_k^w + 2 \leq q$ by parity. So indeed $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). Now let $i \leq k < j$. Then $\langle \tilde{\alpha}, a_k \rangle = \frac{1}{2}$, $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \frac{1}{2}\alpha_{i,j^*}^\vee)$, and $c_k^{s_{\tilde{\alpha}}w} \leq c_k^w + 1$

by (8.11.1)(ii). Since $\nu_p^w(i) \in \{1, \frac{3}{2}\}$ by (8.11.5), we must have $\nu_k^w(i) \in \{1, \frac{3}{2}\}$ by (8.7.2). And since $\nu_i^w(j^*) = 1$, $u(j^*) = 1$, and $k < j$, we must have $\nu_k^w(j^*) \in \{\frac{1}{2}, 1\}$ again by (8.7.2). Therefore we see the following by inspection: that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'); that $\nu_k^{s_{\tilde{\alpha}}w}(\sigma(i)) = \nu_k^w(\sigma(i))$ is a half-integer, so that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP3'); and that

$$\nu_i^w(i) = \frac{3}{2}, \quad \nu_i^w(j^*) = 1, \quad \nu_i^{s_{\tilde{\alpha}}w}(i) = 2, \quad \text{and} \quad \nu_i^{s_{\tilde{\alpha}}w}(j^*) = \frac{1}{2},$$

so that $w < s_{\tilde{\alpha}}w$ by criterion (1) in (8.9.1). We must still show that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). For this, since $i \leq k < j$ and j is the *minimal* element $\geq i+1$ such that $\mu_i(j) = 1$ and $\mu_j(j) = 2$, we have $c_i^w \geq c_k^w$. Hence

$$c_k^{s_{\tilde{\alpha}}w} \leq c_k^w + 1 \leq c_i^w + 1 \leq q,$$

where the last inequality uses (8.12.5)(ii). So indeed $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). Finally let $j \leq k \leq m$. Then $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*} \nu_k^w$ is clearly μ -spin-permissible, using (8.11.4) for (SP3'). Thus $\tilde{\alpha}$ solves our problem when $j < n$, $u(i) = 2$, and $c_i^w \leq c_p^w$.

Now suppose $j < n$ and $u(i) = 1$. Then we take

$$\tilde{\alpha} := \tilde{\alpha}_{i,j;0}.$$

We shall show that $w < s_{\tilde{\alpha}}w$ and that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') and (SP2') for all $0 \leq k \leq m$ by applying (8.11.6). We must show that conditions (1)–(3) are satisfied. Condition (1) is satisfied because $u(i) = 1$ and $u(j) = 2$. For condition (2), note that $\nu_i^w(i) = \frac{1}{2}$ and $\nu_p^w(i) \in \{\frac{1}{2}, 1\}$ by (8.11.5) (applied with $\tilde{\alpha} = \tilde{\alpha}'$). Therefore, by (8.7.2), $\nu_k^w(i) \in \{\frac{1}{2}, 1\}$ for all $i \leq k \leq p$, and in particular for all $i \leq k < j$. And since $\nu_i^w(j) = 1$, we have again by (8.7.2) that $\nu_i^w(j) \leq \frac{3}{2}$ for all $i \leq k < j$. So indeed condition (2) holds. For condition (3), let $i \leq k < j$. Then $c_k^w \leq c_i^w$ by minimality of j , and $c_i^w < q$ by (8.12.5)(ii), which gives us exactly what we need. So (8.11.6) applies. To show that $\tilde{\alpha}$ solves our problem, it remains to show that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies the spin condition for all k . As always, by (8.11.4) we may restrict our attention to $i \leq k < j$. Since by assumption $j < n$, we have that $\nu_k^{s_{\tilde{\alpha}}w}(\sigma(i)) = \nu_k^w(\sigma(i))$ and that $\nu_k^w(\sigma(i))$ is a half-integer. Hence $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies (SP3'). This completes the subcase $j < n$, $u(i) = 1$, and $c_i^w \leq c_p^w$.

Still supposing $c_i^w \leq c_p^w$, now suppose $j > n$. Then $p > n$, so that $u(i) \neq u(\sigma(i))$ by (8.12.3). We take

$$\tilde{\alpha} := \tilde{\alpha}_{n,j;0}.$$

As in the subcase considered in the previous paragraph, we shall first apply (8.11.6) to show that $w < s_{\tilde{\alpha}}w$ and that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1') and (SP2') for all k . Condition (1) in (8.11.6) is satisfied because $u(j) = 2$ and because (8.12.5)(iii) implies that $u(n) = 1$. For condition (2), the value $\nu_k^w(n)$ is constant for varying k with $n \leq k \leq m$. Hence $\nu_k^w(n) = \nu_p^w(n)$, which in turn equals 1 by (8.12.5)(iii), for $n \leq k < j$. And since $\nu_i^w(j) = 1$, we have by (8.7.2) that $\nu_i^w(j) \leq \frac{3}{2}$ for all $i \leq k < j$, and in particular for all $n \leq k < j$. So indeed condition (2) holds. For condition (3), since p is *minimal* such that $\nu_p^{w'}$ is not μ -spin-permissible, we have $c_k^{w'} \leq q$ for all $k < p$, and in particular for all $n \leq k < j$. To make use of this, let us compare c_k^w and $c_k^{w'}$. Since $u(i) \neq u(\sigma(i))$, by (8.11.2) and (8.12.5)(iii) we have the possibilities $u(\sigma(i)) = 1$, $\sigma(i) \leq m$, and $d \in \{-1, 1\}$; or $u(\sigma(i)) = 2$, $\sigma(i) > m$, and $d \in \{-2, 0\}$. Either way $\langle \tilde{\alpha}', a_k \rangle = \pm 1$ and $\nu_k^{w'} = s_{i,\sigma(i)}(\nu_k^w \pm \alpha_{i,\sigma(i)}^\vee)$ for $n \leq k \leq m$, and in

particular for $n \leq k < j$. Now, $\nu_i(i) \in \{\frac{1}{2}, \frac{3}{2}\}$, and by (8.11.5) $\nu_p^w(i) \in \{\frac{1}{2}, 1, \frac{3}{2}\}$. Hence by (8.7.2) $\nu_k^w(i) \in \{\frac{1}{2}, 1, \frac{3}{2}\}$ for all $i \leq k \leq p$, and in particular for all $n \leq k < j$. And we have already observed that $\nu_k^w(n) = 1$ for $n \leq k < j$. Thus we see explicitly that $c_k^{w'} \in \{c_k^w + 1, c_k^w + 2\}$. Hence $c_k^w < c_k^{w'} \leq q$. So condition (3) holds and (8.11.6) applies. To show that $\tilde{\alpha}$ solves our problem, it remains to show that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies the spin condition for all $n \leq k < j$. But for such k we have $\nu_k^w(n) = 1$, $\langle \tilde{\alpha}, a_k \rangle = -\frac{1}{2}$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{n,j}(\nu_k^w - \frac{1}{2}\alpha_{n,j}^\vee) \notin \mathbb{Z}^{2m}$. So $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies (SP3'). With this we have solved our problem in all subcases for which $c_i^w \leq c_p^w$.

Now suppose $c_i^w > c_p^w$. Then by (8.12.5)(i) $i < p < n$ and by (8.4.4)(ii) there exists a minimal element $j \in \{i+1, \dots, p\}$ such that $\mu_i(j) = 0$ and $\mu_p(j) = 1$. Note that evidently $\nu_k^w(j) = 0$ and $\nu_k^w(j^*) = 2$ for all $0 \leq k \leq i$. Moreover, since $\langle \tilde{\alpha}', a_p \rangle$ is a half-integer, (8.11.1)(ii) gives

$$c_p^{w'} \leq c_p^w + 1 \leq c_i^w \leq q.$$

Hence ν_p^w satisfies (SP2'). So it must fail (SP3'). Since this requires $\nu_p^{w'} = s_{i,\sigma(i)}(\nu_p^w + \langle \tilde{\alpha}', a_p \rangle \alpha_{i,\sigma(i)}^\vee)$ to have only integer entries, we conclude that $\nu_p^w(k)$ is a half-integer for $k \in \{i, i^*, \sigma(i), \sigma(i)^*\}$ and an integer for all other k . In particular, since $\mu_p(j) = 1$, we must have $\nu_p^w(j) = \nu_p^w(j^*) = 1$. We shall now prescribe $\tilde{\alpha}$ separately in the subcases $u(i) = 2$ and $u(i) = 1$.

If $u(i) = 2$ then we take

$$\tilde{\alpha} := \tilde{\alpha}_{i,j^*;-1}.$$

For $0 \leq k < i$, we have $\langle \tilde{\alpha}, a_k \rangle = 1$, $\nu_k^w(i) = 1$, and $\nu_k^w(j^*) = 2$. Hence $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \alpha_{i,j^*}^\vee) = \nu_k^w$ is μ -spin-permissible. Next let $i \leq k < j$. Then $\langle \tilde{\alpha}, a_k \rangle = \frac{1}{2}$. Since $u(i) = 2$ we have $\nu_i(i) = \frac{3}{2}$. And since $\nu_p^w(i)$ is a half-integer we must also have $\nu_p^w(i) = \frac{3}{2}$. Hence by (8.7.2) $\nu_k(i) = \frac{3}{2}$. Since $\nu_i^w(j^*) = 2$, (8.7.2) also implies $\nu_k^w(j^*) \in \{\frac{3}{2}, 2\}$. Hence by inspection $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \frac{1}{2}\alpha_{i,j^*}^\vee)$ satisfies (SP1'). And $c_k^{s_{\tilde{\alpha}}w} = c_k^w$, so that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). Moreover $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies (SP3') since $\nu_k^{s_{\tilde{\alpha}}w}(\sigma(i)) = \nu_k^w(\sigma(i))$ is a half-integer. Finally let $j \leq k \leq m$. Then $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}\nu_k^w$, which is clearly μ -spin-permissible, using as always (8.11.4) to address (SP3'). Since $\nu_p^w(i) = \frac{3}{2}$ and $\nu_p^w(j^*) = 1$, we see moreover from criterion (2) in (8.9.1) that $w < s_{\tilde{\alpha}}w$. Thus $\tilde{\alpha}$ solves our problem when $u(i) = 2$ and $c_i^w > c_p^w$.

If $u(i) = 1$ then we take

$$\tilde{\alpha} := \tilde{\alpha}_{i,j;0}.$$

Our argument will parallel the case $u(i) = 2$ just discussed. For $0 \leq k < i$ or $j \leq k \leq m$, we have $\langle \tilde{\alpha}, a_k \rangle = 0$. Hence $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j}\nu_k^w$, which is clearly μ -spin-permissible. Now let $i \leq k < j$. Then $\langle \tilde{\alpha}, a_k \rangle = -\frac{1}{2}$. Since $u(i) = 1$ we have $\nu_i(i) = \frac{1}{2}$. And since $\nu_p^w(i)$ is a half-integer we must also have $\nu_p^w(i) = \frac{1}{2}$. Hence by (8.7.2) $\nu_k(i) = \frac{1}{2}$. Since $\nu_i^w(j) = 0$, (8.7.2) also implies $\nu_k^w(j) \in \{0, \frac{1}{2}\}$. Hence by inspection $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j}(\nu_k^w - \frac{1}{2}\alpha_{i,j}^\vee)$ satisfies (SP1'). And $c_k^{s_{\tilde{\alpha}}w} = c_k^w$, so that $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). Moreover $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies (SP3') since $\nu_k^{s_{\tilde{\alpha}}w}(\sigma(i)) = \nu_k^w(\sigma(i))$ is a half-integer. Since $\nu_p^w(i) = \frac{1}{2}$ and $\nu_p^w(j) = 1$, we see moreover from criterion (2) in (8.9.1) that $w < s_{\tilde{\alpha}}w$. Thus $\tilde{\alpha}$ solves our problem when $u(i) = 1$ and $c_i^w > c_p^w$.

This finishes Case II, and with it the proof of part (i) of the Proposition is complete.

It remains to prove part (ii). So suppose $\nu_0^w(m) = 1$. Since $\nu_0^w \equiv \nu_m^w \bmod Q_{D_m}^\vee$ for any $w \in \widetilde{W}_{D_m}$, we must have $i < m$. If $\sigma(i) = i^*$, then a quick review of Case I above reveals that the affine root $\tilde{\alpha}$ we prescribed there satisfies $\nu_0^{s_{\tilde{\alpha}}w} = \nu_0^w$. So $\tilde{\alpha}$ certainly solves our problem in part (ii).

In the rest of the proof we shall assume that $\sigma(i) \neq i^*$. This assumption places us in the setting of Case II in the proof of part (i), where we used (8.11.2) to produce an affine root $\tilde{\alpha}' = \tilde{\alpha}_{i,\sigma(i);d}$ such that $w' = s_{\tilde{\alpha}'}w$ is naively μ -permissible and $w < w'$. We shall now consider cases based on the possibilities for w' .

Case A: w' is not μ -spin-permissible. Then in every subcase considered in Case II above, we prescribed another affine root $\tilde{\alpha}$ that solved the problem posed in part (i). We claim that $\tilde{\alpha}$ also has the additional property required in part (ii). As in Case II, let p be minimal in $\{0, \dots, m\}$ such that $\nu_p^{w'}$ is not μ -spin-permissible. A quick review of Case II reveals that $\tilde{\alpha}$ is of the form $\tilde{\alpha}_{i,j;0}$ or $\tilde{\alpha}_{i,j^*;-1}$ for some $i < j \leq p$, or $\tilde{\alpha}_{n,j;0}$ for some $n < j \leq p$. Thus to prove the claim it suffices to show that $j \neq m$, since then $\nu_k^{s_{\tilde{\alpha}}w}(m) = \nu_k^w(m)$ for all k .

To show that $j \neq m$, first note that if $p < m$, then of course $j < m$. So we might as well assume $p = m$. Then $c_p^w \equiv q \bmod 2$ and $\langle \tilde{\alpha}', a_p \rangle \in \mathbb{Z}$. Hence by (8.11.4) $\nu_p^{w'}$ satisfies the spin condition. So $\nu_p^{w'}$ must fail (SP2'). Hence

$$q < c_p^{w'} \leq c_p^w + 2 \leq q + 2,$$

where the middle inequality uses (8.11.1). Since $c_p^w \equiv q \bmod 2$, we conclude that $c_p^w = q$. On the other hand, (8.12.3) implies that $u(i) \neq u(\sigma(i))$, so that $c_i^w < q$ by (8.12.2). We conclude that $c_i^w < c_p^w$.

To complete the argument, we compare the entries of ν_i^w and ν_p^w . Since $\nu_p^w = \nu_m^w \in \mathbb{Z}^{2m}$, (8.11.5) implies that $\mu_p^w(i) = \mu_p^w(i^*) = 1$. On the other hand, we have $\mu_i^w(i) = 2$ or $\mu_i^w(i^*) = 2$ according as $u(i) = 2$ or $u(i) = 1$. Either way, since $c_i^w < c_p^w$, we see that there exist at least *two* elements $j' \in \{i+1, \dots, p\}$ such that $\mu_i^w(j') = 1$ and $\mu_p^w(j') = 2$. Such elements are clearly elements of the set S defined in (8.12.6). But when $c_i^w \leq c_p^w$, we defined j to be the *minimal* element of S . Therefore $j < m$, which completes Case A.

Case B: w' is μ -spin-permissible and $\sigma(i) \notin \{m, m+1\}$. Then $\nu_k^{s_{\tilde{\alpha}}w}(m) = \nu_k^w(m)$ for all k , so we may take $\tilde{\alpha} := \tilde{\alpha}'$.

Case C: w' is μ -spin-permissible and $\sigma(i) \in \{m, m+1\}$. In this case the m th entries of ν_0^w and $\nu_0^{w'}$ may differ, so the simple argument of Case B need not apply. Evidently $\mu_0^w(\sigma(i)) = 1$ and $\mu_i^w(\sigma(i)) = 0$, so that $u(\sigma(i)) = 1$; and $n = m$. We consider the subcases $u(i) = 1$ and $u(i) = 2$ separately.

First suppose $u(i) = 1$. Then $\nu_0^w(i) = 0$ and $\tilde{\alpha}'$ equals $\tilde{\alpha}_{i,\sigma(i);-1}$ or $\tilde{\alpha}_{i,\sigma(i);0}$. If $\tilde{\alpha}' = \tilde{\alpha}_{i,\sigma(i);-1}$, then $\langle \tilde{\alpha}', a_0 \rangle = 1$ and $\nu_0^{w'} = s_{i,\sigma(i)}(\nu_0^w + \alpha_{i,\sigma(i)}^\vee) = \nu_0^w$. Hence $\tilde{\alpha}'$ solves our problem. If $\tilde{\alpha}' = \tilde{\alpha}_{i,\sigma(i);0}$, then $\langle \tilde{\alpha}', a_0 \rangle = 0$ and $\nu_0^{w'}(\sigma(i)) = \nu_0^w(i) = 0$. So we must find a different affine root $\tilde{\alpha}$ to solve our problem.

Now, note that (8.11.2) allows us to take $\tilde{\alpha}' = \tilde{\alpha}_{i,\sigma(i);0}$ exactly when $K_i \subset [\sigma(i), i]$. If this inclusion is an equality $K_i = [\sigma(i), i]$, then (8.11.2) also allows us to take $\tilde{\alpha}_{i,\sigma(i);-1}$ for our $\tilde{\alpha}'$. And if $s_{i,\sigma(i);-1}w$ is μ -spin-permissible, then $\tilde{\alpha}_{i,\sigma(i);-1}$

solves our problem exactly as in the previous paragraph. If $s_{i,\sigma(i);-1}w$ is not μ -spin-permissible, then we can run through the argument of Case II with $\tilde{\alpha}_{i,\sigma(i);-1}$ in place of $\tilde{\alpha}_{i,\sigma(i);0}$ to produce a new affine root $\tilde{\alpha}$ that solves the problem posed in part (i). Case A then shows that this $\tilde{\alpha}$ also solves our problem in part (ii).

Still assuming $u(i) = 1$, we therefore reduce to solving our problem under the assumption $K_i \not\subset [\sigma(i), i)$. Since $\sigma(i) \in \{m, m+1\}$, this assumption implies that $m \notin K_i$. Hence $\mu_m^w(i) = 1$, which, since $\nu_m^w \in \mathbb{Z}^{2m}$, in turn implies that $\nu_m^w(i) = 1$. Let

$$\tilde{\beta} := \tilde{\alpha}_{i,\sigma(i);-1}.$$

Let us investigate the extent to which $\tilde{\beta}$ solves our problem. Let $0 \leq k < i$. Then as before $\nu_k^w(i) = 0$, $\nu_k^w(\sigma(i)) = 1$, $\langle \tilde{\beta}, a_k \rangle = 1$, and $\nu_k^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)}(\nu_k^w + \alpha_{i,\sigma(i)}^\vee) = \nu_k^w$, which is μ -spin-permissible. Next let $i \leq k < m$. Then $\nu_k^w(\sigma(i)) = \frac{1}{2}$ and $\langle \tilde{\beta}, a_k \rangle = \frac{1}{2}$. Since $\nu_i^w(i) = \frac{1}{2}$ and $\nu_m^w(i) = 1$, (8.7.2) implies that $\nu_k^w(i) \in \{\frac{1}{2}, 1\}$. Hence by inspection $\nu_k^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)}(\nu_k^w + \frac{1}{2}\alpha_{i,\sigma(i)}^\vee)$ satisfies (SP1'), and $c_k^{s_{\tilde{\beta}}w}$ equals c_k^w or $c_k^w + 1$ according as $\nu_k^w(i)$ equals $\frac{1}{2}$ or 1. Moreover, taking $k = i$ in criterion (1) in (8.9.1), we see that $w < s_{\tilde{\beta}}w$. Finally let $k = m$. If $m = \sigma(i)$, then $\nu_m^w(\sigma(i)) = 1$ and $\langle \tilde{\beta}, a_m \rangle = 1$. Since $\nu_m^w(i) = 1$, we see explicitly that $\nu_m^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)}(\nu_m^w + \tilde{\alpha}_{i,\sigma(i)}^\vee)$ satisfies (SP1') and $c_m^{s_{\tilde{\beta}}w} = c_m^w + 2$. On the other hand, if $m = \sigma(i)^*$, then $\langle \tilde{\beta}, a_m \rangle = 0$ and $\nu_m^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)}\nu_m^w$ is plainly μ -spin-permissible.

We conclude that $s_{\tilde{\beta}}w$ is naively μ -permissible and $w < s_{\tilde{\beta}}w$. If moreover $s_{\tilde{\beta}}w$ is μ -spin-permissible, then as before $\tilde{\beta}$ solves our problem. Otherwise, quite similarly to Case II above, we shall study the way in which $s_{\tilde{\beta}}w$ fails to be μ -spin-permissible to produce the desired affine root $\tilde{\alpha}$. Let $p \in \{0, \dots, m\}$ be minimal such that $\nu_p^{s_{\tilde{\beta}}w}$ is not μ -spin-permissible. It is an immediate consequence of our analysis in the previous paragraph that $i < p$.

The following will be the main tool to solve our problem in present situation.

Lemma 8.12.7. *Let $j \in \{i+1, \dots, p\}$ be such that $\mu_i^w(j) = 1$ and $\mu_p^w(j) = 2$, and let $\tilde{\alpha} := \tilde{\alpha}_{i,j^*;-1}$. Then $s_{\tilde{\alpha}}w$ is μ -spin-permissible and $\nu_0^{s_{\tilde{\alpha}}w}(m) = 1$.*

Proof. First note that $j < m$: this is clear if $p < m$, and if $p = m$ then our above analysis shows that $m = \sigma(i)$ and $\mu_p^w(m) = \mu_p^w(\sigma(i)) = 1 \neq \mu_p^w(j)$. Also note that evidently $\nu_k^w(j) = \nu_k^w(j^*) = 1$ for all $0 \leq k \leq i$, and $u(j^*) = 1$.

To prove the lemma, first let $0 \leq k < i$. Then $\nu_k^w(i) = 0$, $\nu_k^w(j^*) = 1$, $\langle \tilde{\alpha}, a_k \rangle = 1$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \alpha_{i,j^*}^\vee) = \nu_k^w$. Hence $\nu_k^{s_{\tilde{\alpha}}w}$ is μ -spin-permissible with m th entry equal to 1.

Next let $i \leq k < j$. Then $\nu_k^w(i) \in \{\frac{1}{2}, 1\}$, $\langle \tilde{\alpha}, a_k \rangle = \frac{1}{2}$, and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}(\nu_k^w + \frac{1}{2}\alpha_{i,j^*}^\vee)$. Since $\nu_i^w(j^*) = 1$ and $u(j^*) = 1$, we conclude from (8.7.2) that $\nu_k^w(j^*) \in \{\frac{1}{2}, 1\}$. Hence by inspection $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP1'), and $c_k^{s_{\tilde{\alpha}}w}$ equals c_k^w or $c_k^w + 1$ according as $\nu_k^w(i)$ equals $\frac{1}{2}$ or 1. If $\nu_k^w(i) = \frac{1}{2}$ then certainly $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). If $\nu_k^w(i) = 1$, then recall from above that $c_k^{s_{\tilde{\beta}}w} = c_k^w + 1$. Since $k < p$, minimality of p ensures that this is $\leq q$, and again $\nu_k^{s_{\tilde{\alpha}}w}$ satisfies (SP2'). Furthermore, since $j < m$ we have $\nu_k^{s_{\tilde{\alpha}}w}(\sigma(i)) = \nu_k^w(\sigma(i)) = \frac{1}{2}$, so that $\nu_k^{s_{\tilde{\alpha}}w}$ trivially satisfies (SP3').

Finally let $j \leq k \leq m$. Then $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\alpha}}w} = s_{i,j^*}\nu_k^w$ is clearly μ -spin-permissible. $\square$

To enable ourselves to apply the lemma, let us first show that $c_i^w \leq c_p^w$. For let $k > i$ be such that $c_i^w > c_k^w$. If $i < k < m$, then we showed above that $c_k^{s_{\tilde{\beta}}w} \leq c_k^w + 1$. By hypothesis this is $\leq c_i^w$, whence $\nu_k^{s_{\tilde{\beta}}w}$ satisfies (SP2'). To see that $\nu_k^{s_{\tilde{\beta}}w}$ also satisfies (SP3'), suppose that $\nu_k^{s_{\tilde{\beta}}w} \in \mathbb{Z}^{2m}$. Then $\nu_k^w(i) = \nu_k^w(\sigma(i)) = \frac{1}{2}$, $\nu_k^w(i^*) = \nu_k^w(\sigma(i)^*) = \frac{3}{2}$, and all other corresponding entries of ν_k^w and $\nu_k^{s_{\tilde{\beta}}w}$ are equal and in $\mathbb{Z}$. Now, by (8.4.4)(ii) there exists $j \in \{i+1, \dots, k\}$ such that $\mu_i^w(j) = 0$ and $\mu_k^w(j) = 1$. Since $\nu_k^w(j)$ is evidently an integer, it then also equals 1. On the other hand, the formula $\nu_k^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)}(\nu_k^w + \frac{1}{2}\alpha_{i,\sigma(i)}^\vee)$ gives explicitly that $\nu_k^{s_{\tilde{\beta}}w}(\sigma(i)) = 1$. Since $j \in A_k$ and $\sigma(i) \in B_k$, we conclude that $\nu_k^{s_{\tilde{\beta}}w}$ satisfies (SP3'). Thus $\nu_k^{s_{\tilde{\beta}}w}$ is μ -spin-permissible and $p \neq k$. Now suppose $k = m$. Then $c_k^w \equiv q \equiv c_i^w \pmod{2}$ by (8.12.2). Since $c_i^w > c_k^w$, we must then have $c_i^w \geq c_k^w + 2$. Hence by (8.11.1)(i) $c_k^{s_{\tilde{\beta}}w} \leq c_i^w$ and $\nu_k^{s_{\tilde{\beta}}w}$ satisfies (SP2'). As always $\nu_k^{s_{\tilde{\beta}}w}$ satisfies (SP3') by (8.11.4), and we again conclude that $p \neq k$. Thus $c_i^w \leq c_p^w$.

To complete the solution to our problem under our present assumptions, it remains to produce a j as in the lemma and to show that, in the notation of the lemma, $w < s_{\tilde{\alpha}}w$. Of course the vector $\nu_p^{s_{\tilde{\beta}}w}$ fails (SP2') or (SP3'), and we shall treat these cases separately.

Suppose that $\nu_p^{s_{\tilde{\beta}}w}$ fails (SP2'). Then certainly $c_p^{s_{\tilde{\beta}}w} > c_p^w$. If $i < p < m$, then we saw earlier that this requires $\nu_p^w(i) = 1$. If $p = m$, then we have already observed that $\nu_p^w(i) = 1$. Either way, we see in particular that $\mu_p^w(i^*) = 1$. Since $\mu_i^w(i^*) = 2$ and $c_i^w \leq c_p^w$, there then exists an element j such that $\mu_i^w(j) = 1$ and $\mu_p^w(j) = 2$. As usual, the recursion relation (8.3.2) implies that any such j must be contained in $\{i+1, \dots, p\}$, as desired. To see that, in the notation of the lemma, $w < s_{\tilde{\alpha}}w$, note that $\nu_p^w(j^*) \in \{0, \frac{1}{2}\}$ since $\mu_p^w(j) = 2$. So, since $\langle \tilde{\alpha}, a_p \rangle = 0$ and $\nu_p^w(i) = 1$, we get the conclusion by taking $k = p$ in criterion (2) in (8.9.1).

Now suppose that $\nu_p^{s_{\tilde{\beta}}w}$ fails (SP3'). Then $i < p < m$, $\nu_p^{s_{\tilde{\beta}}w} \in \mathbb{Z}^{2m}$, and $c_p^{s_{\tilde{\beta}}w} \not\equiv q \pmod{2}$. Our earlier analysis then shows that $\nu_p^w(i) = \frac{1}{2}$ and $c_p^{s_{\tilde{\beta}}w} = c_p^w$. Since $c_i^w \equiv q \pmod{2}$ by (8.12.2) and $c_i^w \leq c_p^w$ by our claim, we deduce the strict inequality $c_i^w < c_p^w$. Then (8.4.4)(i) gives us our j . To see that, in the notation of the lemma, $w < s_{\tilde{\alpha}}w$, we again use that $\mu_p^w(j) = 2$. Since this time we know that $\nu_p^w(j^*)$ is an integer, we conclude $\nu_p^w(j^*) = 0$. Since $\langle \tilde{\alpha}, a_p \rangle = 0$ and $\nu_p^w(i) = \frac{1}{2}$, we again get the conclusion by taking $k = p$ in criterion (2) in (8.9.1). With this we have completely solved our problem when $u(i) = 1$.

Now suppose $u(i) = 2$. Then $\nu_0^w(i) = 1$ and $\tilde{\alpha}'$ equals $\tilde{\alpha}_{i,\sigma(i);-1}$ or $\tilde{\alpha}_{i,\sigma(i);1}$. Neither possibility for $\tilde{\alpha}'$ solves our problem, since $\nu_0^{s_{\tilde{\alpha}'w}}(\sigma(i))$ equals 0 or 2. We are therefore led to consider

$$\tilde{\beta} := \tilde{\alpha}_{i,\sigma(i)^*;0}.$$

Indeed, for all $0 \leq k < i$, and in particular for $k = 0$, we have $\langle \tilde{\beta}, a_k \rangle = 0$ and $\nu_k^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)^*}\nu_k^w = \nu_k^w$, which by hypothesis has m th entry equal to 1. So $\tilde{\beta}$ solves our problem provided $s_{\tilde{\beta}}w$ is μ -spin-permissible and $w < s_{\tilde{\beta}}w$. Let us investigate the extent to which this is the case. We have just shown that $\nu_k^{s_{\tilde{\beta}}w} = \nu_k^w$ is μ -spin-permissible for $0 \leq k < i$. For $i \leq k < m$, we have $\nu_k^w(i) \in \{1, \frac{3}{2}, 2\}$, $\nu_k^w(\sigma(i)^*) = \frac{3}{2}$, and $\langle \tilde{\beta}, a_k \rangle = -\frac{1}{2}$. Hence by inspection $\nu_k^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)^*}(\nu_k^w - \frac{1}{2}\alpha_{i,\sigma(i)^*}^\vee)$ satisfies (SP1'), and $c_k^{s_{\tilde{\beta}}w}$ equals c_k^w or $c_k^w + 1$ according as $c_k^w \in \{\frac{3}{2}, 2\}$ or $c_k^w = 1$. If

$\nu_k^w(i) \in \{\frac{3}{2}, 2\}$ then certainly $\nu_k^{s_{\tilde{\beta}}w}$ satisfies (SP2'). If $\nu_k^w(i) = 1$, then we use that $\nu_k^{w'} = s_{i,\sigma(i)}(\nu_k^w + \langle \tilde{\alpha}', a_k \rangle \alpha_{i,\sigma(i)}^\vee)$ is μ -spin-permissible. Indeed, condition (SP1') forces $\tilde{\alpha}' = \tilde{\alpha}_{i,\sigma(i);-1}$ and $\langle \tilde{\alpha}', a_k \rangle = \frac{1}{2}$. Since $\nu_k^w(\sigma(i)) = \frac{1}{2}$, we see explicitly that $c_k^{w'} = c_k^w + 1$, which is $\leq q$ by condition (SP2'). Hence $\nu_k^{s_{\tilde{\beta}}w}$ satisfies (SP2') as well. Furthermore, since $\nu_i^w(i) = \nu_i^w(\sigma(i)^*) = \frac{3}{2}$, we get $w < s_{\tilde{\beta}}w$ by taking $k = i$ in criterion (1) in (8.9.1). Finally let $k = m$. Then $\nu_m^w(i) \in \{1, 2\}$. If $m = \sigma(i)^*$, then $\langle \tilde{\beta}, a_m \rangle = 0$ and $\nu_m^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)^*} \nu_m^w$ is plainly μ -spin-permissible. On the other hand, if $m = \sigma(i)$, then $\nu_m^w(\sigma(i)) = \nu_m^w(\sigma(i)^*) = 1$ and $\langle \tilde{\beta}, a_m \rangle = -1$. Hence $\nu_m^{s_{\tilde{\beta}}w} = s_{i,\sigma(i)^*}(\nu_m^w - \alpha_{i,\sigma(i)^*}^\vee)$ satisfies (SP1'), and $c_m^{s_{\tilde{\beta}}w}$ equals c_m^w or $c_m^w + 2$ according as $\nu_m^w(i)$ equals 2 or 1. If $\nu_m^w(i) = 2$ then certainly $\nu_m^{s_{\tilde{\beta}}w}$ satisfies (SP2'). If $\nu_m^w(i) = 1$ then we once more use that $\nu_m^{w'}$ is μ -spin-permissible. Indeed, since $m = \sigma(i)$ we have $\langle \tilde{\alpha}', a_m \rangle = \pm 1$ and $\nu_m^{w'} = s_{i,\sigma(i)}(\nu_m^w \pm \alpha_{i,\sigma(i)}^\vee)$. Hence $c_m^{w'} = c_m^w + 1$, which is $\leq q$ by condition (SP2'). Hence $\nu_m^{s_{\tilde{\beta}}w}$ satisfies (SP2') as well.

We conclude from the previous paragraph that $s_{\tilde{\beta}}w$ fails to solve our problem if and only if $\nu_p^{s_{\tilde{\beta}}w}$ fails the spin condition for some p . In fact it suffices to consider just $p = i$: first let $i \leq p < m$ and suppose that $\nu_p^{s_{\tilde{\beta}}w} \in \mathbb{Z}^{2m}$. Then

$$\nu_p^w(i) = \nu_p^w(\sigma(i)^*) = \frac{3}{2} \quad \text{and} \quad \nu_p^w(\sigma(i)) = \nu_p^w(i^*) = \frac{1}{2};$$

all other entries of ν_p^w are integers;

$$\nu_p^{s_{\tilde{\beta}}w}(i) = 2, \quad \nu_p^{s_{\tilde{\beta}}w}(i^*) = 0, \quad \nu_p^{s_{\tilde{\beta}}w}(\sigma(i)) = 1, \quad \text{and} \quad \nu_p^{s_{\tilde{\beta}}w}(\sigma(i)^*) = 1;$$

and all other corresponding entries of $\nu_p^{s_{\tilde{\beta}}w}$ and ν_p^w are equal. Hence $c_p^{s_{\tilde{\beta}}w} = c_p^w$. If this is $\not\equiv q \pmod{2}$, then, since $\sigma(i) \in \{m, m+1\} \subset B_p$, we conclude that $\nu_p^{s_{\tilde{\beta}}w}$ satisfies the spin condition $\iff$ there exists $j \in A_p$ such that $\nu_p^{s_{\tilde{\beta}}w}(j) = 1$; or equivalently, since such a j is evidently distinct from i and i^* , there exists $j \in A_p$ such that $\nu_p^w(j) = 1$. In particular, our discussion applies when $p = i$, using (8.12.2) for the relation $c_i^w \not\equiv q \pmod{2}$. If such a j exists in this case, then $\nu_k^{s_{\tilde{\beta}}w}(j) = \nu_k^w(j) = \nu_i^w(j) = \nu_i^{s_{\tilde{\beta}}w} = 1$ for all k since i and i^* are the only proper elements in A_i . And then $\nu_p^{s_{\tilde{\beta}}w}$ satisfies the spin condition for all $i \leq p < m$ since $j \in A_p$ and $\sigma(i) \in B_p$. We conclude that $s_{\tilde{\beta}}w$ is μ -spin-permissible $\iff \nu_i^{s_{\tilde{\beta}}w}$ satisfies the spin condition.

So, to complete the proof, let us suppose that $\nu_i^{s_{\tilde{\beta}}w}$ fails the spin condition; or equivalently, that $\nu_i^w(k) \in \{0, 2\}$ for all $k \in A_i \setminus \{i, i^*\}$. We shall make crucial use of the fact that $q < m$. By explicit calculation, we have $\nu_i^{w'}(k) \in \{0, 2\}$ for all $k \in \{i, i^*, \sigma(i), \sigma(i^*)\}$. Hence $\nu_i^{w'}(k) \in \{0, 2\}$ for all $k \in A_i \cup \{m, m+1\}$. Since $q < m$, there must exist a j , which we may moreover take to be minimal, such that $i < j < m$ and $\nu_i^{w'}(j) = \nu_i^w(j) = 1$. Of course $\nu_k^w(j) = 1$ for all $0 \leq k \leq i$. Let

$$\tilde{\gamma} := \tilde{\alpha}_{i,j^*;-1}.$$

For $0 \leq k < i$, we have $\nu_k^w(i) = \nu_k^w(j) = \langle \tilde{\gamma}, a_k \rangle = 1$ and $\nu_k^{s_{\tilde{\gamma}}w} = s_{i,j^*}(\nu_k^w + \alpha_{i,j^*}^\vee)$. Hence $\nu_k^{s_{\tilde{\gamma}}w}$ satisfies (SP1') and $c_k^{s_{\tilde{\gamma}}w} = c_k^w + 2$. But

$$c_k^w + 2 = c_{i-1}^w + 2 = c_i^w + 1 \leq q,$$

where the second equality uses (8.4.3). So $\nu_k^{s\tilde{\gamma}w}$ satisfies (SP2'). And of course $\nu_k^{s\tilde{\gamma}w}$ satisfies the spin condition by (8.11.4). Thus $\nu_k^{s\tilde{\gamma}w}$ is μ -spin-permissible. Next let $j \leq k \leq m$. Then $\langle \tilde{\gamma}, a_k \rangle = 0$ and $\nu_k^{s\tilde{\gamma}w} = s_{i,j^*} \nu_k^w$ is clearly μ -spin-permissible. Finally let $i \leq k < j$. Then $\nu_k^w(i) \in \{1, \frac{3}{2}, 2\}$, $\nu_k^w(j^*) \in \{\frac{1}{2}, 1, \frac{3}{2}\}$ by (8.7.2), $\langle \tilde{\gamma}, a_k \rangle = \frac{1}{2}$, and $\nu_k^{s\tilde{\gamma}w} = s_{i,j^*}(\nu_k^w + \frac{1}{2}\alpha_{i,j^*}^\vee)$. Hence $\nu_k^{s\tilde{\gamma}w}$ satisfies (SP1') provided $\nu_k^w(i) \neq 2$. Let us suppose for the moment that this is the case. Since $\tilde{\alpha}' = \tilde{\alpha}_{i,\sigma(i);\pm 1}$ and $\nu_k^w(\sigma(i)) = \frac{1}{2}$, we see explicitly from the formula $\nu_k^{w'} = s_{i,\sigma(i)}(\nu_k^w + \langle \tilde{\alpha}', a_k \rangle \alpha_{i,\sigma(i)}^\vee)$ that $c_k^{w'} = c_k^w + 1$, which is $\leq q$ by condition (SP2'). Hence

$$c_k^{s\tilde{\gamma}w} \leq c_k^w + 1 \leq q,$$

where the first inequality uses (8.11.1)(ii). Hence $\nu_k^{s\tilde{\gamma}w}$ satisfies (SP2'). And $\nu_k^{s\tilde{\gamma}w}$ trivially satisfies (SP3') because $\nu_k^{s\tilde{\gamma}w}(\sigma(i)) = \nu_k^w(\sigma(i)) = \frac{1}{2}$ is a half-integer. Furthermore, since $\nu_i^w(i) = \frac{3}{2}$ and $\nu_i^w(j^*) = 1$, we see by taking $k = i$ in criterion (1) in (8.9.1) that $w < s\tilde{\gamma}w$.

In conclusion, we have shown that if $\nu_k^w(i) \neq 2$ for all $i \leq k < j$, then $s\tilde{\gamma}w$ is μ -spin-permissible and $w < s\tilde{\gamma}w$. Moreover $\tilde{\gamma}$ then completely solves our problem, since $\nu_k^{s\tilde{\gamma}w}(m) = \nu_k^w(m)$ for all k . So it remains to find a solution to our problem under the assumption that $\nu_k^w(i) = 2$ for some $i < k < j$, which occurs exactly when $i < \sigma^{-1}(i^*) < j$. Minimality of j forces $\nu_i^w(\sigma^{-1}(i^*))$ to equal 0 or 2, and we shall complete the proof by considering these cases separately.

If $\nu_i^w(\sigma^{-1}(i^*)) = 0$ then $\nu_i^w(\sigma^{-1}(i)) = 2$, and we take

$$\tilde{\alpha} := \tilde{\alpha}_{i,\sigma^{-1}(i);-1}.$$

To see that $\tilde{\alpha}$ solves our problem, first let $0 \leq k < i$. Then $\nu_k^w(i) = 1$, $\nu_k^w(\sigma^{-1}(i)) = 2$, $\langle \tilde{\alpha}, a_k \rangle = 1$, and $\nu_k^{s\tilde{\alpha}w} = s_{i,\sigma^{-1}(i)}(\nu_k^w + \alpha_{i,\sigma^{-1}(i)}^\vee) = \nu_k^w$. Hence $\nu_k^{s\tilde{\alpha}w}$ is μ -spin-permissible. Next let $i \leq k < \sigma^{-1}(i^*)$. Then $\nu_k^w(i) = \frac{3}{2}$, $\nu_k^w(\sigma^{-1}(i)) \in \{\frac{3}{2}, 2\}$ by (8.7.2), $\langle \tilde{\alpha}, a_k \rangle = \frac{1}{2}$, and $\nu_k^{s\tilde{\alpha}w} = s_{i,\sigma^{-1}(i)}(\nu_k^w + \frac{1}{2}\alpha_{i,\sigma^{-1}(i)}^\vee)$. Hence by inspection $\nu_k^{s\tilde{\alpha}w}$ satisfies (SP1') and $c_k^{s\tilde{\alpha}w} = c_k^w$, so that $\nu_k^{s\tilde{\alpha}w}$ also satisfies (SP2'). Moreover $\nu_k^{s\tilde{\alpha}w}$ trivially satisfies (SP3') since $\nu_k^{s\tilde{\alpha}w}(\sigma(i)) = \nu_k^w(\sigma(i)) = \frac{1}{2}$ is a half-integer. Finally let $\sigma^{-1}(i^*) \leq k \leq m$. Then $\langle \tilde{\alpha}, a_k \rangle = 0$ and $\nu_k^{s\tilde{\alpha}w} = s_{i,\sigma^{-1}(i)}\nu_k^w$ is clearly μ -spin-permissible. Moreover we have $\nu_{\sigma^{-1}(i^*)}^w(i) = 2$ and $\nu_{\sigma^{-1}(i^*)}^w(\sigma^{-1}(i^*)) = 1$, so that $\nu_{\sigma^{-1}(i^*)}^w(\sigma^{-1}(i)) \in \{1, \frac{3}{2}\}$. Hence $w < s\tilde{\alpha}w$ by taking $k = \sigma^{-1}(i^*)$ in criterion (2) in (8.9.1). Of course $\nu_k^{s\tilde{\alpha}w}(m) = \nu_k^w(m)$ for all k , and we conclude that $\tilde{\alpha}$ solves our problem.

If $\nu_i^w(\sigma^{-1}(i^*)) = 2$ then $\nu_i^w(\sigma^{-1}(i)) = 0$, and we take

$$\tilde{\alpha} := \tilde{\alpha}_{i,\sigma^{-1}(i);1}.$$

To see that $\tilde{\alpha}$ solves our problem, first let $0 \leq k < i$. Then $\nu_k^w(i) = 1$, $\nu_k^w(\sigma^{-1}(i)) = 0$, $\langle \tilde{\alpha}, a_k \rangle = -1$, and $\nu_k^{s\tilde{\alpha}w} = s_{i,\sigma^{-1}(i)}(\nu_k^w - \alpha_{i,\sigma^{-1}(i)}^\vee) = \nu_k^w$. Hence $\nu_k^{s\tilde{\alpha}w}$ is μ -spin-permissible. Next let $i \leq k < \sigma^{-1}(i^*)$. Then $\nu_k^w(i) = \frac{3}{2}$, $\nu_k^w(\sigma^{-1}(i)) \in \{0, \frac{1}{2}\}$ by (8.7.2), $\langle \tilde{\alpha}, a_k \rangle = -\frac{3}{2}$, and $\nu_k^{s\tilde{\alpha}w} = s_{i,\sigma^{-1}(i)}(\nu_k^w - \frac{3}{2}\alpha_{i,\sigma^{-1}(i)}^\vee)$. Hence by inspection $\nu_k^{s\tilde{\alpha}w}$ satisfies (SP1') and $c_k^{s\tilde{\alpha}w} \leq c_k^w + 1$. To see that $\nu_k^{s\tilde{\alpha}w}$ satisfies (SP2'), we use, as in earlier parts of the proof, that $\nu_k^{w'}$ does: indeed $q \geq c_k^{w'} = c_k^w + 1$ by (8.11.3) (applied with $\tilde{\alpha} = \tilde{\alpha}'$), which gives us what we need. Moreover $\nu_k^{s\tilde{\alpha}w}$ trivially satisfies (SP3') since $\nu_k^{s\tilde{\alpha}w}(\sigma(i)) = \nu_k^w(\sigma(i)) = \frac{1}{2}$ is a half-integer. Finally let $\sigma^{-1}(i^*) \leq k \leq m$. Then $\nu_k^w(i) = 2$, $\nu_k^w(\sigma^{-1}(i)) = 0$, $\langle \tilde{\alpha}, a_k \rangle = -2$, and

$\nu_k^{s_{\tilde{\alpha}}w} = s_{i, \sigma^{-1}(i)}(\nu_k^w - 2\alpha_{i, \sigma^{-1}(i)}^\vee) = \nu_k^w$. Hence $\nu_k^{s_{\tilde{\alpha}}}$ is μ -spin-permissible. To see that $w < s_{\tilde{\alpha}}w$, note that $\sigma^{-1}(i)$ is evidently proper. So there must exist a k such that $\nu_k^w(\sigma^{-1}(i)) \neq 0$. By what we have just seen, such a k must satisfy $i \leq k < \sigma^{-1}(i^*)$, $\nu_k^w(\sigma^{-1}(i)) = \frac{1}{2}$, and $\nu_k^w(i) = \frac{3}{2}$. Hence $w < s_{\tilde{\alpha}}w$ by criterion (1) in (8.9.1). Finally $\nu_k^{s_{\tilde{\alpha}}w}(m) = \nu_k^w(m)$ for all k , and we conclude that $\tilde{\alpha}$ solves our problem. This completes the proof in Case C when $u(i) = 2$, and with it the entire proposition is proved. $\square$

This finally completes the proof of (8.2.2), and, equivalently, of Theorem 1.5.

8.13. Parahoric variants and faces of type I. Our aims for the rest of §8 are to extend (8.2.2) and (8.6.4) to the general parahoric case (in §8.14 and §8.16, respectively) and to characterize μ -admissibility in terms of μ -vertexwise admissibility (in §8.15). In this subsection we shall lay some of the needed groundwork.

For F a subfacet of the base alcove A_{D_m} , let W_{F, D_m} denote the common stabilizer and pointwise fixer of F in W_{aff, D_m} , as in §5.1. To simplify matters for ourselves, we introduce the affine transformations on $\mathcal{A}_{D_m}$

$$(8.13.1) \quad \begin{aligned} \theta_0 &: (x_1, \dots, x_{2m}) \mapsto (x_{2m} - 1, x_2, \dots, x_{2m-1}, x_1 + 1), \\ \theta_m &: (x_1, \dots, x_{2m}) \mapsto (x_1, \dots, x_{m-1}, x_{m+1}, x_m, x_{m+2}, \dots, x_{2m}). \end{aligned}$$

Then θ_0 and θ_m induce automorphisms of the affine root system of $\mathcal{B}_{D_m}$, and θ_0 (resp. θ_m) interchanges the points a_0 and $a_{0'}$ (resp. a_m and $a_{m'}$) and fixes a_k for all other $k \in \{0, 0', 2, \dots, m-2, m, m'\}$. In particular, θ_0 and θ_m send A_{D_m} to itself. It follows that both θ_0 and θ_m induce automorphisms of $\widetilde{W}_{D_m}$ which preserve the A_{D_m} -Bruhat order. The force of this in the next two subsections will be that, by applying θ_0 or θ_m (or both) as needed, we shall incur no loss of generality by considering only subfacets F of A_{D_m} with the property that

$$(8.13.2) \quad a_{0'} \in \overline{F} \implies a_0 \in \overline{F} \quad \text{and} \quad a_{m'} \in \overline{F} \implies a_m \in \overline{F}.$$

Given a subfacet F of A_{D_m} satisfying (8.13.2), let

$$I := \{k \in \{0, \dots, m\} \mid a_k \in \overline{F}\}.$$

Then I is a nonempty subset of $\{0, \dots, m\}$ satisfying

$$(8.13.3) \quad 1 \in I \implies 0 \in I \quad \text{and} \quad m-1 \in I \implies m \in I.$$

Indeed, if $a_1 \in \overline{F}$ (resp. $a_{m-1} \in \overline{F}$), then so is a_0 (resp. a_m), simply because a_0 (resp. a_m) is in the closure of the facet containing a_1 (resp. a_{m-1}). Moreover I uniquely determines F , since $a_{0'}$ (resp. $a_{m'}$) is also in the closure of the facet containing a_1 (resp. a_{m-1}), and any subfacet of A_{D_m} is specified by which of the vertices $a_0, a_{0'}, a_2, \dots, a_{m-2}, a_m, a_{m'}$ are in its closure. In this way we obtain a bijection between the subfacets F of A_{D_m} satisfying (8.13.2) and the nonempty subsets I of $\{0, \dots, m\}$ satisfying (8.13.3). Given such I , let

$$(8.13.4) \quad W_{I, D_m} := \{w \in W_{\text{aff}, D_m} \mid wa_k = a_k \text{ for all } k \in I\}.$$

Of course

$$W_{I, D_m} = W_{F, D_m}$$

for the unique subfacet F of A_{D_m} determined by I in the way we have just described.

Our aim for the rest of the subsection is to interpret the set $\widetilde{W}_{D_m}/W_{I,D_m}$ in terms of faces of type I , for nonempty $I \subset \{0, \dots, m\}$ satisfying (8.13.3). Define a homomorphism

$$\begin{aligned} \varepsilon: \widetilde{W}_{D_m} &\longrightarrow \mathbb{Z}/2\mathbb{Z} \\ w &\longmapsto \sum_{i=1}^m \nu_0^w(i) \bmod 2. \end{aligned}$$

Thus the affine Weyl group $W_{\text{aff}, D_m} = Q_{D_m}^\vee \rtimes S_{2m}^\circ$ may be characterized as the subset of elements $w \in \widetilde{W}_{D_m}$ for which $\nu_0^w + (\nu_0^w)^* = \mathbf{0}$ and $\varepsilon(w) = 0$.

Definition 8.13.5. Let I be a nonempty subset of $\{0, \dots, m\}$ (not necessarily satisfying (8.13.3)). We define the subset $\mathcal{F}_{I,D_m}$ of $\{\text{faces of type } I\} \times \mathbb{Z}/2\mathbb{Z}$ as follows. If $0, m \in I$, then we define

$$\mathcal{F}_{I,D_m} := \{(\mathbf{v}, \gamma) \mid \mu_0^\mathbf{v} \equiv \mu_m^\mathbf{v} \bmod Q_{D_m}^\vee \text{ and } \gamma = \varepsilon(t_{\mu_0^\mathbf{v}})\}.$$

If $0 \in I$ and $m \notin I$ (resp. $m \in I$ and $0 \notin I$), then we define

$$\mathcal{F}_{I,D_m} := \{(\mathbf{v}, \gamma) \mid \gamma = \varepsilon(t_{\mu_0^\mathbf{v}})\} \quad (\text{resp. } \mathcal{F}_{I,D_m} := \{(\mathbf{v}, \gamma) \mid \gamma = \varepsilon(t_{\mu_m^\mathbf{v}})\}).$$

If $0, m \notin I$, then we define

$$\mathcal{F}_{I,D_m} := \{\text{faces of type } I\} \times \mathbb{Z}/2\mathbb{Z}.$$

If $0, m \in I$ and $(\mathbf{v}, \gamma) \in \mathcal{F}_{I,D_m}$, then of course we also have $\gamma = \varepsilon(t_{\mu_m^\mathbf{v}})$. The group $\widetilde{W}_{D_m}$ acts naturally on $\mathcal{F}_{I,D_m}$ via the rule $w \cdot (\mathbf{v}, \gamma) = (w \cdot \mathbf{v}, \varepsilon(w) + \gamma)$. For a nonempty subset J of I , the natural $\widetilde{W}_{D_m}$ -equivariant map

$$\{\text{faces of type } I\} \longrightarrow \{\text{faces of type } J\}$$

induces a natural $\widetilde{W}_{D_m}$ -equivariant map

$$\mathcal{F}_{I,D_m} \longrightarrow \mathcal{F}_{J,D_m}.$$

Furthermore, if exactly one of $0, m$ is in I (resp. both 0 and m are in I), then the $\widetilde{W}_{D_m}$ -equivariant projection

$$\mathcal{F}_{I,D_m} \longrightarrow \{\text{faces of type } I\}$$

is a bijection (resp. is an injection whose image consists of all faces $\mathbf{v}$ such that $\mu_0^\mathbf{v} \equiv \mu_m^\mathbf{v} \bmod Q_{D_m}^\vee$).

Now suppose that I satisfies (8.13.3), let $w \in W_{I,D_m}$, and consider the standard face ω_I of type $(2m, I)$. The affine Weyl group W_{aff, D_m} is naturally a subgroup of the affine Weyl group

$$\{v \in \mathbb{Z}^{2m} \mid \Sigma v = 0\} \rtimes S_{2m}$$

for the root datum attached to GL_{2m} , and the points $\omega_k \in \mathbb{R}^{2m}$ for $k \in \mathbb{Z}$ are all contained in minimal facets of an alcove for GL_{2m} . For $k \in I$, since w fixes a_k and a_k is the midpoint of ω_k and ω_{-k} , it follows that w fixes ω_k and ω_{-k} . Hence w fixes ω_I . Since moreover $\varepsilon(W_{I,D_m}) \subset \varepsilon(W_{\text{aff}, D_m}) = 0$, we get a well-defined $\widetilde{W}_{D_m}$ -equivariant map

$$\begin{aligned} f: \widetilde{W}_{D_m}/W_{I,D_m} &\longrightarrow \mathcal{F}_{I,D_m} \\ wW_{I,D_m} &\longmapsto w \cdot (\omega_I, 0). \end{aligned}$$

Our main result for the subsection is the following.

Lemma 8.13.6. *Let I be a nonempty subset of $\{0, \dots, m\}$ satisfying (8.13.3). Then the map $f: \widetilde{W}_{D_m}/W_{I,D_m} \rightarrow \mathcal{F}_{I,D_m}$ is a bijection.*

Proof. First suppose that $w \in \widetilde{W}_{D_m}$ fixes $(\omega_I, 0)$. Then $\varepsilon(w) = 0$; since w fixes some vector in $\mathbb{R}^{2m}$, w must satisfy $\nu_0^w + (\nu_0^w)^* = \mathbf{0}$; and, since w fixes ω_k and ω_{-k} for all $k \in I$, w fixes their midpoint a_k for all $k \in I$. Hence $w \in W_{I,D_m}$. This proves that the map f is injective.

To see that f is surjective when 0 or m is in I , we need simply note that $\widetilde{W}_{D_m}$ acts transitively on the image of $\mathcal{F}_{I,D_m}$ under the injection

$$\mathcal{F}_{I,D_m} \hookrightarrow \{\text{faces of type } I\}.$$

If $0, m \notin I$, then we note that the element

$$w := \theta_0 \theta_m = t_{(-1, 0(2m-2), 1)}(1, 2m)(m, m+1) \in \widetilde{W}_{D_m},$$

where we use cycle notation for elements in S_{2m}° , fixes ω_I and satisfies $\varepsilon(w) = 1$. So the surjectivity of f for such I now follows from the fact that $\widetilde{W}_{D_m}$ acts transitively on faces of type I . $\square$

8.14. μ -admissibility and μ -spin-permissibility in the parahoric case. In this subsection we generalize (8.2.2) to the general parahoric case. For subfacets F, F' of A_{D_m} , we write

$$\text{Adm}_{F', F, D_m}(\mu)$$

for the μ -admissible set in $W_{F', D_m} \setminus \widetilde{W}_{D_m}/W_{F, D_m}$ (5.3.1). When $F = F' = A_{D_m}$, we abbreviate this to $\text{Adm}_{D_m}(\mu)$.

Recall $\theta_0, \theta_m \in \text{Aff}(\mathcal{A}_{D_m})$ from (8.13.1), and let $\theta \in \{\theta_0, \theta_m, \theta_0 \theta_m\}$. Then $\theta \in \widetilde{W}_{2m}$ (4.1.2), and θ stabilizes the base alcove A_{D_m} . Hence for any $\mu \in X_{*D_m}$, conjugation by θ induces an automorphism

$$W_{F', D_m} \setminus \widetilde{W}_{D_m}/W_{F, D_m} \xrightarrow{\sim} W_{\theta F', D_m} \setminus \widetilde{W}_{D_m}/W_{\theta F, D_m}$$

identifying

$$\text{Adm}_{F', F, D_m}(\mu) \xrightarrow{\sim} \text{Adm}_{\theta F', \theta F, D_m}(\sigma \cdot \mu),$$

where σ denotes the linear part of θ . When μ is the cocharacter (8.0.1), since $q < m$ we have $S_{2m}^\circ \mu = S_{2m}^* \mu$, and therefore the admissible set on the right-hand side of the last display equals $\text{Adm}_{\theta F', \theta F, D_m}(\mu)$.

In the case that $F \preceq F' \preceq A_{D_m}$, we conclude that in studying admissible sets for this μ , it is harmless to assume that F and F' satisfy property (8.13.2), since we can always apply a suitable $\theta \in \{\theta_0, \theta_m, \theta_0 \theta_m\}$ to place ourselves in this situation. Thus we shall assume that W_{F, D_m} (resp. W_{F', D_m}) is of the form W_{I, D_m} (resp. W_{J, D_m}) for some nonempty $I \subset J \subset \{0, \dots, m\}$ satisfying property (8.13.3).

Now recall that for $w \in \widetilde{W}_{D_m}/W_{I, D_m}$, we get well-defined vectors $\mu_k^w = w\omega_k - \omega_k$ and $\nu_k^w = wa_k - a_k$ for $k \in 2m \pm I$, as discussed in the previous subsection.

Definition 8.14.1. Let μ be the cocharacter (8.0.1), and let $I \subset J$ be nonempty subsets of $\{0, \dots, m\}$ satisfying property (8.13.3). We say that $w \in W_{J, D_m} \setminus \widetilde{W}_{D_m}/W_{I, D_m}$ is μ -spin-permissible if $w \equiv t_\mu \pmod{W_{\text{aff}, D_m}}$ and for one, hence any, representative $\tilde{w}$ of w in $\widetilde{W}_{D_m}/W_{I, D_m}$, $\mu_k^{\tilde{w}}$ satisfies conditions (SP1), (SP2), and (SP3) in (8.2.1) for all $k \in I$. We write

$$\text{SPerm}_{J, I, D_m}(\mu)$$

for the set of μ -spin-permissible elements in $W_{J,D_m} \backslash \widetilde{W}_{D_m} / W_{I,D_m}$. We abbreviate this to $\text{SPerm}_{I,D_m}(\mu)$ when $J = \{0, \dots, m\}$, and to $\text{SPerm}_{D_m}(\mu)$ when $I = J = \{0, \dots, m\}$.

We shall see in (8.14.4) below (with an assist from (8.5.7)) that the definition really is independent of the choice of representative $\tilde{w} \in \widetilde{W}_{D_m} / W_{I,D_m}$. Let us first make some other remarks.

First, the condition $w \equiv t_\mu \bmod W_{\text{aff},D_m}$ is well-defined because $W_{I,D_m}, W_{J,D_m} \subset W_{\text{aff},D_m}$. If 0 (resp. m) is in I , then this condition is moreover redundant, since (SP3) requires $w \equiv t_{\mu_0^w} \equiv t_\mu \bmod W_{\text{aff},D_m}$ (resp. $w \equiv t_{\mu_m^w} \equiv t_\mu \bmod W_{\text{aff},D_m}$). Of course, for any I satisfying (8.13.3), w is equivalently μ -spin-permissible if $w \equiv t_\mu \bmod W_{\text{aff},D_m}$ and $\nu_k^{\tilde{w}}$ satisfies (SP1'), (SP2'), and (SP3') for all $k \in I$, for any representative $\tilde{w} \in \widetilde{W}_{D_m} / W_{I,D_m}$, by (8.5.7).

To prepare for (8.14.4), we first prove the following.

Lemma 8.14.2. *Let $x \in \widetilde{W}_{D_m}$, and write $x = t_{\nu_0^x} \tau$, with $\nu_0^x \in X_{*D_m}$ and τ in the Weyl group S_{2m}° . Suppose that $xa_k = a_k$ for some $k \in \{1, \dots, m-1\}$. Then the sets A_k and B_k defined in (4.3.6) are τ -stable.*

Proof. Suppose by contradiction that τ carries some element $i \in A_k$ into B_k . Since $a_k = ((-\frac{1}{2})^{(k)}, 0^{(2m-2k)}, (\frac{1}{2})^{(k)})$, we obtain

$$(\tau a_k - a_k)(\tau(i)) = (\tau a_k)(\tau(i)) = a_k(i) = \pm \frac{1}{2}.$$

But

$$(8.14.3) \quad a_k = xa_k = \nu_0^x + \tau a_k.$$

Hence $\tau a_k - a_k = -\nu_0^x \in X_{*D_m} \subset \mathbb{Z}^{2m}$, a contradiction. $\square$

Lemma 8.14.4. *Let $w, w' \in \widetilde{W}_{D_m} / W_{I,D_m}$, and suppose that $w' = xw$ for some $x \in W_{I,D_m}$. Then for $k \in I$, ν_k^w satisfies (SP1') (resp. (SP2'), resp. (SP3')) $\iff \nu_k^{w'}$ satisfies (SP1') (resp. (SP2'), resp. (SP3')). In particular, w is μ -spin-permissible $\iff w'$ is μ -spin-permissible.*

Proof. Let $k \in I$, and write $x = t_{\nu_0^x} \tau$ with τ in the Weyl group S_{2m}° . Then, using (8.14.3) to substitute for a_k in the middle equality,

$$\nu_k^{w'} = xwa_k - a_k = \tau wa_k - \tau a_k = \tau \nu_k^w.$$

So it is immediate that $\nu_k^{w'}$ satisfies (SP1') (resp. (SP2')) $\iff \nu_k^w$ satisfies (SP1') (resp. (SP2')). To consider the spin condition, we see from the display that $\nu_k^{w'} \in \mathbb{Z}^{2m} \iff \nu_k^w \in \mathbb{Z}^{2m}$. Let us suppose this is so. The display implies that $\nu_k^{w'} \equiv \nu_k^w \bmod Q_{D_m}^\vee$. If these are both $\equiv \mu$, then they trivially satisfy the spin condition. So suppose they are $\not\equiv \mu$. Since $\nu_k^{w'} = \tau \nu_k^w$ and τ permutes the sets A_k and B_k separately by (8.14.2), we conclude at once that $\nu_k^{w'}$ satisfies the spin condition $\iff \nu_k^w$ does, which completes the proof. $\square$

Our main result for the subsection is the following. We write $\text{Adm}_{J,I,D_m}(\mu)$ for the set of μ -admissible elements in $W_{J,D_m} \backslash \widetilde{W}_{D_m} / W_{I,D_m}$. We abbreviate this to $\text{Adm}_{I,D_m}(\mu)$ when $J = \{0, \dots, m\}$, and to $\text{Adm}_{D_m}(\mu)$ when $I = J = \{0, \dots, m\}$.

Theorem 8.14.5. *Let μ be the cocharacter (8.0.1), and let $I \subset J \subset \{0, \dots, m\}$ be nonempty sets satisfying (8.13.3). Then*

$$\text{Adm}_{J,I,D_m}(\mu) = \text{SPerm}_{J,I,D_m}(\mu).$$

Proof. The natural map $\widetilde{W}_{D_m}/W_{I,D_m} \rightarrow W_{J,D_m} \setminus \widetilde{W}_{D_m}/W_{I,D_m}$ induces a surjection

$$\text{Adm}_{I,D_m}(\mu) \twoheadrightarrow \text{Adm}_{J,I,D_m}(\mu)$$

by definition, and a surjection

$$\text{SPerm}_{I,D_m}(\mu) \twoheadrightarrow \text{SPerm}_{J,I,D_m}(\mu)$$

by (8.14.4). Thus it suffices to establish the equality $\text{Adm}_{I,D_m}(\mu) = \text{SPerm}_{I,D_m}(\mu)$ in $\widetilde{W}_{D_m}/W_{I,D_m}$. When $I = \{0, \dots, m\}$, the result is given by (8.2.2). For general I , we consider the natural map $\pi: \widetilde{W}_{D_m} \rightarrow \widetilde{W}_{D_m}/W_{I,D_m}$. Again by definition, π induces a surjection $\text{Adm}_{D_m}(\mu) \twoheadrightarrow \text{Adm}_{I,D_m}(\mu)$. So the result follows from the fact that π also induces a surjection $\text{SPerm}_{D_m}(\mu) \twoheadrightarrow \text{SPerm}_{I,D_m}(\mu)$, which we prove in (8.14.6) below. $\square$

In fact we shall formulate (8.14.6) in terms of faces of type I , or more precisely, in terms of the sets $\mathcal{F}_{I,D_m}$ (8.13.5) for varying I . Recall that in the previous subsection, for nonempty $I \subset \{0, \dots, m\}$ satisfying (8.13.3), we established a $\widetilde{W}_{D_m}$ -equivariant bijection $\widetilde{W}_{D_m}/W_{I,D_m} \xrightarrow{\sim} \mathcal{F}_{I,D_m}$, compatible with the natural projections on both sides as we pass from I to a subset of I . To prove (8.14.6), it will be most natural to consider $\mathcal{F}_{I,D_m}$ for *arbitrary* nonempty $I \subset \{0, \dots, m\}$.

For any nonempty I , $\mathcal{F}_{I,D_m}$ is a subset of $\{\text{faces of type } I\} \times \mathbb{Z}/2\mathbb{Z}$, and we say that $(\mathbf{v}, \gamma) \in \mathcal{F}_{I,D_m}$ is μ -spin-permissible if $\gamma = \varepsilon(\mu)$ and $\mu_k^\mathbf{v}$ satisfies (SP1), (SP2), and (SP3) for all $k \in I$. If I satisfies (8.13.3), then this definition of μ -spin-permissibility coincides with the one in $\widetilde{W}_{D_m}/W_{I,D_m}$ via the identification $\widetilde{W}_{D_m}/W_{I,D_m} \xrightarrow{\sim} \mathcal{F}_{I,D_m}$. For any I , we write $\text{SPerm}_{\mathcal{F}_{I,D_m}}(\mu)$ for the set of μ -spin-permissible elements in $\mathcal{F}_{I,D_m}$.

If $\mathbf{v}$ is a 2-face of type I and $k \in 2m\mathbb{Z} \pm I$, then let l denote the unique element in I that is congruent mod $2m$ to k or $-k$. Then by (4.3.2) and (4.3.4), μ_k equals μ_l or $2 - \mu_l^*$. Hence

- μ_k satisfies (SP1) $\iff \mu_l$ does;
- when these equivalent conditions hold, and using (4.4.3) in the case that $\mu_k = 2 - \mu_l^*$, μ_k satisfies (SP2) $\iff \mu_l$ does; and
- μ_k satisfies (SP3) $\iff \mu_l$ does.

In particular, an element $(\mathbf{v}, \varepsilon(\mu)) \in \mathcal{F}_{I,D_m}$ is μ -spin-permissible $\iff \mu_k$ satisfies (SP1), (SP2), and (SP3) for all $k \in 2m\mathbb{Z} \pm I$.

Lemma 8.14.6. *Let $I \subset J$ be nonempty subsets of $\{0, \dots, m\}$. Then the natural map $\mathcal{F}_{J,D_m} \rightarrow \mathcal{F}_{I,D_m}$ carries $\text{SPerm}_{\mathcal{F}_{J,D_m}}(\mu)$ surjectively onto $\text{SPerm}_{\mathcal{F}_{I,D_m}}(\mu)$.*

Proof. To prove the lemma, it suffices to assume that

$$2m\mathbb{Z} \pm J = (2m\mathbb{Z} \pm I) \amalg (2m\mathbb{Z} \pm \{i+1\})$$

for some $i \in 2m\mathbb{Z} \pm I$. Let $(\mathbf{v}, \varepsilon(\mu)) \in \text{SPerm}_{\mathcal{F}_{I,D_m}}(\mu)$, with $\mathbf{v} = (v_k)_{k \in 2m\mathbb{Z} \pm I}$. To lift this element to $\text{SPerm}_{\mathcal{F}_{J,D_m}}(\mu)$, our task is to extend $\mathbf{v}$ to a face of type J for which $\mu_k^\mathbf{v}$ satisfies (SP1), (SP2), and (SP3) for all $k \in J$. Of course, once we have extended $\mathbf{v}$, it suffices to check just that $\mu_{i+1}^\mathbf{v}$ satisfies (SP1), (SP2), and (SP3).

Let j denote the minimal element in $2m\mathbb{Z} \pm I$ which is $> i$. Then $j - i \geq 2$. Let π denote the canonical projection

$$\pi: \{\text{faces of type } J\} \longrightarrow \{\text{faces of type } I\}.$$

Kottwitz and Rapoport [KR, Lem. 10.3] show that there is a bijection³

$$\begin{aligned} \pi^{-1}(\mathbf{v}) &\xrightarrow{\sim} \{v \in \mathbb{Z}^{2m} \mid v_i \geq v \geq v_j \text{ and } \Sigma v = \Sigma v_i - 1\} \\ (v_k)_{k \in 2m\mathbb{Z} \pm J} &\longmapsto v_{i+1}. \end{aligned}$$

Thus to extend $\mathbf{v}$ to a face of type J is to define v_{i+1} subject to the conditions in the definition of the target set in the display. To ensure that moreover

$$\mu_{i+1} := v_{i+1} - \omega_{i+1}$$

satisfies conditions (SP1), (SP2), and (SP3), we shall largely follow Görtz's proof of surjectivity between μ -permissible sets for GL_n [G3, Lem. 8]. Indeed Görtz shows that one can always find a v_{i+1} that extends $\mathbf{v}$ to a face of type J such that $\mu_{i+1} \in \text{Conv}(S_{2m}\mu)$; this last condition is equivalent, for our μ , to μ_{i+1} satisfying (SP1) and (SP2). Unfortunately μ_{i+1} need not satisfy the spin condition, which will cause us some complications. For any k , let $\tilde{k}$ denote the unique element in $\{1, \dots, 2m\}$ congruent to $k \bmod 2m$. We have that $v_j(\widetilde{i+1})$ equals $v_i(\widetilde{i+1})$ or $v_i(\widetilde{i+1}) - 1$, and we shall consider these cases separately.

Case 1: $v_j(\widetilde{i+1}) = v_i(\widetilde{i+1}) - 1$. Let $v_{i+1} := v_i - e_{\widetilde{i+1}}$. Then v_{i+1} extends $\mathbf{v}$ to a face of type J , and $\mu_{i+1} = \mu_i^\vee$. Hence μ_{i+1} satisfies (SP1) and (SP2). If μ_{i+1} also satisfies (SP3) then we are done, so suppose μ_{i+1} fails (SP3). Then μ_{i+1} is self-dual and $\mu_{i+1} \not\equiv \mu \bmod Q_{D_m}^\vee$. Let

$$c_i := \#\{k \mid \mu_i(k) = 2\},$$

and define c_{i+1} likewise. By (8.4.2) $c_i = c_{i+1} \not\equiv q \bmod 2$. Hence $c_i < q$.

Now, since μ_i satisfies (SP3), there must exist $l_1 \in A_i$ and $l_2 \in B_i$ such that $\mu_i(l_1) = \mu_i(l_2) = 1$. And since μ_{i+1} fails (SP3), we must have $l_1, l_2 \in A_{i+1}$ and $\mu_{i+1}(k) \in \{0, 2\}$ for all $k \in B_{i+1}$, or the same statement with A_{i+1} and B_{i+1} interchanged. For convenience, let us assume the former; one proceeds in an analogous way in case of the latter. Then evidently $A_{i+1} = A_i \amalg \{l_2, l_2^*\}; \{l_2, l_2^*\} = \{\widetilde{i+1}, (\widetilde{i+1})^*\}$; and $B_{i+1} = B_i \setminus \{\widetilde{i+1}, (\widetilde{i+1})^*\}$. To proceed with our proof in Case 1, we shall consider separately the possibilities that $\mu_j^\vee = \mu_{i+1}$ and $\mu_j^\vee \neq \mu_{i+1}$.

Suppose that $\mu_j^\vee \neq \mu_{i+1}$. Then, since $\Sigma\mu_j^\vee = \Sigma\mu_{i+1}$, there exists $l \in \{1, \dots, 2m\}$ such that $\mu_j^\vee(l) = \mu_{i+1}(l) - 1$. This requires $l \notin \{\widetilde{i+1}, \widetilde{i+2}, \dots, \widetilde{j}\}$, whence $v_j(l) = v_{i+1}(l) - 1$. We define $v'_{i+1} := v_i - e_l$. Then v'_{i+1} extends $\mathbf{v}$ to a face of type J , and we must show that $\mu'_{i+1} := v'_{i+1} - \omega_{i+1}$ satisfies (SP1), (SP2), and (SP3). All of this follows easily from the formula

$$(8.14.7) \quad \mu'_{i+1} = \mu_i^\vee + e_{\widetilde{i+1}} - e_l.$$

Indeed, for (SP1), we need note only that $\mu_i^\vee$ and $\mu_j^\vee$ satisfy (SP1) by hypothesis, that

$$\mu'_{i+1}(\widetilde{i+1}) = \mu_i^\vee(\widetilde{i+1}) + 1 = 2,$$

³Here our description of the target set differs from that of Kottwitz–Rapoport due to our differing conventions in defining faces of type I .

and that

$$\mu'_{i+1}(l) = \mu_i^{\mathbf{v}}(l) - 1 = \mu_{i+1}(l) - 1 = \mu_j^{\mathbf{v}}(l) \in \{0, 1, 2\}.$$

For (SP2), recalling from earlier that $c_i < q$, (8.14.7) gives

$$c'_{i+1} := \#\{k \mid \mu'_{i+1}(k) = 2\} \leq c_i + 1 \leq q.$$

And for (SP3), μ'_{i+1} is self-dual only if $l = (\widetilde{i+1})^*$, in which case $c'_{i+1} = c_i + 1$. Then $\mu'_{i+1} \equiv \mu \pmod{Q_{D_m}^{\vee}}$ by (8.4.2).

Now suppose that $\mu_j^{\mathbf{v}} = \mu_{i+1}$. Then $v_j(k) = v_i(k) - 1$ for exactly the elements $k \in \{\widetilde{i+1}, \widetilde{i+2}, \dots, \widetilde{j}\}$. Since $A_i \subset A_{i+1}$, we have $\widetilde{i+1} \in \{1, \dots, m\}$, and

$$B_{i+1} = \{\widetilde{i+2}, \widetilde{i+3}, \dots, 2m - (\widetilde{i+1})\}.$$

(To be clear, we interpret this as $B_{i+1} = \emptyset$ if $\widetilde{i+1} = m$.) If $\widetilde{j} \in B_{i+1}$, then $B_j \subset B_{i+1}$, so that $\mu_j^{\mathbf{v}}(k) \in \{0, 2\}$ for all $k \in B_j$. Since $\mu_j^{\mathbf{v}} = \mu_{i+1} \not\equiv \mu \pmod{Q_{D_m}^{\vee}}$, this would violate the spin condition. So $\widetilde{j} \notin B_{i+1}$. By minimality of $j \in 2m\mathbb{Z} \pm I$, we therefore have $\widetilde{j} = 2m - (\widetilde{i+1}) + 1 = (\widetilde{i+1})^*$. We now define

$$v'_{i+1} := v_i - e_{\widetilde{j}} = v_i - e_{(\widetilde{i+1})^*}.$$

Then v'_{i+1} extends $\mathbf{v}$ to a face of type J , and one sees just as in the previous paragraph that $v'_{i+1} - \omega_{i+1}$ satisfies (SP1), (SP2), and (SP3). This completes the proof in Case 1.

Case 2: $v_j(\widetilde{i+1}) = v_i(\widetilde{i+1})$. Let $X := \{k \mid v_j(k) \neq v_i(k)\}$. Working with respect to the Kottwitz–Rapoport conventions on faces of type I , Görtz shows in his proof of [G3, Lem. 8]⁴ that by taking l to be any element of X for which

$$\mu_i^{\mathbf{v}}(l) = \min\{\mu_i^{\mathbf{v}}(k)\}_{k \in X},$$

the vector $v_{i+1} := v_i + e_l$ extends $\mathbf{v}$ to a face of type J for which μ_{i+1} satisfies (SP1) and (SP2). Translating this to our conventions, we get that if l is any element of X for which

$$\mu_i^{\mathbf{v}}(l) = \max\{\mu_i^{\mathbf{v}}(k)\}_{k \in X},$$

then the vector $v_{i+1} := v_i - e_l$ extends $\mathbf{v}$ to a face of type J for which μ_{i+1} satisfies (SP1) and (SP2). If μ_{i+1} is not self-dual, or if μ_{i+1} is self-dual and $\equiv \mu \pmod{Q_{D_m}^{\vee}}$, then it trivially satisfies (SP3). So let us suppose μ_{i+1} is self-dual and $\not\equiv \mu \pmod{Q_{D_m}^{\vee}}$.

Before continuing, let us note that $\#X = j - i$ and $\widetilde{i+1} \notin X$. Hence there exists $k' \in X \setminus \{\widetilde{i+1}, \widetilde{i+2}, \dots, \widetilde{j}\}$. Since $\mu_j^{\mathbf{v}}(k') = \mu_i^{\mathbf{v}}(k') - 1$ and $\mu_i^{\mathbf{v}}$ and $\mu_j^{\mathbf{v}}$ satisfy (SP1), we conclude that $\max\{\mu_i^{\mathbf{v}}(k)\}_{k \in X}$ equals 1 or 2.

To proceed with the proof in Case 2, first suppose that l is the unique element in X such that $\mu_i^{\mathbf{v}}(l) \neq 0$. We shall show that μ_{i+1} satisfies the spin condition. Since $\mu_j^{\mathbf{v}}$ satisfies (SP1) and $\widetilde{i+1} \notin X$, we must have

$$X \setminus \{l\} \subset \{\widetilde{i+2}, \widetilde{i+3}, \dots, \widetilde{j}\},$$

and by counting cardinalities we see that this inclusion is an equality. Hence $\mu_{i+1} = \mu_j^{\mathbf{v}}$, and $\mu_i^{\mathbf{v}}(k) = \mu_{i+1}(k) = \mu_j^{\mathbf{v}}(k) = 0$ for all $k \in \{\widetilde{i+2}, \widetilde{i+3}, \dots, \widetilde{j}\}$. Since $\mathbf{v}$ is

⁴For the reader's convenience, we note that Görtz takes permutations to act on vectors on the *right*, so that $\sigma \cdot \sum_i b_i e_i = \sum_i b_i e_{\sigma^{-1}(i)}$ ($b_i \in \mathbb{R}$). It also appears that, in his notation, he means to define m to equal $\sigma(\widetilde{m})$, not $\sigma(\overline{m})$; the changes to his proof needed to accommodate this correction are trivial.

a 2-face and $\mu_j^\vee$ is self-dual, we must have that $\{\widetilde{i+2}, \widetilde{i+3}, \dots, \widetilde{j}\}$ is contained in $\{1, \dots, m\}$ or in $\{m+1, \dots, 2m\}$. Let us assume the former; one proceeds in an analogous way in case of the latter. Then $A_j = \{1, \dots, \widetilde{j}, \widetilde{j}^*, \dots, 2m\}$ and $B_j = \{\widetilde{j}+1, \dots, 2m - \widetilde{j}\}$. Since $\mu_j^\vee$ satisfies the spin condition and $\mu_j^\vee = \mu_{i+1} \not\equiv \mu \pmod{Q_{D_m}^\vee}$, there exist $k_1 \in A_j$ and $k_2 \in B_j$ such that $\mu_j^\vee(k_1) = \mu_j^\vee(k_2) = 1$. Evidently $k_1 \in A_j \setminus \{\widetilde{i+2}, \dots, \widetilde{j}, \widetilde{j}^*, \dots, (\widetilde{i+2})^*\}$. It follows that $\widetilde{i+1} \in \{1, \dots, m\}$, that $A_{i+1} = \{1, \dots, \widetilde{i+1}, (\widetilde{i+1})^*, \dots, m\}$, and that $B_{i+1} = \{\widetilde{i+2}, \dots, 2m - (\widetilde{i+1})\}$. Plainly $k_1 \in A_{i+1}$ and $k_2 \in B_{i+1}$, so that μ_{i+1} satisfies (SP3).

Finally suppose that there exists another element $l' \in X$ distinct from l and such that $\mu_i^\vee(l') \neq 0$. Of course if μ_{i+1} already satisfies (SP3) then there is nothing further to do, so let us suppose that it does not. Let $v'_{i+1} := v_i - e_{l'}$ and $\mu'_{i+1} := v'_{i+1} - \omega_{i+1}$, and define c_{i+1} and c'_{i+1} as in Case 1. Then v'_{i+1} extends $\mathbf{v}$ to another face of type J , $c_{i+1} \not\equiv q \pmod{2}$ by (8.4.2), and $c_{i+1} < q$. Our task is to show that μ'_{i+1} satisfies (SP1), (SP2), and (SP3). We have

$$\mu'_{i+1} = \mu_i^\vee + e_{\widetilde{i+1}} - e_{l'} = \mu_{i+1} + e_l - e_{l'}.$$

Since $\mu_i^\vee$ and μ_{i+1} satisfy (SP1) and the elements $\widetilde{i+1}$, l , l' are pairwise distinct, these expressions make clear that $0 \leq \mu'_{i+1}(k) \leq 2$ for all k except possibly $k = \widetilde{l'}$; and since $\mu_i^\vee(l') \in \{1, 2\}$ by hypothesis, we have $\mu'_{i+1}(l') \in \{0, 1\}$. Hence μ'_{i+1} satisfies (SP1). Since $c_{i+1} < q$, we also read off from the display that

$$c'_{i+1} \leq c_i + 1 \leq q,$$

so that μ'_{i+1} satisfies (SP2). It remains to see that μ'_{i+1} satisfies (SP3). But since μ_{i+1} is self-dual by assumption, μ'_{i+1} is self-dual only if $l' = l^*$. In this case $c'_{i+1} = c_i \pm 1$ and $\mu'_{i+1} \equiv \mu \pmod{Q_{D_m}^\vee}$ by (8.4.2), which completes the proof. $\square$

Remark 8.14.8. For $I \subset J$ nonempty and satisfying (8.13.3), we have shown directly, in (8.14.4) and (8.14.6), that the map

$$\text{SPerm}_{D_m}(\mu) \longrightarrow \text{SPerm}_{J,I,D_m}(\mu)$$

is surjective. This statement can be sharpened: for $w \in \text{SPerm}_{J,I,D_m}(\mu)$, let ${}^Jw^I$ denote the minimal length representative of w in $\widetilde{W}_{D_m}$. Then ${}^Jw^I \in \text{SPerm}_{D_m}(\mu)$. Indeed, since we've already shown that the displayed map is surjective, the sharpened statement is an immediate consequence of the fact that $\text{SPerm}_{D_m}(\mu)$ is closed in the Bruhat order (8.10.1).

8.15. Vertexwise admissibility. In this subsection we recall the notion of μ -vertexwise admissibility for any cocharacter μ in any based root datum, and we show that for μ the cocharacter (8.0.1) in $\mathcal{R}_{D_m}$, the notions of μ -admissibility and μ -vertexwise admissibility coincide.

Let $\mathcal{R}$ be a based root datum, let $\widetilde{W}$ denote its extended affine Weyl group, and let μ be a cocharacter for $\mathcal{R}$. Fix a base alcove $\mathbf{a}$, and let $\mathbf{f} \preceq \mathbf{f}'$ be subfacets of $\mathbf{a}$. The choice of $\mathbf{a}$ endows $W_{\mathbf{f}'} \backslash \widetilde{W} / W_{\mathbf{f}}$ with a Bruhat order, as in §5.2. For any subfacet $\mathbf{v} \preceq \mathbf{f}$, the natural map

$$\rho_{\mathbf{f}', \mathbf{f}; \mathbf{v}} : W_{\mathbf{f}'} \backslash \widetilde{W} / W_{\mathbf{f}} \longrightarrow W_{\mathbf{v}} \backslash \widetilde{W} / W_{\mathbf{v}}$$

carries (by definition) the set $\text{Adm}_{\mathbf{f}', \mathbf{f}}(\mu)$ surjectively to the set $\text{Adm}_{\mathbf{v}, \mathbf{v}}(\mu)$. We then have the following definition.

Definition 8.15.1 ([PRS, Def. 4.5.2]). For any subfacets $\mathbf{f} \preceq \mathbf{f}' \preceq \mathbf{a}$, the μ -vertexwise admissible set is the subset of $W_{\mathbf{f}'} \backslash \widetilde{W} / W_{\mathbf{f}}$

$$\text{Adm}_{\mathbf{f}', \mathbf{f}}^{\text{vert}}(\mu) := \bigcap_{\substack{\text{minimal} \\ \mathbf{v} \preceq \mathbf{f}}} \rho_{\mathbf{f}', \mathbf{f}; \mathbf{v}}^{-1}(\text{Adm}_{\mathbf{v}, \mathbf{v}}(\mu)).$$

In other words, an element in $W_{\mathbf{f}'} \backslash \widetilde{W} / W_{\mathbf{f}}$ is μ -vertexwise admissible if its image in $W_{\mathbf{v}} \backslash \widetilde{W} / W_{\mathbf{v}}$ is μ -admissible for all minimal subfacets $\mathbf{v}$ of $\mathbf{f}$. When $\mathcal{R}$ is semisimple, that is, when the root lattice has finite index in the character group, the minimal subfacets of $\mathbf{f}$ are vertices, i.e. points, which motivates the terminology. Strictly speaking, vertexwise admissibility is only defined in [PRS] when $\mathbf{f} = \mathbf{f}'$, but the added bit of generality we allow here will cause us no trouble, at least in the particular situations we consider later.

Since $\rho_{\mathbf{f}', \mathbf{f}; \mathbf{v}}$ carries $\text{Adm}_{\mathbf{f}', \mathbf{f}}(\mu)$ to $\text{Adm}_{\mathbf{v}, \mathbf{v}}(\mu)$, there is a tautological inclusion

$$(8.15.2) \quad \text{Adm}_{\mathbf{f}', \mathbf{f}}(\mu) \subset \text{Adm}_{\mathbf{v}, \mathbf{v}}^{\text{vert}}(\mu).$$

It is conjectured in [PRS, Conj. 4.5.3] (at least for $\mathbf{f} = \mathbf{f}'$) that in many cases this inclusion is an equality.⁵ Indeed no examples whatsoever are known where equality fails; and equality is known to hold in all situations arising from local models that have been studied,⁶ and furthermore whenever μ -admissibility and μ -permissibility are equivalent in $W_{\mathbf{f}'} \backslash \widetilde{W} / W_{\mathbf{f}}$. In some sense the notion of μ -vertexwise admissibility is intended as a replacement for μ -permissibility (which of course is not equivalent to μ -admissibility in general), in that μ -vertexwise admissibility is a manifestly “vertexwise” condition which is supposed to characterize μ -admissibility, at least in many cases.

Now let us specialize to the case $\mathcal{R} = \mathcal{R}_{D_m}$ and μ the cocharacter (8.0.1). We shall show that the inclusion (8.15.2) is indeed an equality. We write $\text{Adm}_{F', F, D_m}^{\text{vert}}(\mu)$ for the μ -vertexwise admissible set in $W_{F', D_m} \backslash \widetilde{W}_{D_m} / W_{F, D_m}$.

Theorem 8.15.3. *Let μ be the cocharacter (8.0.1), and let $F \preceq F' \preceq A_{D_m}$. Then*

$$\text{Adm}_{F', F, D_m}(\mu) = \text{Adm}_{F', F, D_m}^{\text{vert}}(\mu).$$

Proof. After applying one or both of the automorphisms θ_0 and θ_m (8.13.1) as needed, we may assume that F and F' satisfy property (8.13.2), as discussed at the beginning of §8.14. Thus we may assume that $W_{F, B_m} = W_{I, B_m}$ and $W_{F', B_m} = W_{J, B_m}$ for some nonempty subsets $I \subset J \subset \{0, \dots, m\}$ satisfying property (8.13.3). The proof will now follow fairly easily from our characterization of μ -admissibility in terms of μ -spin-permissibility (8.14.5).

We must show that $\text{Adm}_{F', F, D_m}^{\text{vert}}(\mu) \subset \text{Adm}_{F', F, D_m}(\mu)$. So let $w \in \text{Adm}_{F', F, D_m}^{\text{vert}}(\mu)$. Let $\tilde{w}$ be any lift of w to $\widetilde{W}_{D_m}$. By (8.14.5), we must show that $\tilde{w} \equiv t_\mu \pmod{W_{\text{aff}, D_m}}$ and that $\nu_k^{\tilde{w}}$ satisfies (SP1'), (SP2'), and (SP3') for all $k \in I$. Of course μ -vertexwise admissibility immediately implies the first of these conditions.

⁵We refer the reader to [PRS] for the precise formulation of the conjecture, which is given in group-theoretic terms.

⁶In the situation arising from the local models studied in this paper, the equality $\text{Adm}_{\mathbf{f}', \mathbf{f}}(\mu) = \text{Adm}_{\mathbf{f}', \mathbf{f}}^{\text{vert}}(\mu)$ will be proved in §9.7.

Now, the minimal subfacets of F are the lines

$$\begin{aligned} a_k + \mathbb{R} \cdot (1, \dots, 1) & \text{ for } k \in I \setminus \{1, m-1\}; \\ a_{0'} + \mathbb{R} \cdot (1, \dots, 1) & \text{ if } 1 \in I; \\ a_{m'} + \mathbb{R} \cdot (1, \dots, 1) & \text{ if } m-1 \in I. \end{aligned}$$

So by vertexwise admissibility and (8.14.5) applied to $W_{\{k\}, D_m} \backslash \widetilde{W}_{D_m} / W_{\{k\}, D_m}$, we conclude that $\nu_k^{\widetilde{w}}$ satisfies (SP1'), (SP2'), and (SP3') for $k \in I \setminus \{1, m-1\}$. It remains to consider the possibility that 1, $m-1$, or both is in I .

If $1 \in I$, then $a_0, a_{0'} \in \overline{F}$. Let $W_{\{0'\}, D_m}$ denote the stabilizer of $a_{0'}$ in W_{aff, D_m} . Since $a_{0'}$ is a special vertex, we have

$$\widetilde{W}_{D_m} = X_{*D_m} \rtimes W_{\{0'\}, D_m}$$

and

$$W_{\{0'\}, D_m} \backslash \widetilde{W}_{D_m} / W_{\{0'\}, D_m} \cong W_{\{0'\}, D_m} \backslash X_{*D_m},$$

which we identify with the set $X_{*D_m}^+$ of dominant elements in X_{*D_m} . For $x = t_{\nu_0^x} \sigma$ with $\nu_0^x \in X_{*D_m}$ and $\sigma \in S_{2m}^o$, we have $t_{a_{0'} - \sigma a_{0'}} \sigma \in W_{\{0'\}, D_m}$. Hence $x = (t_{\nu_0^x + \sigma a_{0'} - a_{0'}}, t_{a_{0'} - \sigma a_{0'}} \sigma)$ according to the semidirect product decomposition $\widetilde{W}_{D_m} = X_{*D_m} \rtimes W_{\{0'\}, D_m}$. Hence the map

$$\widetilde{W}_{D_m} \longrightarrow W_{\{0'\}, D_m} \backslash \widetilde{W}_{D_m} / W_{\{0'\}, D_m} \cong X_{*D_m}^+$$

sends

$$x \longmapsto \text{dominant form of } \nu_{0'}^x,$$

where

$$\nu_{0'}^x := \nu_0^x + \sigma a_{0'} - a_{0'} = x a_{0'} - a_{0'}.$$

The Bruhat order on $W_{\{0'\}, D_m} \backslash \widetilde{W}_{D_m} / W_{\{0'\}, D_m}$ is just given by the dominance order on $X_{*D_m}^+$. Thus our assumption that the image of $\widetilde{w}$ in $W_{\{0'\}, D_m} \backslash \widetilde{W}_{D_m} / W_{\{0'\}, D_m}$ is μ -admissible says exactly that $\nu_{0'}^{\widetilde{w}}$ satisfies (SP1') and (SP2') and $\nu_{0'}^{\widetilde{w}} \equiv \mu \pmod{Q_{D_m}^\vee}$. This implies that $\nu_{0'}^{\widetilde{w}} \in \text{Conv}(S_{2m}^o \mu)$. Since we also have $0 \in I$ and we have already shown that $\nu_0^{\widetilde{w}}$ satisfies (SP1')–(SP3'), we conclude from (8.6.3) that $\nu_1^{\widetilde{w}}$ satisfies (SP1')–(SP3').

If $m-1 \in I$, then one shows in a similar fashion that $\nu_{m-1}^{\widetilde{w}}$ satisfies (SP1')–(SP3'), using a_m in place of a_0 and $a_{m'}$ in place of $a_{0'}$. $\square$

Remark 8.15.4. Our original definition of μ -spin-permissibility (8.2.1), and its parahoric variant (8.14.1), is formulated in analogy with the definition of the Rapoport–Zink local model [RZ], which is in terms of totally ordered *chains* (or more generally, multichains) of lattices. However, from a root-theoretic perspective it is unnatural to emphasize the vectors μ_1^w and μ_{m-1}^w , or what amounts to the same, the vectors ν_1^w and ν_{m-1}^w , since the points a_1 and a_{m-1} are not contained in minimal facets. Instead it is more natural to consider the vectors $wa_k - a_k$ for all k in some nonempty subset I of $\{0, 0', 2, 3, \dots, m-2, m, m'\}$. We have just shown in the preceding theorem that if $F \preceq F'$ are the subfacets of A_{D_m} whose closures contain the vertices a_k for respective nonempty subsets $I \subset J$ of $\{0, 0', 2, 3, \dots, m-2, m, m'\}$, then an element w is in $\text{Adm}_{F', F, D_m}(\mu) \iff w \equiv t_\mu \pmod{W_{\text{aff}, D_m}}$ and $\widetilde{w}a_k - a_k$ satisfies (SP1'), (SP2'), and (SP3') for all $k \in I$, for any representative $\widetilde{w} \in \widetilde{W}_{D_m} / W_{F, D_m}$; here for $k \in \{0, 0', m, m'\}$, the

spin condition (SP3') just means $\tilde{w}a_k - a_k \equiv \mu \bmod Q_{D_m}^\vee$ (which is equivalent to $w \equiv t_\mu \bmod W_{\text{aff}, D_m}$).

8.16. μ -permissibility in the parahoric case. In this final subsection of §8, we generalize (8.6.4) to give a characterization of μ -permissibility in $\mathcal{R}_{D_m}$ in the general parahoric case. See (5.3.2) for the general definition. We write

$$\text{Perm}_{F', F, D_m}(\mu)$$

for the set of μ -permissible elements in $W_{F', D_m} \backslash \widetilde{W}_{D_m} / W_{F, D_m}$, and we abbreviate this to $\text{Perm}_{D_m}(\mu)$ when $F = F' = A_{B_m}$. As has been the case previously, we shall incur no loss of generality by restricting to subfacets of the base alcove A_{D_m} satisfying property (8.13.2), so that the corresponding subgroups of W_{aff, D_m} are of the form W_{I, D_m} .

Proposition 8.16.1. *Let μ be the cocharacter (8.0.1), let $I \subset J$ be nonempty subsets of $\{0, \dots, m\}$ satisfying (8.13.3), and let $w \in W_{J, D_m} \backslash \widetilde{W}_{D_m} / W_{I, D_m}$. Then w is μ -permissible $\iff w \equiv t_\mu \bmod W_{\text{aff}, D_m}$ and for one, hence any, representative $\tilde{w}$ of w in $\widetilde{W}_{D_m} / W_{I, D_m}$, $\nu_k^{\tilde{w}}$ satisfies (SP1') and (SP2') for all $k \in I$, and (SP3') for all $k \in I \cap \{0, 1, m-1, m\}$.*

Proof. As in the proof of (8.6.4), this is obvious from (8.6.1) and (8.6.3). $\square$

We conclude by generalizing (8.6.6) to the general parahoric case.

Proposition 8.16.2. *Let μ be the cocharacter (8.0.1), let $F \preceq F' \preceq A_{D_m}$, and suppose that $q \leq 2$. Then*

$$\text{Adm}_{F', F, D_m}(\mu) = \text{Perm}_{F', F, D_m}(\mu).$$

Proof. For $\theta \in \{\theta_0, \theta_m, \theta_0\theta_m\}$, conjugation by θ induces isomorphisms

$$\text{Adm}_{F', F, D_m}(\mu) \xrightarrow{\sim} \text{Adm}_{\theta F', \theta F, D_m}(\mu)$$

and

$$\text{Perm}_{F', F, D_m}(\mu) \xrightarrow{\sim} \text{Perm}_{\theta F', \theta F, D_m}(\mu).$$

So it suffices to prove the proposition in the case that F and F' satisfy (8.13.2). This is done as in (8.6.6), using (8.14.5) in place of (8.2.2). $\square$

9. TYPE B COMBINATORICS

In this final section of the paper, we extend all of the combinatorial results from the previous section to type B . The key result is (9.6.1), which characterizes μ -admissibility in terms of a type B analog of μ -spin-permissibility, where μ is the cocharacter below; it is this result that is fed directly into the proofs of Theorems 1.3 and 1.4, as explained earlier in the proof of (7.5.1).

We fix an integer $m \geq 1$. In §9.1 we use the symbol μ to denote an arbitrary cocharacter, but from §9.2 on we take

$$(9.0.1) \quad \mu := (2^q, 1^{2m+1-2q}, 0^{(q)}), \quad 0 \leq q \leq m,$$

except where specified to the contrary. For $i \in \{1, \dots, 2m+1\}$, we write $i^* = 2m+2-i$. We denote by $e_1, \dots, e_{2m+1}$ the standard basis in $\mathbb{Z}^{2m+1}$.

9.1. Steinberg fixed-point root data. We begin §9 with some generalities on Steinberg fixed-point root data. We take as our main references the papers of Steinberg [St], Kottwitz–Rapoport [KR], and Haines–Ngô [HN], especially [HN, §9].

Let us briefly recall what we need from the theory of Steinberg fixed-point root data. Let $\mathcal{R} = (X^*, X_*, R, R^\vee, \Pi)$ be a reduced based root datum.⁷ Attached to $\mathcal{R}$ are its Weyl group W , its affine Weyl group W_{aff} , and its extended affine Weyl group $\widetilde{W} := X_* \rtimes W$, as in §5.1. An *automorphism* Θ of $\mathcal{R}$ is an automorphism Θ of the abelian group X_* that stabilizes the set of coroots $R^\vee$ and such that the subsets $R, \Pi \subset X^*$ are stable under the dual automorphism Θ^* of X^* induced by Θ and the perfect pairing $X^* \times X_* \rightarrow \mathbb{Z}$. It follows that any automorphism of $\mathcal{R}$ induces automorphisms of W, W_{aff} , and $\widetilde{W}$. Attached to Θ is the *Steinberg fixed-point root datum* $\mathcal{R}^{[\Theta]} = (X^{*[\Theta]}, X_*^{[\Theta]}, R^{[\Theta]}, R^{\vee[\Theta]}, \Pi^{[\Theta]})$; this is a reduced based root datum described explicitly in [HN, §9]. We systematically use a superscript $[\Theta]$ to denote the analogs for $\mathcal{R}^{[\Theta]}$ of all the objects defined for $\mathcal{R}$. For our purposes, we shall just mention that $\widetilde{W}^{[\Theta]}$ is naturally a subgroup of $\widetilde{W}$, with $W^{[\Theta]}$ and $W_{\text{aff}}^{[\Theta]}$ equal to the respective fixed-point subgroups W^Θ and W_{aff}^Θ , and that the cocharacter group $X_*^{[\Theta]}$ is the subgroup of X_*

$$X_*^{[\Theta]} := \{x \in X_* \mid \Theta(x) \equiv x \bmod Z\},$$

where $Z := \{x \in X_* \mid \langle \alpha, x \rangle = 0 \text{ for all } \alpha \in R\}$.

Let $\mathcal{A} := X_* \otimes \mathbb{R}$ and $\mathcal{A}^{[\Theta]} := X_*^{[\Theta]} \otimes \mathbb{R}$ denote the respective apartments for $\mathcal{R}$ and $\mathcal{R}^{[\Theta]}$. For each facet $\mathbf{f}$ in $\mathcal{A}$, let

$$\mathbf{f}^{[\Theta]} := \mathbf{f} \cap \mathcal{A}^{[\Theta]}.$$

We then have the following obvious generalization of [HN, Prop. 9.3], which follows from the fact that the affine root hyperplanes for $\mathcal{R}^{[\Theta]}$ in $\mathcal{A}^{[\Theta]}$ are exactly the intersections of the affine root hyperplanes for $\mathcal{R}$ with $\mathcal{A}^{[\Theta]}$.

Lemma 9.1.1.

- (i) For any facet $\mathbf{f}$ in $\mathcal{A}$, the set $\mathbf{f}^{[\Theta]}$ is either empty or a facet in $\mathcal{A}^{[\Theta]}$.
- (ii) Every facet in $\mathcal{A}^{[\Theta]}$ is of the form $\mathbf{f}^{[\Theta]}$ for a unique facet $\mathbf{f}$ in $\mathcal{A}$.
- (iii) The rule $\mathbf{a} \mapsto \mathbf{a}^{[\Theta]}$ is a bijection from the set of alcoves in $\mathcal{A}$ that meet $\mathcal{A}^{[\Theta]}$ to the set of alcoves in $\mathcal{A}^{[\Theta]}$. $\square$

As something of a contrast to (iii), it can happen that non-minimal facets in $\mathcal{A}$ intersect with $\mathcal{A}^{[\Theta]}$ to give minimal facets in $\mathcal{A}^{[\Theta]}$.

For any facet $\mathbf{f}$ in $\mathcal{A}$ that meets $\mathcal{A}^{[\Theta]}$, it follows from the lemma that the stabilizer group $W_{\mathbf{f}^{[\Theta]}} \subset W_{\text{aff}}^{[\Theta]} = W_{\text{aff}}^\Theta$ is given by

$$W_{\mathbf{f}^{[\Theta]}} = W_{\mathbf{f}}^\Theta,$$

where as in §5.1 $W_{\mathbf{f}}$ denotes the stabilizer group of $\mathbf{f}$ in W_{aff} . We then have the following.

Lemma 9.1.2. *Let $\mathbf{f}_1$ and $\mathbf{f}_2$ be facets in $\mathcal{A}$ that meet $\mathcal{A}^{[\Theta]}$. Then the natural map*

$$W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}} \longrightarrow W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$$

is an injection.

⁷Strictly speaking, [HN] also assumes that $\mathcal{R}$ is irreducible, but we shall not cite any facts from [HN] that require this assumption.

Note that in the special case that $\mathbf{f}_1 = \mathbf{f}_2$ is a special minimal facet in $\mathcal{A}$ meeting $\mathcal{A}^{[\Theta]}$, the displayed sets of double cosets identify with the sets of dominant cocharacters in $X_*^{[\Theta]}$ and X_* , respectively, relative to the choice of a positive Weyl chamber in $\mathcal{A}$ that meets $\mathcal{A}^{[\Theta]}$; and the displayed map is then the natural injection on dominant cocharacters.

Proof of (9.1.2). Let us begin with a few generalities. By a “hyperplane” in $\mathcal{A}$ we shall mean an “affine root hyperplane for $\mathcal{R}$.” For any subset A in $\mathcal{A}$, we say that a hyperplane H is an A -hyperplane if H contains A ; we let $\mathcal{H}_A$ denote the (possibly empty) set of A -hyperplanes in $\mathcal{A}$; and we let W_A denote the (possibly trivial) subgroup of W_{aff} generated by the reflections across the A -hyperplanes. We refer to the connected components of the set $\mathcal{A} \setminus \bigcup_{H \in \mathcal{H}_A} H$ as A -chambers. By general properties of reflection groups, W_A acts simply transitively on the set of A -chambers. If $A \subset \mathcal{A}^{[\Theta]}$, then Θ acts on $\mathcal{H}_A$, and hence on the set of A -chambers; an A -chamber C meets $\mathcal{A}^{[\Theta]}$ if and only if C is Θ -stable; and W_A^Θ acts simply transitively on the set of Θ -stable A -chambers.

To prove the lemma, our problem is to show that if $w' = x_1 w x_2$ with $w, w' \in \widetilde{W}^{[\Theta]}$, $x_1 \in W_{\mathbf{f}_1}$, and $x_2 \in W_{\mathbf{f}_2}$, then there exist $y_1 \in W_{\mathbf{f}_1}^\Theta$ and $y_2 \in W_{\mathbf{f}_2}^\Theta$ such that $w' = y_1 w y_2$. Let $\mathbf{a}$ be an alcove in $\mathcal{A}$ meeting $\mathcal{A}^{[\Theta]}$ and such that $\mathbf{f}_2 \preceq \mathbf{a}$. Let $\Omega_{\mathbf{a}}$ (resp. $\Omega_{\mathbf{a}^{[\Theta]}}$) denote the stabilizer of $\mathbf{a}$ (resp. $\mathbf{a}^{[\Theta]}$) in $\widetilde{W}$ (resp. $\widetilde{W}^{[\Theta]}$), so that $\widetilde{W} = W_{\text{aff}} \rtimes \Omega_{\mathbf{a}}$ and $\widetilde{W}^{[\Theta]} = W_{\text{aff}}^\Theta \rtimes \Omega_{\mathbf{a}^{[\Theta]}}$. Write $w = x\omega$ and $w' = x'\omega'$ with $x, x' \in W_{\text{aff}}^\Theta$ and $\omega, \omega' \in \Omega_{\mathbf{a}^{[\Theta]}}$. Then $\omega = \omega'$ and

$$x'\omega = w' = x_1 w x_2 = x_1 x \omega x_2 \omega^{-1} \omega.$$

We have $\omega x_2 \omega^{-1} \in W_{\omega \mathbf{f}_2}$, and $\omega \mathbf{f}_2$ meets $\mathcal{A}^{[\Theta]}$ since $\omega \in \widetilde{W}^{[\Theta]}$. Thus it suffices to solve the analogous problem (with $\omega \mathbf{f}_2$ in place of $\mathbf{f}_2$) for the equation $x' = x_1 x (\omega x_2 \omega^{-1})$. So, since $\mathbf{f}_2$ is arbitrary, we might as well assume that $w, w' \in W_{\text{aff}}^\Theta$ to begin with, which we shall do for the rest of the proof.

Consider the alcoves $x_1 w \mathbf{a}$ and $w' \mathbf{a}$, which share the common subfacet $x_1 w \mathbf{f}_2$. Let A denote the intersection of the $\mathbf{f}_1$ -hyperplanes containing $x_1 w \mathbf{f}_2$. If A is empty, which is to say that there are no $\mathbf{f}_1$ -hyperplanes containing $x_1 w \mathbf{f}_2$, then $x_1 w \mathbf{a}$ and $w' \mathbf{a}$ must be contained in the same $\mathbf{f}_1$ -chamber C , since any hyperplane that separates two facets in $\mathcal{A}$ must contain all the common subfacets of both. Then C and $x_1^{-1}C$ both meet $\mathcal{A}^{[\Theta]}$, since they respectively contain $w' \mathbf{a}$ and $w \mathbf{a}$. Hence $x_1 \in W_{\mathbf{f}_1}^\Theta$, and we conclude immediately that $x_2 \in W_{\mathbf{f}_2}^\Theta$ as well.

If A is nonempty, then tautologically $\mathbf{f}_1 \subset A$, whence $W_A \subset W_{\mathbf{f}_1}$. Let z_1 be the unique element in W_A such that $z_1 x_1 w \mathbf{a}$ is contained in the same A -chamber as $w' \mathbf{a}$. Since $x_1 w \mathbf{f}_2 \subset A$ and A is fixed by the elements of W_A , we have $x_1 w \mathbf{f}_2 \preceq z_1 x_1 w \mathbf{a}$. Since we also have $x_1 w \mathbf{f}_2 \preceq w' \mathbf{a}$, we conclude from the definition of A that $z_1 x_1 w \mathbf{a}$ and $w' \mathbf{a}$ must be contained in the same $\mathbf{f}_1$ -chamber. As in the previous paragraph, it follows that $y_1 := z_1 x_1 \in W_{\mathbf{f}_1}^\Theta$. Since the alcoves that share the subfacet $x_1 w \mathbf{f}_2$ with $y_1 w \mathbf{a}$ are precisely those of the form $y_1 w y \mathbf{a}$ for $y \in W_{\mathbf{f}_2}$, there exists $y_2 \in W_{\mathbf{f}_2}$ such that $y_1 w y_2 = w'$, and it follows immediately that y_2 is Θ -fixed as well. $\square$

Now fix an alcove $\mathbf{a}$ in $\mathcal{A}$, and consider the associated Bruhat order $\leq$ and length function on $\widetilde{W}$, as in §5.2. Let $\mathbf{f}_1$ and $\mathbf{f}_2$ be subfacets of $\mathbf{a}$. Then every double coset $w \in W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ contains a unique representative ${}^{\mathbf{f}_1} w {}^{\mathbf{f}_2}$ of minimal length in $\widetilde{W}$, which moreover has the property that ${}^{\mathbf{f}_1} w {}^{\mathbf{f}_2} \leq w'$ for all $w' \in w$. In the Steinberg fixed-point setting, we then have the following result.

Lemma 9.1.3. *Let $\mathbf{f}_1, \mathbf{f}_2 \preceq \mathbf{a}$, and suppose that all three meet $\mathcal{A}^{[\Theta]}$. Let $w \in W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ be a double coset contained in the image of $W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}}$. Then $\mathbf{f}_1 w \mathbf{f}_2 \in \widetilde{W}^{[\Theta]}$.*

Proof. This is proved for example in [Sm3, Lem. 6.2.1]. $\square$

We use the lemma to obtain the following double coset version of a result of Kottwitz–Rapoport and Haines–Ngô.

Theorem 9.1.4. *Let $\mathbf{a}, \mathbf{f}_1$, and $\mathbf{f}_2$ be as in (9.1.3), and let $\leq^{[\Theta]}$ denote the Bruhat order on $W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}}$ determined by the alcove $\mathbf{a}^{[\Theta]} \subset \mathcal{A}^{[\Theta]}$. Regard $W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}}$ as a subset of $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$ via (9.1.2), and let $w, x \in W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}}$. Then*

$$w \leq^{[\Theta]} x \text{ in } W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}} \iff w \leq x \text{ in } W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}.$$

Proof. By definition of the Bruhat order on $W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2}$,

$$w \leq x \text{ in } W_{\mathbf{f}_1} \backslash \widetilde{W} / W_{\mathbf{f}_2} \iff \mathbf{f}_1 w \mathbf{f}_2 \leq \mathbf{f}_1 x \mathbf{f}_2 \text{ in } \widetilde{W}.$$

By (9.1.3), $\mathbf{f}_1 w \mathbf{f}_2, \mathbf{f}_1 x \mathbf{f}_2 \in \widetilde{W}^{[\Theta]}$. Hence by the Iwahori version of the theorem [HN, Prop. 9.6] (which generalizes the version of the theorem for affine Weyl groups proved by Kottwitz–Rapoport [KR, Prop. 2.3]), the right-hand condition in the display holds

$$\iff \mathbf{f}_1 w \mathbf{f}_2 \leq^{[\Theta]} \mathbf{f}_1 x \mathbf{f}_2 \text{ in } \widetilde{W}^{[\Theta]}.$$

By the Iwahori version of the theorem again, $\mathbf{f}_1 w \mathbf{f}_2$ and $\mathbf{f}_1 x \mathbf{f}_2$ are the minimal length representatives in $\widetilde{W}^{[\Theta]}$ of their respective $(W_{\mathbf{f}_1^{[\Theta]}}, W_{\mathbf{f}_2^{[\Theta]}})$ -cosets. Hence the last display holds

$$\iff w \leq^{[\Theta]} x \text{ in } W_{\mathbf{f}_1^{[\Theta]}} \backslash \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}}. \quad \square$$

For use later on when we come to μ -permissibility, we cite the following result, which is proved for example in [Sm3, Lem. 6.5.1].

Lemma 9.1.5. *Let μ be any cocharacter in $X_*^{[\Theta]}$. Then*

$$\text{Conv}(W^{[\Theta]}\mu) = \text{Conv}(W\mu) \cap \mathcal{A}^{[\Theta]}. \quad \square$$

9.2. The root datum $\mathcal{R}_{B_m}$. In this subsection we define the root datum $\mathcal{R}_{B_m}$. Let

$$\begin{aligned} X_{*B_m} &:= X_{*2m+1} \\ &= \left\{ (x_1, \dots, x_{2m+1}) \in \mathbb{Z}^{2m+1} \mid \begin{array}{l} x_1 + x_{2m+1} = x_2 + x_{2m} = \dots = \\ = x_m + x_{m+2} = 2x_{m+1} \end{array} \right\}, \end{aligned}$$

where we recall X_{*2m+1} from (4.1.1), and

$$X_{B_m}^* := \mathbb{Z}^{2m+1} / \{(x_1, \dots, x_m, -2 \sum_{i=1}^m x_i, x_m, \dots, x_1)\}.$$

Then the standard dot product on $\mathbb{Z}^{2m+1}$ induces a perfect pairing $X_{B_m}^* \times X_{*B_m} \rightarrow \mathbb{Z}$. For $1 \leq i, j \leq 2m+1$ with $j \neq i, i^*$, let

$$\begin{aligned} \alpha_{i,j}: \quad X_{*B_m} &\longrightarrow \mathbb{Z} \\ (x_1, \dots, x_{2m+1}) &\longmapsto x_i - x_j \end{aligned}$$

and

$$\alpha_{i,j}^\vee := \begin{cases} e_i - e_j + e_{j^*} - e_{i^*}, & i, j \neq m+1; \\ 2e_i - 2e_{i^*}, & j = m+1; \\ -2e_j + 2e_{j^*}, & i = m+1. \end{cases}$$

Then we may regard $\alpha_{i,j}$ as an element of $X_{B_m}^* \cong \text{Hom}(X_{*B_m}, \mathbb{Z})$, and we let

$$(9.2.1) \quad \Phi_{B_m} := \{\alpha_{i,j}\}_{j \neq i, i^*} \subset X_{B_m}^* \quad \text{and} \quad \Phi_{B_m}^\vee := \{\alpha_{i,j}^\vee\}_{j \neq i, i^*} \subset X_{*B_m}.$$

The objects so defined form a root datum

$$\mathcal{R}_{B_m} := (X_{B_m}^*, X_{*B_m}, \Phi_{B_m}, \Phi_{B_m}^\vee),$$

which is the root datum for split GO_{2m+1} . We take as simple roots

$$\alpha_{1,2}, \alpha_{2,3}, \dots, \alpha_{m,m+1}.$$

The Weyl group of $\mathcal{R}_{B_m}$ identifies canonically with S_{2m+1}^* . As discussed in §5.1, we then have the extended affine Weyl group

$$\widetilde{W}_{B_m} := X_{*B_m} \rtimes S_{2m+1}^*.$$

The affine Weyl group W_{aff, B_m} is the subgroup $Q_{B_m}^\vee \rtimes S_{2m+1}^* \subset \widetilde{W}_{B_m}$, where $Q_{B_m}^\vee \subset X_{*B_m}$ is coroot lattice. Explicitly,

$$Q_{B_m}^\vee = \left\{ (x_1, \dots, x_{2m+1}) \in X_{*B_m} \mid \begin{array}{l} x_i + x_{i^*} = 0 \text{ for all } i \text{ and} \\ x_1 + \dots + x_m \text{ is even} \end{array} \right\}.$$

Let

$$\mathcal{A}_{B_m} := X_{*B_m} \otimes_{\mathbb{Z}} \mathbb{R} \subset \mathbb{R}^{2m+1}.$$

We take as positive Weyl chamber the chamber in $\mathcal{A}_{B_m}$ on which the simple roots are all positive. We take as our base alcove

$$(9.2.2) \quad A_{B_m} := \left\{ (x_1, \dots, x_{2m+1}) \in \mathbb{R}^{2m+1} \mid \begin{array}{l} x_1 + x_{2m+1} = \dots = x_m + x_{m+2} = 2x_{m+1} \\ \text{and } x_1, x_{2m+1} - 1 < x_2 < x_3 < \dots < x_{m+1} \end{array} \right\};$$

note that this is the unique alcove contained in the Weyl chamber *opposite* the positive chamber and whose closure contains the origin. The minimal facets of A_{B_m} are the lines

$$a + \mathbb{R} \cdot (1, \dots, 1)$$

for a one of the vertices

$$(9.2.3) \quad \begin{aligned} a_k &:= \left((-\tfrac{1}{2})^{(k)}, 0^{(2m+1-2k)}, (\tfrac{1}{2})^{(k)} \right) \quad \text{for } k = 0, 2, 3, \dots, m, \\ a_{0'} &:= (-1, 0^{2m-1}, 1). \end{aligned}$$

As discussed in §5.2, our choice of A_{B_m} endows W_{aff, B_m} and $\widetilde{W}_{B_m}$ with Bruhat orders. We also define

$$a_1 := \left(-\tfrac{1}{2}, 0^{(2m-1)}, \tfrac{1}{2} \right).$$

Note that in terms of our earlier notation (4.1.2), we have $\widetilde{W}_{B_m} = \widetilde{W}_{2m}$ on the nose. Thus for each $w \in \widetilde{W}_{B_m}$ we get we get a face $\mathbf{v}$ of type $(2m+1, \{0, \dots, m\})$ by letting w act on the standard face $\omega_{\{0, \dots, m\}}$, as in §4.2. We write μ_k^w for the vector $\mu_k^\vee$ attached to this face,

$$\mu_k^w := w\omega_k - \omega_k \quad \text{for } k \in \mathbb{Z}.$$

The rule $w \mapsto w \cdot \omega_{\{0, \dots, m\}}$ identifies $\widetilde{W}_{B_m}$ with the faces of type $(2m+1, \{0, \dots, m\})$.

9.3. Relation to $\mathcal{R}_{D_{m+1}}$. Consider the root datum $\mathcal{R}_{D_{m+1}}$ defined in §8.1, and let $\Theta := \theta_{m+1}$ be the automorphism on $X_{*D_{m+1}}$ defined in (8.13.1) (with m replaced by $m+1$),

$$\Theta: (x_1, \dots, x_{2m+2}) \mapsto (x_1, \dots, x_m, x_{m+2}, x_{m+1}, x_{m+3}, x_{m+4}, \dots, x_{2m+2}).$$

Then Θ is an automorphism of $\mathcal{R}_{D_{m+1}}$ in the sense of §9.1, where we take the roots (8.1.3) (with m replaced by $m+1$) as simple roots in $\Phi_{D_{m+1}}$. The map on cocharacters

$$\begin{aligned} \iota: X_{*B_m} &\xrightarrow{\quad\quad\quad} X_{*D_{m+1}} \\ (x_1, \dots, x_{2m+1}) &\mapsto (x_1, \dots, x_m, x_{m+1}, x_{m+1}, x_{m+2}, \dots, x_{2m+1}) \end{aligned}$$

identifies $X_{*B_m} \xrightarrow{\sim} X_{*D_{m+1}}^{[\Theta]}$, and we leave it to the reader to verify that ι induces $\mathcal{R}_{B_m} \simeq \mathcal{R}_{D_{m+1}}^{[\Theta]}$.

Regarding $\mathcal{A}_{B_m} = X_{*B_m} \otimes \mathbb{R}$ as a subspace of $\mathcal{A}_{D_{m+1}} = X_{*D_{m+1}} \otimes \mathbb{R}$ via $\iota \otimes \text{id}_{\mathbb{R}}$, note that $A_{B_m} = A_{D_{m+1}} \cap \mathcal{A}_{B_m}$. Moreover, $\iota \otimes \text{id}_{\mathbb{R}}$ identifies each of the points $a_0, a_0', a_2, \dots, a_m$ defined in (9.2.3) with the point in $\mathcal{A}_{D_{m+1}}$ denoted by the same symbol, as defined in (8.1.4) and (8.5.1) (with m replaced by $m+1$).

The Θ -fixed Weyl group $(S_{2m+2}^{\circ})^{[\Theta]} = (S_{2m+2}^{\circ})^{\Theta}$ consists of the $\sigma \in S_{2m+2}^{\circ}$ such that $\sigma(m+1) \in \{m+1, m+2\}$. It identifies with S_{2m+1}^* via restriction to X_{*B_m} along ι . Let us describe the induced embedding $S_{2m+1}^* \hookrightarrow S_{2m+2}^{\circ}$ more explicitly. Given $\sigma \in S_{2m+1}^*$, σ necessarily fixes $m+1$, and we view σ as a permutation on $\{1, \dots, m, m+3, \dots, 2m+2\}$ by identifying

$$\begin{aligned} \{1, \dots, m, m+2, \dots, 2m+1\} &\xrightarrow{\sim} \{1, \dots, m, m+3, \dots, 2m+2\} \\ i &\longmapsto \begin{cases} i, & i \leq m; \\ i+1, & m+2 \leq i. \end{cases} \end{aligned}$$

To define σ on all of $\{1, \dots, 2m+2\}$, we then declare that σ fixes or interchanges $m+1$ and $m+2$ as needed to make σ even in S_{2m+2} .

Writing ι' for the embedding $S_{2m+1} \hookrightarrow S_{2m+2}^{\circ}$ just described, ι and ι' combine to specify the embedding

$$\phi: \widetilde{W}_{B_m} = X_{*B_m} \rtimes S_{2m+1}^* \xrightarrow{\iota \rtimes \iota'} X_{*D_{m+1}} \rtimes S_{2m+2}^{\circ} = \widetilde{W}_{D_{m+1}}.$$

Let $w \in \widetilde{W}_{B_m}$ and $0 \leq k \leq m$. In terms of our identifications of $\widetilde{W}_{B_m}$ and $\widetilde{W}_{D_{m+1}}$ with faces of type $(2m+1, \{0, \dots, m\})$ and $(2m+2, \{0, \dots, m+1\})$, respectively, ϕ sends

$$(9.3.1) \quad \begin{aligned} \mu_k^w &= (x_1, \dots, x_{2m+1}) \mapsto \\ &(x_1, \dots, x_m, x_{m+1}, x_{m+1}, x_{m+2}, \dots, x_{2m+1}) = \mu_k^{\phi(w)}. \end{aligned}$$

In this way ϕ identifies $\widetilde{W}_{B_m}$ with the set of $x \in \widetilde{W}_{D_m}$ such that $\mu_k^x(m+1) = \mu_k^x(m+2)$ for all $0 \leq k \leq m$. (Note however that for such an x , one may have $\mu_{m+1}^x(m+1) \neq \mu_{m+1}^x(m+2)$.)

9.4. Parahoric variants. We continue to embed

$$X_{*B_m} \subset X_{*D_{m+1}}, \quad \mathcal{A}_{B_m} \subset \mathcal{A}_{D_{m+1}}, \quad \text{and} \quad \widetilde{W}_{B_m} \subset \widetilde{W}_{D_{m+1}}$$

as in the previous subsection. Let F be a subfacet of the base alcove A_{B_m} , and let W_{F, B_m} denote the common stabilizer and pointwise fixer of F in W_{aff, B_m} . By

(9.1.1) F is of the form $\tilde{F} \cap \mathcal{A}_{B_m}$ for a uniquely determined subfacet $\tilde{F}$ of $A_{D_{m+1}}$, and

$$W_{F,B_m} = W_{\tilde{F},D_{m+1}}^\Theta = W_{\tilde{F},D_{m+1}} \cap W_{\text{aff},B_m} = W_{\tilde{F},D_{m+1}} \cap \widetilde{W}_{B_m}.$$

The affine automorphism θ_0 (8.13.1) on $\mathcal{A}_{D_{m+1}}$ stabilizes $\mathcal{A}_{B_m}$ and interchanges a_0 and $a_{0'}$, while fixing a_k for all $2 \leq k \leq m$. In analogy with §8.13, by applying θ_0 as needed, we shall therefore incur no loss of generality by working only with facets F with the property that if $a_{0'} \in F$, then $a_0 \in F$.

Continuing the analogy with §8.13, given a subfacet $F \preceq A_{B_m}$ such that a_0 is a vertex of F if $a_{0'}$ is, let

$$I := \{k \in \{0, \dots, m\} \mid a_k \in \overline{F}\}$$

and

$$(9.4.1) \quad W_{I,B_m} := \{w \in W_{\text{aff},B_m} \mid wa_k = a_k \text{ for all } k \in I\}.$$

Then, arguing as in §8.13, the association $F \mapsto I$ gives a bijection from the set of such subfacets F to the set of nonempty subsets I of $\{0, \dots, m\}$ with the property that

$$(9.4.2) \quad 1 \in I \implies 0 \in I;$$

and for F corresponding to I , we have

$$W_{F,B_m} = W_{I,B_m}.$$

In terms of our notation (8.13.4) (with m replaced by $m+1$), we have

$$W_{I,B_m} = W_{\tilde{I},D_{m+1}}^\Theta = W_{\tilde{I},D_{m+1}} \cap \widetilde{W}_{B_m},$$

where $\tilde{I}$ is the nonempty subset of $\{0, \dots, m+1\}$

$$(9.4.3) \quad \tilde{I} := \begin{cases} I \cup \{m+1\}, & m \in I; \\ I, & m \notin I. \end{cases}$$

By the same reasoning as in the paragraph before (8.13.6), the elements of W_{I,B_m} fix the standard face ω_I of type $(2m+1, I)$. Thus to any $w \in \widetilde{W}_{B_m}/W_{I,B_m}$ we may attach a well-defined face $w \cdot \omega_I$ of type $(2m+1, I)$, and we write

$$\mu_k^w := w\omega_k - \omega_k \quad \text{for } k \in I.$$

The embedding $\phi: \widetilde{W}_{B_m} \subset \widetilde{W}_{D_{m+1}}$ induces an embedding

$$\widetilde{W}_{B_m}/W_{I,B_m} \subset \widetilde{W}_{D_{m+1}}/W_{\tilde{I},D_{m+1}},$$

and of course the corresponding map on μ_k -vectors is again given by (9.3.1), for $k \in I$. By definition of W_{I,B_m} , for each $w \in \widetilde{W}_{B_m}/W_{I,B_m}$ we also get a well-defined vector

$$\nu_k^w := wa_k - a_k \quad \text{for } k \in I.$$

If $1 \in I$, then the elements of W_{I,B_m} also fix $a_{0'}$ (because $a_{0'}$ is a vertex of the facet containing a_1), and we define

$$\nu_{0'}^w := wa_{0'} - a_{0'}$$

for $w \in \widetilde{W}_{B_m}/W_{I,B_m}$.

9.5. μ -spin-permissibility. In this subsection we define μ -spin-permissibility in $\widetilde{W}_{B_m}$ and its parahoric variants. We continue with the notation of the previous subsections. Recall the set $A_k \subset \{1, \dots, 2m+1\}$ from (4.3.6).

Definition 9.5.1. Let μ be the cocharacter (9.0.1), and let $I \subset J$ be nonempty subsets of $\{0, \dots, m\}$ satisfying property (9.4.2). We say that $w \in W_{J,B_m} \setminus \widetilde{W}_{B_m}/W_{I,B_m}$ is μ -spin-permissible if $w \equiv t_\mu \bmod W_{\text{aff},B_m}$ and for one, hence any, representative $\tilde{w}$ of w in $\widetilde{W}_{B_m}/W_{I,B_m}$, $\tilde{w}$ satisfies the following conditions for all $k \in I$.

- (sp1) $\mu_k^{\tilde{w}} + (\mu_{-k}^{\tilde{w}})^* = \mathbf{2}$ and $\mathbf{0} \leq \mu_k^{\tilde{w}} \leq \mathbf{2}$.
- (sp2) $\#\{j \mid \mu_k^{\tilde{w}}(j) = 2\} \leq q$.
- (sp3) (spin condition) If $\mu_k^{\tilde{w}}$ is self-dual and $\mu_k^{\tilde{w}} \not\equiv \mu \bmod Q_{B_m}^\vee$, then there exists $j \in A_k$ such that $\mu_k^{\tilde{w}}(j) = 1$.

We write

$$\text{SPerm}_{J,I,B_m}(\mu)$$

for the set of μ -spin-permissible elements in $W_{J,B_m} \setminus \widetilde{W}_{B_m}/W_{I,B_m}$. We abbreviate this to $\text{SPerm}_{I,B_m}(\mu)$ when $J = \{0, \dots, m\}$, and to $\text{SPerm}_{B_m}(\mu)$ when $I = J = \{0, \dots, m\}$.

In analogy with the type D situation, we shall verify in a moment that the definition does not depend on the choice of representative $\tilde{w}$. Let us first make a couple of other remarks.

First, the condition $w \equiv t_\mu \bmod W_{\text{aff},B_m}$ is well-defined because $W_{J,B_m}, W_{I,B_m} \subset W_{\text{aff},B_m}$. And, as in the type D setting, it is implied by the spin condition when $\mathbf{0} \in I$, since $A_0 = \emptyset$.

Second, in contrast with the spin condition (SP3) for $\mathcal{B}_{D_m}$, the set B_k does not appear in (sp3). Indeed if (sp1) holds, then necessarily $\mu_k^{\tilde{w}}(m+1) = 1$ by (4.3.8). Since $m+1 \in B_k$ for k in the range $0 \leq k \leq m$, the existence of a $j' \in B_k$ such that $\mu_k^{\tilde{w}}(j') = 1$ is then automatic. This point can also be understood via the following lemma.

Lemma 9.5.2. *Let I be a nonempty subset of $\{0, \dots, m\}$ satisfying property (9.4.2), let $w \in \widetilde{W}_{B_m}/W_{I,B_m}$, and consider the embedding*

$$\phi: \widetilde{W}_{B_m}/W_{I,B_m} \hookrightarrow \widetilde{W}_{D_{m+1}}/W_{\tilde{I},D_{m+1}},$$

with $\tilde{I}$ as in (9.4.3). Then

- (i) $w \equiv t_\mu \bmod W_{\text{aff},B_m} \iff \phi(w) \equiv t_{\iota(\mu)} \bmod W_{\text{aff},D_{m+1}}$;
- (ii) for $k \in I$, μ_k^w satisfies (sp1) $\iff \mu_k^{\phi(w)}$ satisfies (SP1); and
- (iii) for $k \in I$, if μ_k^w satisfies (sp1), then μ_k^w satisfies (sp2) (resp. (sp3)) relative to $\mu \iff \mu_k^{\phi(w)}$ satisfies (SP2) (resp. (SP3)) relative to $\iota(\mu)$.

Furthermore ϕ gives an identification

$$\text{SPerm}_{I,B_m}(\mu) = \text{SPerm}_{\tilde{I},D_{m+1}}(\iota(\mu)) \cap \widetilde{W}_{B_m}/W_{I,B_m}.$$

Proof. Part (i) is obvious because, by general properties of fixed-point root data, $W_{\text{aff},B_m} = W_{\text{aff},D_{m+1}} \cap \widetilde{W}_{B_m}$. Parts (ii) and (iii) are obvious from the explicit form of the map on μ_k -vectors (9.3.1). Thus the real content of the lemma is contained in the final asserted equality, which is an immediate consequence of (i)–(iii) and the following lemma. $\square$

Lemma 9.5.3. *Under the assumptions of the previous lemma, suppose that $m \in I$, μ_m^w satisfies (sp1) and (sp2), and $w \equiv t_\mu \bmod W_{\text{aff}, B_m}$. Then μ_m^w also satisfies (sp3), and $\mu_{m+1}^{\phi(w)}$ satisfies (SP1)–(SP3).*

Proof. Before getting started, we note that since μ_m^w satisfies (sp1), we have

$$(9.5.4) \quad \mu_m^{\phi(w)}(m+1) = \mu_m^{\phi(w)}(m+2) = 1.$$

Let us first show that μ_m^w satisfies (sp3), or, equivalently, that $\mu_m^{\phi(w)}$ satisfies (SP3). Recall the integer $c_m^{\phi(w)}$ from (8.4.1). If $c_m^{\phi(w)} \equiv q \bmod 2$, then $\mu_m^{\phi(w)}$ trivially satisfies (SP3) by (8.4.2). If $c_m^{\phi(w)} \not\equiv q \bmod 2$, then $c_m^{\phi(w)} < q$ since $\mu_m^{\phi(w)}$ satisfies (SP2). And since $q \leq m$, $\#A_m = 2m$, and $\mu_k^{\phi(w)}$ satisfies (SP1), there must exist $j \in A_m$ such that $\mu_k^{\phi(w)}(j) = 1$. This combines with (9.5.4) to show that $\mu_m^{\phi(w)}$ satisfies (SP3).

It remains to show that $\mu_{m+1}^{\phi(w)}$ satisfies (SP1)–(SP3). It follows from the form of the embedding $\iota': S_{2m+1}^* \hookrightarrow S_{2m+2}^\circ$ in §9.3 and the recursion relation (8.3.2) that

$$\mu_{m+1}^{\phi(w)} = \mu_m^{\phi(w)} \quad \text{or} \quad \mu_{m+1}^{\phi(w)} = \mu_m^{\phi(w)} + e'_{m+1} - e'_{m+2},$$

where $e'_1, \dots, e'_{2m+2}$ denotes the standard basis in $\mathbb{Z}^{2m+2}$. Since $\mu_m^{\phi(w)}$ satisfies (SP1), we conclude, with the help of (9.5.4) in the second case, that $\mu_{m+1}^{\phi(w)}$ satisfies (SP1). Furthermore, since $x \equiv t_{\mu_{m+1}^x} \bmod W_{\text{aff}, D_{m+1}}$ for any $x \in \widetilde{W}_{D_{m+1}}/\widetilde{W}_{\tilde{I}, D_{m+1}}$, we find that $t_{\mu_{m+1}^{\phi(w)}} \equiv \phi(w) \equiv t_{\iota(\mu)} \bmod W_{\text{aff}, D_{m+1}}$. Hence $\mu_{m+1}^{\phi(w)}$ automatically satisfies (SP3), and by (8.4.2)

$$c_{m+1}^{\phi(w)} \equiv q \bmod 2.$$

Since $\mu_m^{\phi(w)}$ satisfies (SP2), we have

$$c_m^{\phi(w)} \leq q,$$

and by (8.4.3)

$$c_{m+1}^{\phi(w)} \leq c_m^{\phi(w)} + 1.$$

These last three displays imply $c_{m+1}^{\phi(w)} \leq q$, which completes the proof. $\square$

Using (9.5.2), it is now easy to see that for $w \in W_{J, B_m} \setminus \widetilde{W}_{B_m}/W_{I, B_m}$, conditions (sp1)–(sp3) are independent of the choice of representative $\tilde{w} \in \widetilde{W}_{B_m}/W_{I, B_m}$. Indeed, $\phi(W_{J, B_m}) \subset W_{\tilde{J}, D_{m+1}}$, and we then need note only that conditions (SP1)–(SP3), for $k \in \tilde{I}$, are independent of the choice of element in $W_{\tilde{J}, D_{m+1}}\phi(\tilde{w}) \subset \widetilde{W}_{D_{m+1}}/\widetilde{W}_{\tilde{I}, D_{m+1}}$ by (8.5.7) and (8.14.4).

Let us conclude the subsection by reformulating conditions (sp1)–(sp3) in terms of the vector ν_k^w . Let $w \in \widetilde{W}_{B_m}/W_{I, B_m}$, and consider the following conditions for $k \in I$.

- (sp1') $\nu_k^w + (\nu_k^w)^* = \mathbf{2}$ and $\mathbf{0} \leq \nu_k^w \leq \mathbf{2}$.
- (sp2') $\#\{j \mid \nu_k^w(j) = 2\} + \#\{j \mid \nu_k^w(j) \notin \mathbb{Z}\}/4 \leq q$.
- (sp3') (spin condition) If $\nu_k^w \in \mathbb{Z}^{2m}$ and $\nu_k^w \not\equiv \mu \bmod Q_{B_m}^\vee$, then there exists $j \in A_k$ such that $\nu_k^w(j) = 1$.

The following equivalences are all proved as in (8.5.7), or can be deduced from (8.5.7) via the embedding $\widetilde{W}_{B_m} \hookrightarrow \widetilde{W}_{D_{m+1}}$.

Lemma 9.5.5. *Let $w \in \widetilde{W}_{B_m}/W_{I,B_m}$ and $k \in I$*

- (i) μ_k^w satisfies (sp1) $\iff \nu_k^w$ satisfies (sp1').
- (ii) *Suppose that the equivalent conditions in (i) hold. Then μ_k^w satisfies (sp2) $\iff \nu_k^w$ satisfies (sp2').*
- (iii) μ_k^w satisfies (sp3) $\iff \nu_k^w$ satisfies (sp3'). □

9.6. μ -spin-permissibility and μ -admissibility. In this subsection we prove the equivalence of μ -spin-permissibility and μ -admissibility for $\mathcal{R}_{B_m}$, for μ the cocharacter (9.0.1), in analogy with the result for $\mathcal{R}_{D_m}$ (8.14.5). We continue with the notation of the previous subsections. We write $\text{Adm}_{J,I,B_m}(\mu)$ for the set of μ -admissible elements in $W_{J,B_m} \setminus \widetilde{W}_{B_m}/W_{I,B_m}$, and we abbreviate this to $\text{Adm}_{B_m}(\mu)$ when $I = J = \{0, \dots, m\}$.

Theorem 9.6.1. *Let μ be the cocharacter (9.0.1), and let $I \subset J$ be nonempty subsets of $\{0, \dots, m\}$ satisfying property (9.4.2). Then*

$$\text{Adm}_{J,I,B_m}(\mu) = \text{SPerm}_{J,I,B_m}(\mu).$$

Proof. This will follow easily from our results in type D and from (9.1.4). For notational convenience, we regard X_{*B_m} as a subgroup of $X_{*D_{m+1}}$, and $\widetilde{W}_{B_m}$ as a subgroup of $\widetilde{W}_{D_{m+1}}$. The inclusion $\text{Adm}_{J,I,B_m}(\mu) \subset \text{SPerm}_{J,I,B_m}(\mu)$ is an immediate consequence of (8.14.5), (9.1.4), and (9.5.2).

To prove the reverse inclusion, let $w \in \text{SPerm}_{J,I,B_m}(\mu)$. Then

$$w \in \text{SPerm}_{\tilde{J}, \tilde{I}, D_{m+1}}(\mu)$$

by (9.5.2). Let $\tilde{J}w\tilde{I}$ denote the minimal length representative of w in $\widetilde{W}_{D_{m+1}}$. Then

$$\tilde{J}w\tilde{I} \in \text{SPerm}_{D_{m+1}}(\mu) \cap \widetilde{W}_{B_m}$$

by (8.14.8) and (9.1.3). By applying (8.12.1)(ii) repeatedly, we eventually find a translation element $t_\nu \in \text{SPerm}_{D_{m+1}}(\mu)$ such that $\tilde{J}x\tilde{I} \leq t_\nu$ in $\widetilde{W}_{D_{m+1}}$ and $\nu(m+1) = \nu(m+2) = 1$ (regarding ν as a $(2m+2)$ -tuple), i.e. $t_\nu \in \widetilde{W}_{B_m}$. Hence $t_\nu \in \text{SPerm}_{B_m}(\mu)$ by (9.5.2), and $\tilde{J}w\tilde{I} \leq t_\nu$ in $\widetilde{W}_{B_m}$ by (9.1.4). Since the μ -spin-permissible translation elements in $\widetilde{W}_{B_m}$ are μ -admissible by the same argument as in (8.2.3), we conclude that $w \in \text{Adm}_{J,I,B_m}(\mu)$. □

The key point in the above proof, besides the fact (9.1.4) that the Bruhat order for $\mathcal{R}_{B_m}$ is inherited from the Bruhat order for $\mathcal{R}_{D_{m+1}}$, is that given $w \in \widetilde{W}_{B_m}$ such that $w \leq t_\nu$ in $\widetilde{W}_{D_{m+1}}$ for some $\nu \in S_{2m+2}^\circ\mu$, there in fact exists $\nu' \in S_{2m+1}^*\mu$ such that $w \leq t_{\nu'}$; or in other words, that $\text{Adm}_{B_m}(\mu) = \widetilde{W}_{B_m} \cap \text{Adm}_{D_{m+1}}(\mu)$. We conjecture that the same holds for any Steinberg fixed-point root datum, as follows.

Conjecture 9.6.2. *Let $\mathcal{R} = (X^*, X_*, R, R^\vee, \Pi)$ be a based root datum and Θ an automorphism of $\mathcal{R}$, as in §9.1. Let $\mathcal{A}$ denote the apartment of $\mathcal{R}$, let $\mathbf{a}$ be an alcove in $\mathcal{A}$ meeting $\mathcal{A}^{[\Theta]}$, and let $\mathbf{f}_1, \mathbf{f}_2 \preceq \mathbf{a}$ be subfacets also meeting $\mathcal{A}^{[\Theta]}$. Then for any cocharacter $\mu \in X_*^{[\Theta]}$,*

$$\text{Adm}_{\mathbf{f}_1^{[\Theta]}, \mathbf{f}_2^{[\Theta]}, \mathcal{R}^{[\Theta]}}(\mu) = \text{Adm}_{\mathbf{f}_1, \mathbf{f}_2, \mathcal{R}}(\mu) \cap (W_{\mathbf{f}_1^{[\Theta]}} \setminus \widetilde{W}^{[\Theta]} / W_{\mathbf{f}_2^{[\Theta]}}),$$

where the Bruhat orders in the admissible sets are specified by $\mathbf{a}^{[\Theta]}$ and $\mathbf{a}$, respectively.

The content of the conjecture lies in the inclusion $\supset$, as the forward inclusion $\subset$ is a trivial consequence of (9.1.4). It suffices to prove the conjecture just in the Iwahori case $\mathbf{f}_1 = \mathbf{f}_2 = \mathbf{a}$, since the sets $\text{Adm}_{\mathcal{R}^{[\Theta]}}(\mu)$ and $\text{Adm}_{\mathcal{R}}(\mu) \cap \widetilde{W}^{[\Theta]}$ respectively surject onto the left-hand and right-hand sets in the display. If $\mathcal{R}$ is irreducible of type A , then the conjecture has been proved by Haines and Ngô for any μ [HN, Th. 1, Prop. 9.7]. The conjecture also holds whenever μ -admissibility and μ -permissibility are equivalent in $\widetilde{W}$ and in $\widetilde{W}^{[\Theta]}$, since it is an easy consequence of (9.1.5) that $\text{Perm}_{\mathcal{R}}(\mu) \cap \widetilde{W}^{[\Theta]} \subset \text{Perm}_{\mathcal{R}^{[\Theta]}}(\mu)$. In particular, the conjecture is known for *minuscule* μ when the root datums involve only types A , B , C , and D ; see [PRS, Prop. 4.4.5(iii)]. As we remarked above, we proved the conjecture for $\mathcal{R} = \mathcal{R}_{D_{m+1}}$, $\mathcal{R}^{[\Theta]} = \mathcal{R}_{B_m}$, and μ the cocharacter (9.0.1) in the course of proving (9.6.1). Our proof of this relied crucially on (8.12.1)(ii). If however the conjecture were already known, then it would have obviated our need for (8.12.1)(ii) in the first place.

9.7. Vertexwise admissibility. In this subsection we prove the equivalence of μ -admissibility and μ -vertexwise admissibility in $\mathcal{R}_{B_m}$ for the cocharacter μ (9.0.1), in analogy with (8.15.3). See (8.15.1) for the general definition of μ -vertexwise admissibility. We write $\text{Adm}_{F',F,B_m}^{\text{vert}}(\mu)$ for the μ -vertexwise admissible set in $W_{F',B_m} \setminus \widetilde{W}_{B_m} / W_{F,B_m}$.

Theorem 9.7.1. *Let μ be the cocharacter (9.0.1), and let $F \preceq F' \preceq A_{B_m}$. Then*

$$\text{Adm}_{F',F,B_m}(\mu) = \text{Adm}_{F',F,B_m}^{\text{vert}}(\mu).$$

Proof. The argument is very similar to the proof of (8.15.3). After applying the automorphism θ_0 if necessary, we may assume that if $a_{0'}$ is a vertex of F , then so is a_0 , and likewise for F' . Thus we may assume that $W_{F,B_m} = W_{I,B_m}$ and $W_{F',B_m} = W_{J,B_m}$ for some nonempty subsets $I \subset J \subset \{0, \dots, m\}$ satisfying property (9.4.2).

We must show that $\text{Adm}_{F',F,B_m}^{\text{vert}}(\mu) \subset \text{Adm}_{F',F,B_m}(\mu)$. If $w \in \text{Adm}_{F',F,B_m}^{\text{vert}}(\mu)$ and $\tilde{w}$ is any representative of w in $\widetilde{W}_{B_m} / W_{F,B_m}$, then we see right away that $w \equiv t_\mu \bmod W_{\text{aff},B_m}$ and, by (9.6.1) applied with $I = J = \{k\}$ for $k \in I \setminus \{1\}$, that $\mu_k^{\tilde{w}}$ satisfies (sp1)–(sp3) for $k \in I \setminus \{1\}$. Thus, by (9.6.1) again, it remains only to show that $\mu_1^{\tilde{w}}$ satisfies (sp1)–(sp3) in case $1 \in I$. This is done by using that the image of w in $W_{\{0'\},B_m} \setminus \widetilde{W}_{B_m} / W_{\{0'\},B_m}$ is μ -admissible, as in the proof of (8.15.3), where $W_{\{0'\},B_m}$ denotes the stabilizer in W_{aff,B_m} of the facet $a_{0'} + \mathbb{R} \cdot (1, \dots, 1)$. $\square$

Remark 9.7.2. Upon identifying $\widetilde{W}_G$ with $\widetilde{W}_{B_m}$ as in (5.7.4), the theorem was conjectured by Pappas and Rapoport in [PR4, Conj. 4.2].

Remark 9.7.3. In analogy with (8.15.4), from a root-theoretic point of view it would be more natural to define μ -spin-permissibility in terms of the vectors μ_k^w , or equivalently ν_k^w , for $k \in \{0, 0', 2, 3, \dots, m\}$, since the points a_k for such k are vertices of the base alcove. The theorem shows that $\text{Adm}_{F',F,B_m}(\mu)$ consists of all elements w such that $w \equiv t_\mu \bmod W_{\text{aff},B_m}$ and for one, hence any, representative $\tilde{w}$ of w in $\widetilde{W}_{B_m} / W_{F,B_m}$, $\tilde{w}a_k - a_k$ satisfies (sp1')–(sp3') for all a_k which are vertices of F . Here, analogously to (8.15.4), the spin condition (sp3') for $\tilde{w}a_{0'} - a_{0'}$ means that $\tilde{w}a_{0'} - a_{0'} \equiv \mu \bmod Q_{B_m}^\vee$ (which is equivalent to $w \equiv t_\mu \bmod W_{\text{aff},B_m}$).

9.8. μ -permissibility. In this final subsection of the paper, we shall characterize μ -permissibility in $\mathcal{R}_{B_m}$, for μ the cocharacter (9.0.1), in terms of the conditions (sp1)–(sp3). Everything we shall do will be in close analogy with §§8.6 and 8.16. We write

$$\text{Perm}_{F', F, B_m}(\mu)$$

for the set of μ -permissible elements in $W_{F', B_m} \backslash \widetilde{W}_{B_m} / W_{F, B_m}$, and we abbreviate this to $\text{Perm}_{B_m}(\mu)$ when $F = F' = A_{B_m}$. As usual, it is harmless to restrict to restrict to subfacets F of A_{B_m} with the property that a_0 is a vertex of F if $a_{0'}$ is.

Proposition 9.8.1. *Let μ be the cocharacter (9.0.1), let $I \subset J$ be nonempty subsets of $\{0, \dots, m\}$ satisfying property (9.4.2), and let $w \in W_{J, B_m} \backslash \widetilde{W}_{B_m} / W_{I, B_m}$. Then w is μ -permissible $\iff w \equiv t_\mu \bmod W_{\text{aff}, B_m}$ and for one, hence any, representative $\tilde{w}$ of w in $\widetilde{W}_{B_m} / W_{I, B_m}$, $\nu_k^{\tilde{w}}$ satisfies (sp1') and (sp2') for all $k \in I$, and (sp3') for all $k \in I \cap \{0, 1\}$.*

Proof. This is obvious from the following lemma. $\square$

Lemma 9.8.2. *Keep the notation of (9.8.1), and let $\text{Conv}(S_{2m+1}^* \mu)$ denote the convex hull in $\mathcal{A}_{B_m}$ of the Weyl orbit $S_{2m+1}^* \mu$.*

- (i) *Let $k \in I$. Then $\nu_k^{\tilde{w}}$ satisfies (sp1') and (sp2') $\iff \nu_k^{\tilde{w}} \in \text{Conv}(S_{2m+1}^* \mu)$.*
- (ii) *Suppose that $1 \in I$. Then $\nu_0^{\tilde{w}}$ and $\nu_1^{\tilde{w}}$ satisfy (sp1')–(sp3') $\iff w \equiv t_\mu \bmod W_{\text{aff}, B_m}$ and $\nu_0^{\tilde{w}}, \nu_{0'}^{\tilde{w}} \in \text{Conv}(S_{2m+1}^* \mu)$.*

Proof. Upon identifying $\mathcal{R}_{B_m}$ with $\mathcal{R}_{D_{m+1}}^{[\Theta]}$ as in §9.3, everything follows from the analogous results (8.6.1) and (8.6.3) for $\mathcal{R}_{D_{m+1}}$, the equivalence of the (sp) and (SP) conditions, and the fact that $\text{Conv}(S_{2m+1}^* \mu) = \text{Conv}(S_{2m+1}^\circ \mu) \cap \mathcal{A}_{B_m}$ by (9.1.5). $\square$

Example 9.8.3. Recall the element $w \in \widetilde{W}_{D_4}$ from (8.6.5) which is contained in $\text{Perm}_{D_4}(\mu)$ but not in $\text{Adm}_{D_4}(\mu)$, for μ the cocharacter $(2^{(2)}, 1, 1, 0^{(3)}) \in X_{*D_4}$. Upon embedding $\widetilde{W}_{B_3} \subset \widetilde{W}_{D_4}$, we find that $w, t_\mu \in \widetilde{W}_{B_3}$, and that $w \in \text{Perm}_{B_m}(\mu)$ but $w \notin \text{Adm}_{B_m}(\mu)$. As in (8.6.5), this example easily generalizes to show that $\text{Adm}_{B_m}(\mu) \neq \text{Perm}_{B_m}(\mu)$ whenever $m, q \geq 3$. We remark that Haines and Ngô [HN, Th. 3] have previously shown that in any irreducible root datum $\mathcal{R}$ not of type A and of rank ≥ 4 , there exist cocharacters μ for which $\text{Adm}_{\mathcal{R}}(\mu) \neq \text{Perm}_{\mathcal{R}}(\mu)$.

In general, μ -permissible sets need not be well-behaved with regard to Steinberg fixed-point root data, in the sense that for an arbitrary cocharacter μ in $\mathcal{R}^{[\Theta]}$, the intersection $\text{Perm}_{\mathcal{R}}(\mu) \cap \widetilde{W}^{[\Theta]}$ may only be properly contained in $\text{Perm}_{\mathcal{R}^{[\Theta]}}(\mu)$, not equal. However, for μ the cocharacter (9.0.1), such bad behavior does not arise.

Proposition 9.8.4. *Let $F \preceq F'$ be subfacets of A_{B_m} , and let $\tilde{F}$ and $\tilde{F}'$ be the respective facets in $\mathcal{A}_{D_m}$ containing each of them. Then*

$$\text{Perm}_{F', F, B_m}(\mu) = \text{Perm}_{\tilde{F}', \tilde{F}, D_m}(\mu) \cap (W_{F', B_m} \backslash \widetilde{W}_{B_m} / W_{F, B_m}).$$

Proof. The inclusion $\supset$ holds by general properties of Steinberg fixed-point root data. To prove the forward inclusion $\subset$, by applying the automorphism θ_0 if necessary, we may assume that $a_{0'}$ is a vertex of F if a_0 is, and likewise for F' . The conclusion then follows at once from (9.5.2), (9.5.3), and our explicit descriptions of $\text{Perm}_{\tilde{F}', \tilde{F}, D_m}(\mu)$ (8.16.1) and $\text{Perm}_{F', F, B_m}(\mu)$ (9.8.1). $\square$

As a consequence of (9.8.4), (8.16.2), and the fact that we have proved (9.6.2) in the case at hand, we conclude the following.

Corollary 9.8.5. *Let μ be the cocharacter (9.0.1), let $F \preceq F' \preceq A_{B_m}$, and suppose that $q \leq 2$. Then*

$$\mathrm{Adm}_{F',F,B_m}(\mu) = \mathrm{Perm}_{F',F,B_m}(\mu). \quad \square$$

REFERENCES

- [Bo] M. Borovoi, *Abelian Galois cohomology of reductive groups*, Mem. Amer. Math. Soc. **132** (1998), no. 626, viii+50 pp.
- [B] N. Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.
- [G1] U. Görtz, *On the flatness of models of certain Shimura varieties of PEL-type*, Math. Ann. **321** (2001), no. 3, 689–727.
- [G2] ———, *On the flatness of local models for the symplectic group*, Adv. Math. **176** (2003), no. 1, 89–115.
- [G3] ———, *Topological flatness of local models in the ramified case*, Math. Z. **250** (2005), no. 4, 775–790.
- [HN] T. J. Haines and Ngô B. C., *Alcoves associated to special fibers of local models*, Amer. J. Math. **124** (2002), no. 6, 1125–1152.
- [HR] T. Haines and M. Rapoport, *On parahoric subgroups*, appendix to: G. Pappas and M. Rapoport, *Twisted loop groups and their affine flag varieties*, Adv. Math. **219** (2008), no. 1, 118–198.
- [Ki] M. Kisin, *Integral models for Shimura varieties of abelian type*, J. Amer. Math. Soc. **23** (2010), 967–1012.
- [Ko1] R. E. Kottwitz, *Points on some Shimura varieties over finite fields*, J. Amer. Math. Soc. **5** (1992), no. 2, 373–444.
- [Ko2] ———, *Isocrystals with additional structure. II*, Compositio Math. **109** (1997), no. 3, 255–339.
- [KR] R. Kottwitz and M. Rapoport, *Minuscule alcoves for GL_n and GSp_{2n}* , Manuscripta Math. **102** (2000), no. 4, 403–428.
- [P] G. Pappas, *On the arithmetic moduli schemes of PEL Shimura varieties*, J. Algebraic Geom. **9** (2000), no. 3, 577–605.
- [PR1] G. Pappas and M. Rapoport, *Local models in the ramified case. I. The EL-case*, J. Algebraic Geom. **12** (2003), no. 1, 107–145.
- [PR2] ———, *Local models in the ramified case. II. Splitting models*, Duke Math. J. **127** (2005), no. 2, 193–250.
- [PR3] ———, *Twisted loop groups and their affine flag varieties*, with an appendix by T. Haines and Rapoport, Adv. Math. **219** (2008), no. 1, 118–198.
- [PR4] ———, *Local models in the ramified case. III. Unitary groups*, J. Inst. Math. Jussieu **8** (2009), no. 3, 507–564.
- [PRS] G. Pappas, M. Rapoport, and B. Smithling, *Local models of Shimura varieties, I. Geometry and combinatorics*, to appear in *Handbook of Moduli*. [arXiv:1011.5551v3](#) [math.AG].
- [R] M. Rapoport, *A guide to the reduction modulo p of Shimura varieties*, Astérisque **298** (2005), 271–318 (English, with English and French summaries). Automorphic forms. I.
- [RZ] M. Rapoport and Th. Zink, *Period spaces for p -divisible groups*, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996.
- [R] T. Richarz, *Schubert varieties in twisted affine flag varieties and local models*, preprint. [arXiv:1011.5416v1](#) [math.AG].
- [Sm1] B. D. Smithling, *Topological flatness of orthogonal local models in the split, even case. I*, Math. Ann. **350** (2011), no. 2, 381–416.
- [Sm2] ———, *Admissibility and permissibility for minuscule cocharacters in orthogonal groups*, Manuscripta Math. **136** (2011), no. 3–4, 295–314.
- [Sm3] ———, *Topological flatness of local models for ramified unitary groups. I. The odd dimensional case*, Adv. Math. **226** (2011), no. 4, 3160–3190.
- [St] R. Steinberg, *Endomorphisms of linear algebraic groups*, Memoirs of the American Mathematical Society, No. 80, American Mathematical Society, Providence, R.I., 1968.

- [T] J. Tits, *Reductive groups over local fields*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 29–69.
- [V] A. Vasiu, *Geometry of Shimura varieties of Hodge type over finite fields*, Higher-dimensional geometry over finite fields, NATO Sci. Peace Secur. Ser. D Inf. Commun. Secur., vol. 16, IOS, Amsterdam, 2008, pp. 197–243.
- [Z] X. Zhu, *On the coherence conjecture of Pappas and Rapoport*, preprint. [arXiv:1012.5979v2 \[math.AG\]](#).

UNIVERSITY OF TORONTO, DEPARTMENT OF MATHEMATICS, 40 ST. GEORGE ST., TORONTO,
ON M5S 2E4, CANADA

E-mail address: `bds@math.toronto.edu`